

Two-time Green function method in quantum electrodynamics of high-Z few-electron atoms

V. M. Shabaev

Department of Physics, St. Petersburg State University,
Oulianovskaya Street 1, Petrodvorets, 198504 St. Petersburg,
Russia

The two-time Green function method in quantum electrodynamics of high-Z few-electron atoms is described in detail. This method provides a simple procedure for deriving formulas for the energy shift of a single level and for the energies and wave functions of degenerate and quasi-degenerate states. It also allows one to derive formulas for the transition and scattering amplitudes. Application of the method to resonance scattering processes yields a systematic theory for the spectral line shape. The practical ability of the method is demonstrated by deriving formulas for the QED and interelectronic-interaction corrections to energy levels and transition and scattering amplitudes in one-, two-, and three-electron atoms. Numerical calculations of the Lamb shift, the hyperfine splitting, the bound-electron g factor, and the radiative recombination cross section in heavy ions are also reviewed.

PACS number(s): 12.20.-m, 12.20.Ds, 31.30. Jv

Contents

I	Introduction	4
II	Energy levels of atomic systems	5
A	2 <i>N</i> -time Green function	5
B	Two-time Green function (TTGF) and its analytical properties	8
C	Energy shift of a single level	10
D	Perturbation theory for degenerate and quasidegenerate levels	13
E	Practical calculations	16
1	Zeroth order approximation	16
2	One-electron atom	17
3	Atom with one electron over closed shells	19
4	Two-electron atom	20
5	General rules for practical calculations	21
6	Two-photon exchange diagrams for the ground state of a heliumlike atom	22
7	Self-energy and vacuum-polarization screening diagrams	25
8	Quasidegenerate states	26
F	Nuclear recoil corrections	30
III	Transition probabilities and cross sections of scattering processes	32
A	Photon emission by an atom	33
B	Transition probability in a one-electron atom	36
1	Zeroth order approximation	36
2	QED corrections of first order in α	37
C	Radiative recombination of an electron with an atom	39
D	Radiative recombination of an electron with a high- <i>Z</i> hydrogenlike atom	42
1	Zeroth order approximation	42
2	Interelectronic-interaction corrections of first order in $1/Z$	43
E	Photon scattering by an atom	46
F	Resonance scattering: Spectral line shape	48
G	Resonance photon scattering by a one-electron atom	50
IV	Numerical evaluations of QED and interelectronic-interaction corrections in heavy ions	51
A	Methods of numerical evaluations and renormalization procedure	51
B	Energy levels in heavy ions	52
1	Hydrogenlike ions	52
2	Heliumlike ions	54
3	Lithiumlike ions	54
C	Hyperfine splitting and bound-electron <i>g</i> factor	55
1	Hyperfine splitting in hydrogenlike ions	55
2	Hyperfine splitting in lithiumlike ions	56
3	Bound-electron <i>g</i> factor	57
D	Radiative recombination of an electron with a heavy ion	58
1	QED corrections to the radiative recombination of an electron with a bare nucleus	58
2	Interelectronic-interaction effect on the radiative recombination of an electron with a heavy heliumlike ion	61
3	Resonance recombination of an electron with a heavy hydrogenlike ion	62
V	Conclusion	64
	APPENDIXES	66
A	QED in the Heisenberg representation	66
B	Singularities of the two-time Green function in a finite order of perturbation theory	66
C	Two-time Green function in terms of the Fourier transform of the 2<i>N</i>-time Green function	68

D	Matrix elements of the two-time Green function between one-determinant wave functions	69
E	Double spectral representation for the two-time Green function describing a transition process	69

I. INTRODUCTION

A great progress in experimental investigations of high-Z few-electron systems (see, e.g., [1,2]) stimulated theorists to perform accurate calculations for these systems in the framework of quantum electrodynamics (QED). The calculations of QED and interelectronic-interaction corrections in high-Z few-electron systems are conveniently divided into two stages. The first stage consists in deriving formal expressions for these corrections from the first principles of QED. The second comprises numerical evaluations of these expressions. The present paper is mainly focused on the first stage. As to the numerical calculations, we give only a short overview of them in this paper. For more details we refer to [3–7].

Historically, the first method suitable for deriving the formal expressions for the energy shift of a bound-state level was formulated by Gell-Mann, Low, and Sucher [8,9]. This method is based on introducing an adiabatically damped factor, $\exp(-\lambda|t|)$, in the interaction Hamiltonian and expressing the energy shift in terms of so-called adiabatic S_λ matrix elements. Due to its simple formulation, the Gell-Mann–Low–Sucher formula for the energy shift gained wide spreading in the literature related to high-Z few-electron systems [10–17]. However, the practical use of this method showed that it has several serious drawbacks. One of them is the very complicated derivation of the formal expressions for so-called *reducible* diagrams. By “reducible diagrams” we denote those diagrams where an intermediate-state energy of the atom coincides with the reference-state energy. (This terminology is quite natural since it can be considered as an extension of the definitions introduced by Dyson [18] and by Bethe and Salpeter [19] to high-Z few-electron atoms.) As to *irreducible* diagrams, i.e. those diagrams where the intermediate-state energies differ from the reference-state energy, the derivation of the formal expressions can easily be reduced to the “usual ($\lambda = 0$) S -matrix” elements in each method, including the Gell-Mann–Low–Sucher method as well (see, e.g., [12,16]). Another serious drawback of the Gell-Mann–Low–Sucher method is the fact that this method requires special investigation of the renormalization procedure since the adiabatic S_λ -matrix suffers from ultraviolet divergences. The adiabatically damped factor, $\exp(-\lambda|t|)$, is non-covariant and, therefore, the ultraviolet divergences can not be removed from S_λ if $\lambda \neq 0$. However, from the physical point of view one may expect these divergences to cancel each other in the expression for the energy shift. Therefore, they may be disregarded in the calculation of the energy shift for a single level. For the case of degenerate levels, however, this problem remains since we can not expect that the standard renormalization procedure makes the secular operator finite in the ultraviolet limit [11,13]. In addition, we should note that at present there is no formalism based on the Gell-Mann–Low–Sucher approach which would provide a proper treatment of quasidegenerate levels. Also no formalism in the framework of this approach was developed for calculation of the transition or scattering amplitudes up to now. The same difficulties emerge in the evolution operator method developed in Refs. [20–24]. Attempts to solve some of these problems by modifying the Gell-Mann–Low–Sucher method were recently undertaken in [25–27].

Another way to formulate a perturbation theory for high-Z few-electron systems consists in using Green functions. These functions contain the complete information about the energy levels and the transition and scattering amplitudes. In this way, the renormalization problem does not appear, since Green functions can be renormalized from the very beginning (see, e.g., [28]). Up to now, various versions of the Green function formalism were developed which differ from each other by the methods of extracting the physical information from Green functions, i.e. the energy levels and the transition and scattering amplitudes. One of these methods was worked out in [29–33]. It was successfully employed in many practical calculations [34–46]. Since one of the key elements of this method consists in using two-time Green functions, in what follows we will call it the two-time Green function (TTGF) method. This method, which provides a solution of all the problems appearing in the other formalisms indicated above, will be considered in detail in the present work.

As to other versions of the Green function method [13,17,47–55], a detailed discussion of them would be beyond the scope of the present paper. We note only that some of these methods are also based on employing two-time Green functions but yield other forms of perturbation theory. In Refs. [13,48–51], the two-time Green functions were used for constructing quasipotential equations for high-Z few-electron systems. This corresponds to the perturbation theory in the Brillouin-Wigner form while the method of Refs. [29–33] yields the perturbation theory in the Rayleigh-Schrödinger form. Various versions of the Bethe-Salpeter equation derived from the $2N$ -time Green function formalism for high-Z few-electron systems can be found in [13,52]. In [53,54] the perturbation theory in the Rayleigh-Schrödinger form is constructed for the case of a one-electron system where the problem of relative time coordinates for the electrons does not occur.

The relativistic unit system ($\hbar = c = 1$) and the Heaviside charge unit ($\alpha = \frac{e^2}{4\pi}, e < 0$) are used in the paper.

II. ENERGY LEVELS OF ATOMIC SYSTEMS

In this section we formulate the perturbation theory for the calculation of the energy levels in high- Z few-electron atoms. In these systems the number of electrons denoted by N is much smaller than the nuclear charge number Z . It follows that the interaction of the electrons with each other and with the quantized electromagnetic field is much smaller (by factors $1/Z$ and α , respectively) than the interaction of the electrons with the Coulomb field of the nucleus. Therefore, it is natural to assume that in zeroth approximation the electrons interact only with the Coulomb field of the nucleus and obey the Dirac equation

$$(-i\boldsymbol{\alpha} \cdot \boldsymbol{\nabla} + \beta m + V_C(\mathbf{x}))\psi_n(\mathbf{x}) = \varepsilon_n \psi_n(\mathbf{x}). \quad (2.1)$$

The interaction of the electrons with each other and with the quantized electromagnetic field is accounted for by perturbation theory. In this way we obtain quantum electrodynamics in the Furry picture. It should be noted that we could start also with the Dirac equation with an effective potential $V_{\text{eff}}(\mathbf{x})$ which approximately describes the interaction with the other electrons. In this case the interaction with the potential $\delta V(\mathbf{x}) = V_C(\mathbf{x}) - V_{\text{eff}}(\mathbf{x})$ must be accounted for perturbatively. Using the effective potential provides an extension of the theory to many-electron atoms where, for instance, a local version of the Hartree-Fock potential can be used as $V_{\text{eff}}(\mathbf{x})$. However, for simplicity, in what follows we will assume that in zeroth approximation the electrons interact only with the Coulomb field of the nucleus.

In the present paper we will mainly consider the perturbation theory with the standard QED vacuum. The transition to the formalism in which closed shells are regarded as belonging to the vacuum is realized by replacing $i0$ with $-i0$ in the electron propagator denominators corresponding to the closed shells.

Before formulating the perturbation theory for calculations of the interelectronic interaction and radiative corrections to the energy levels, we consider standard equations of the Green function approach in quantum electrodynamics.

A. $2N$ -time Green function

It can be shown that the complete information about the energy levels of an N -electron atom is contained in the Green function defined as

$$G(x'_1, \dots, x'_N; x_1, \dots, x_N) = \langle 0 | T \psi(x'_1) \cdots \psi(x'_N) \bar{\psi}(x_N) \cdots \bar{\psi}(x_1) | 0 \rangle, \quad (2.2)$$

where $\psi(x)$ is the electron-positron field operator in the Heisenberg representation, $\bar{\psi}(x) = \psi^\dagger \gamma^0$, and T is the time-ordered product operator. The basic equations of quantum electrodynamics in the Heisenberg representation are summarized in Appendix A. Equation (2.2) presents a standard definition of the $2N$ -time Green function which is a fundamental object of quantum electrodynamics. It can be shown (see, e.g., [28,56]) that in the interaction representation the Green function is given by

$$\begin{aligned} G(x'_1, \dots, x'_N; x_1, \dots, x_N) &= \frac{\langle 0 | T \psi_{\text{in}}(x'_1) \cdots \psi_{\text{in}}(x'_N) \bar{\psi}_{\text{in}}(x_N) \cdots \bar{\psi}_{\text{in}}(x_1) \exp \left\{ -i \int d^4 z \mathcal{H}_I(z) \right\} | 0 \rangle}{\langle 0 | T \exp \left\{ -i \int d^4 z \mathcal{H}_I(z) \right\} | 0 \rangle} \end{aligned} \quad (2.3)$$

$$\begin{aligned} &= \left\{ \sum_{m=0}^{\infty} \frac{(-i)^m}{m!} \int d^4 y_1 \cdots d^4 y_m \langle 0 | T \psi_{\text{in}}(x'_1) \cdots \psi_{\text{in}}(x'_N) \bar{\psi}_{\text{in}}(x_N) \cdots \bar{\psi}_{\text{in}}(x_1) \right. \\ &\quad \left. \times \mathcal{H}_I(y_1) \cdots \mathcal{H}_I(y_m) | 0 \rangle \right\} \left\{ \sum_{l=0}^{\infty} \frac{(-i)^l}{l!} \int d^4 z_1 \cdots d^4 z_l \langle 0 | T \mathcal{H}_I(z_1) \cdots \mathcal{H}_I(z_l) | 0 \rangle \right\}^{-1} \end{aligned} \quad (2.4)$$

where

$$\mathcal{H}_I(x) = \frac{e}{2} [\bar{\psi}_{\text{in}}(x) \gamma_\mu, \psi_{\text{in}}(x)] A_{\text{in}}^\mu(x) - \frac{\delta m}{2} [\bar{\psi}_{\text{in}}(x), \psi_{\text{in}}(x)] \quad (2.5)$$

is the interaction Hamiltonian. The commutators in equation (2.5) refer to operators only. The first term in (2.5) describes the interaction of the electron-positron field with the quantized electromagnetic field and the second one is the mass renormalization counterterm. We consider here that the interaction of the electrons with the Coulomb field of the nucleus is included in the unperturbed Hamiltonian, i.e. the Furry picture. However, there is also an alternative method to get the Furry picture. In that method the interaction with the Coulomb field of the nucleus is

included in the interaction Hamiltonian and the Furry picture is obtained by summing infinite sequences of Feynman diagrams describing the interaction of the electrons with the Coulomb potential. As a result of this summation, the free-electron propagators are replaced by bound-electron propagators. This method is very convenient for studying processes involving continuum-electron states. It will be used in the section concerning the radiative recombination process.

The Green function G is constructed by perturbation theory according to equation (2.4). This is carried out with the aid of the Wick theorem (see, e.g., [28]). According to this theorem the time-ordered product of field operators is equal to the sum of normal-ordered products with all possible contractions between the operators

$$T(ABCD \dots) = N(ABCD \dots) + N(A^a B^a C D \dots) + N(A^a B C^a D \dots) + \text{all possible contractions}, \quad (2.6)$$

where N is the normal-ordered product operator and the superscripts denote the contraction between the corresponding operators. The contraction between neighbouring operators is defined by

$$A^a B^a = T(AB) - N(AB) = \langle 0 | T(AB) | 0 \rangle. \quad (2.7)$$

If the contracted operators are boson operators, they can be put one next to another. If the contracted operators are fermion operators, they also can be put one next to another but in this case the expression must be multiplied with the parity of the permutation of the fermion operators. Since in the Green function the vacuum expectation value is calculated, only the term with all operators contracted remains on the right-hand side of equation (2.6). In contrast to the free-electron QED, in the Furry picture the time-ordered product of two fermion operators must be defined also for the equal-time case to obtain the correct vacuum polarization terms. As was noticed in [57], the definition

$$T[A(t)B(t)] = \frac{1}{2}A(t)B(t) - \frac{1}{2}B(t)A(t) \quad (2.8)$$

provides the following simple rule for dealing with the interaction operator. It can be written as

$$\mathcal{H}_I(x) = e \bar{\psi}_{\text{in}}(x) \gamma_\mu \psi_{\text{in}}(x) A_{\text{in}}^\mu(x) - \delta m \bar{\psi}_{\text{in}}(x) \psi_{\text{in}}(x) \quad (2.9)$$

and then the Wick theorem is applied with contractions between all operators, including equal-time operators. We note that the problem of the definition of the time-ordered product of fermion operators at equal times does not appear at all if the alternative method for obtaining the Furry picture discussed above is employed.

The contractions between the electron-positron fields and between the photon fields lead to the following propagators

$$\langle 0 | T \psi_{\text{in}}(x) \bar{\psi}_{\text{in}}(y) | 0 \rangle = \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_n \frac{\psi_n(\mathbf{x}) \bar{\psi}_n(\mathbf{y})}{\omega - \varepsilon_n(1 - i0)} \exp[-i\omega(x^0 - y^0)] \quad (2.10)$$

and

$$\langle 0 | T A_{\text{in}}^\mu(x) A_{\text{in}}^\nu(y) | 0 \rangle = -ig^{\mu\nu} \int \frac{d^4 k}{(2\pi)^4} \frac{\exp[-ik \cdot (x - y)]}{k^2 + i0}. \quad (2.11)$$

Here the Feynman gauge is considered. In equation (2.10) the index n runs over all bound and continuum states.

The denominator in equation (2.3) describes unobservable vacuum-vacuum transitions and, as can be shown (see, e.g., [28]), it cancels disconnected vacuum-vacuum subdiagrams from the numerator. Therefore, we can simply omit all diagrams containing disconnected vacuum-vacuum subdiagrams in the numerator and replace the denominator by 1.

In practical calculations of the Green function it is convenient to work with the Fourier transform with respect to time variables,

$$\begin{aligned} G((p_1^0, \mathbf{x}'_1), \dots, (p_N^0, \mathbf{x}'_N); (p_1^0, \mathbf{x}_1), \dots, (p_N^0, \mathbf{x}_N)) \\ = (2\pi)^{-2N} \int_{-\infty}^{\infty} dx_1^0 \dots dx_N^0 dx_1'^0 \dots dx_N'^0 \\ \times \exp(ip_1'^0 x_1'^0 + \dots + ip_N'^0 x_N'^0 - ip_1^0 x_1^0 - \dots - ip_N^0 x_N^0) \\ \times G(x'_1, \dots, x'_N; x_1, \dots, x_N). \end{aligned} \quad (2.12)$$

For the Green function $G((p_1^0, \mathbf{x}'_1), \dots, (p_N^0, \mathbf{x}'_N); (p_1^0, \mathbf{x}_1), \dots, (p_N^0, \mathbf{x}_N))$, the following Feynman rules can be derived:

(1) External electron line

$$\overline{\text{x}} \longleftarrow \text{y} \quad \frac{i}{2\pi} S(\omega, \text{x}, \text{y}),$$

where

$$S(\omega, \text{x}, \text{y}) = \sum_n \frac{\psi_n(\text{x}) \bar{\psi}_n(\text{y})}{\omega - \varepsilon_n(1 - i0)}, \quad (2.13)$$

$\psi_n(\text{x})$ are solutions of the Dirac equation (2.1).

(2) Internal electron line

$$\overline{\text{x}} \longleftarrow \text{y} \quad \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \quad S(\omega, \text{x}, \text{y}).$$

(3) Disconnected electron line

$$\overline{\text{x}} \longleftarrow \text{y} \quad \frac{i}{2\pi} S(\omega, \text{x}, \text{y}) \delta(\omega - \omega').$$

(4) Internal photon line

$$\text{x} \text{---} \text{y} \quad \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \quad D_{\rho\sigma}(\omega, \text{x} - \text{y}),$$

where, for zero photon mass, $D_{\rho\sigma}(\omega, \text{x} - \text{y})$ is given by

$$D_{\rho\sigma}(\omega, \text{x} - \text{y}) = -g_{\rho\sigma} \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{\exp(i\mathbf{k} \cdot (\text{x} - \text{y}))}{\omega^2 - \mathbf{k}^2 + i0} \quad (2.14)$$


in the Feynman gauge and by

$$D_{00}(\omega, \text{x} - \text{y}) = \frac{1}{4\pi|\text{x} - \text{y}|}, \quad D_{i0} = D_{0i} = 0 \quad (i = 1, 2, 3), \quad (2.15)$$

$$D_{il}(\omega, \text{x} - \text{y}) = \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{\exp(i\mathbf{k} \cdot (\text{x} - \text{y}))}{\omega^2 - \mathbf{k}^2 + i0} \left(\delta_{il} - \frac{k_i k_l}{\mathbf{k}^2} \right) \quad (i, l = 1, 2, 3), \quad (2.16)$$


in the Coulomb gauge. (In this work we assume that the Coulomb gauge is used only for diagrams which do not involve a renormalization procedure. The renormalization in the Coulomb gauge is discussed in Refs. [58,59].)

(5) Vertex




$$-2\pi i e \gamma^\rho \delta(\omega_1 - \omega_2 - \omega_3) \int d\mathbf{x}.$$

(6) The mass counterterm



$$2\pi i \delta(\omega - \omega') \delta m \int d\mathbf{x}.$$

- (7) Symmetry factor $(-1)^P$, where P is the parity of the permutation of the final electron coordinates with respect to the initial ones.
(8) Factor (-1) for every closed electron loop.
(9) If, in addition, an external potential $\delta V(\mathbf{x})$ is present, an additional vertex appears,



$$-2\pi i \gamma^0 \delta(\omega - \omega') \int d\mathbf{x} \delta V(\mathbf{x}).$$

In principle, the Green function G contains the complete information about the energy levels of the atomic system. This can be shown by deriving the spectral representation for G . However, it is a hard task to extract this information directly from G since it depends on $2(N-1)$ relative times (energies) in the time (energy) representation. As we will see in the next section, the two-time Green function defined as

$$\tilde{G}(t', t) \equiv G(t'_1 = t'_2 = \dots t'_N \equiv t'; t_1 = t_2 = \dots t_N \equiv t) \quad (2.17)$$

also contains the complete information about the energy levels, and it is a much simpler task to extract the energy levels from $\tilde{G}$.

B. Two-time Green function (TTGF) and its analytical properties

Let us introduce the Fourier transform of the two-time Green function by

$$\mathcal{G}(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) \delta(E - E') = \frac{1}{2\pi i} \frac{1}{N!} \int_{-\infty}^{\infty} dx^0 dx'^0 \exp(iE'x'^0 - iEx^0) \\ \times \langle 0 | T \psi(x'^0, \mathbf{x}'_1) \dots \psi(x'^0, \mathbf{x}'_N) \bar{\psi}(x^0, \mathbf{x}_N) \dots \bar{\psi}(x^0, \mathbf{x}_1) | 0 \rangle, \quad (2.18)$$

where, as in (2.2), the Heisenberg representation for the electron-positron field operators is used. Defined by equation (2.18) for real E , the Green function $\mathcal{G}$ can be continued analytically to the complex E plane. Analytical properties of this type of Green functions in the complex E plane were studied in various fields of physics (see, e.g., [60–62]). In quantum field theory they were considered in detail by Logunov and Tavkhelidze in [63] (see also [64]), where the two-time Green function was employed for constructing a quasipotential equation. To study the analytical properties of the two-time Green function we derive the spectral representation for $\mathcal{G}$. Using the time-shift transformation rule for the Heisenberg operators (see Appendix A)

$$\psi(x^0, \mathbf{x}) = \exp(iHx^0) \psi(0, \mathbf{x}) \exp(-iHx^0) \quad (2.19)$$

and the equations

$$H|n\rangle = E_n|n\rangle, \quad \sum_n |n\rangle\langle n| = I, \quad (2.20)$$

where H is the Hamiltonian of the system in the Heisenberg representation, we find

$$\mathcal{G}(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) \delta(E - E') \\ = \frac{1}{2\pi i} \frac{1}{N!} \int_{-\infty}^{\infty} dx^0 dx'^0 \exp(iE'x'^0 - iEx^0) \\ \times \left\{ \theta(x'^0 - x^0) \sum_n \exp[i(E_0 - E_n)(x'^0 - x^0)] \langle 0 | \psi(0, \mathbf{x}'_1) \dots \psi(0, \mathbf{x}'_N) | n \rangle \right. \\ \times \langle n | \bar{\psi}(0, \mathbf{x}_N) \dots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle + (-1)^{N^2} \theta(x^0 - x'^0) \sum_n \exp[i(E_0 - E_n)(x^0 - x'^0)] \\ \left. \times \langle 0 | \bar{\psi}(0, \mathbf{x}_N) \dots \bar{\psi}(0, \mathbf{x}_1) | n \rangle \langle n | \psi(0, \mathbf{x}'_1) \dots \psi(0, \mathbf{x}'_N) | 0 \rangle \right\}. \quad (2.21)$$

Assuming, for simplicity, $E_0 = 0$ (it corresponds to choosing the vacuum energy as the origin of reference) and taking into account that

$$\begin{aligned} \int_{-\infty}^{\infty} dx^0 dx'^0 \theta(x'^0 - x^0) \exp[-iE_n(x'^0 - x^0)] \exp[i(E'x'^0 - Ex^0)] \\ = 2\pi\delta(E' - E) \frac{i}{E - E_n + i0}, \end{aligned} \quad (2.22)$$

$$\begin{aligned} \int_{-\infty}^{\infty} dx^0 dx'^0 \theta(x^0 - x'^0) \exp[-iE_n(x^0 - x'^0)] \exp[i(E'x'^0 - Ex^0)] \\ = -2\pi\delta(E' - E) \frac{i}{E + E_n - i0}, \end{aligned} \quad (2.23)$$

we obtain

$$\mathcal{G}(E) = \sum_n \frac{\Phi_n \bar{\Phi}_n}{E - E_n + i0} - (-1)^N \sum_n \frac{\Xi_n \bar{\Xi}_n}{E + E_n - i0}, \quad (2.24)$$

where the variables $\mathbf{x}'_1, \dots, \mathbf{x}'_N, \mathbf{x}_1, \dots, \mathbf{x}_N$ are implicit and

$$\Phi_n(\mathbf{x}_1, \dots, \mathbf{x}_N) = \frac{1}{\sqrt{N!}} \langle 0 | \psi(0, \mathbf{x}_1) \cdots \psi(0, \mathbf{x}_N) | n \rangle, \quad (2.25)$$

$$\Xi_n(\mathbf{x}_1, \dots, \mathbf{x}_N) = \frac{1}{\sqrt{N!}} \langle n | \psi(0, \mathbf{x}_1) \cdots \psi(0, \mathbf{x}_N) | 0 \rangle. \quad (2.26)$$

In equation (2.24) the summation runs over all bound and continuum states of the system of the interacting fields. Let us introduce the functions

$$A(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) = \sum_n \delta(E - E_n) \Phi_n(\mathbf{x}'_1, \dots, \mathbf{x}'_N) \bar{\Phi}_n(\mathbf{x}_1, \dots, \mathbf{x}_N), \quad (2.27)$$

$$B(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) = \sum_n \delta(E - E_n) \Xi_n(\mathbf{x}'_1, \dots, \mathbf{x}'_N) \bar{\Xi}_n(\mathbf{x}_1, \dots, \mathbf{x}_N). \quad (2.28)$$

These functions satisfy the conditions

$$\begin{aligned} \int_{-\infty}^{\infty} dE A(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{1}{N!} \langle 0 | \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) \\ &\quad \times \bar{\psi}(0, \mathbf{x}_N) \cdots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle, \end{aligned} \quad (2.29)$$

$$\begin{aligned} \int_{-\infty}^{\infty} dE B(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{1}{N!} \langle 0 | \bar{\psi}(0, \mathbf{x}_N) \cdots \bar{\psi}(0, \mathbf{x}_1) \\ &\quad \times \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) | 0 \rangle. \end{aligned} \quad (2.30)$$

In terms of these functions, the equation (2.24) is

$$\mathcal{G}(E) = \int_0^{\infty} dE' \frac{A(E')}{E - E' + i0} - (-1)^N \int_0^{\infty} dE' \frac{B(E')}{E + E' - i0}, \quad (2.31)$$

where we have omitted the variables $\mathbf{x}_1, \dots, \mathbf{x}_N, \mathbf{x}'_1, \dots, \mathbf{x}'_N$ and have taken into account that $A(E') = B(E') = 0$ for $E' < 0$ since $E_n \geq 0$. In fact, due to charge conservation, only states with an electric charge of eN contribute to A in the sum over n in the right-hand side of equation (2.27) and only states with an electric charge of $-eN$ contribute to B in the sum over n in the right-hand side of equation (2.28). This can easily be shown by using the following commutation relations

$$[Q, \psi(x)] = -e\psi(x), \quad [Q, \bar{\psi}(x)] = e\bar{\psi}(x), \quad (2.32)$$

where Q is the charge operator in the Heisenberg representation. Therefore, the equation (2.31) can be written as

$$\mathcal{G}(E) = \int_{E_{\min}^{(+)}}^{\infty} dE' \frac{A(E')}{E - E' + i0} - (-1)^N \int_{E_{\min}^{(-)}}^{\infty} dE' \frac{B(E')}{E + E' - i0}, \quad (2.33)$$

where $E_{\min}^{(+)}$ is the minimal energy of states with electric charge eN and $E_{\min}^{(-)}$ is the minimal energy of states with electric charge $-eN$. So far we considered $\mathcal{G}(E)$ for real E . The equation (2.33) shows that the Green function $\mathcal{G}(E)$ is the sum of Cauchy-type integrals. Using the fact that the integrals $\int_{E_{\min}^{(+)}}^{\infty} dE' A(E')$ and $\int_{E_{\min}^{(-)}}^{\infty} dE' B(E')$ converge (see equations (2.29), (2.30)), one can show with the help of standard mathematical methods that the equation

$$\mathcal{G}(E) = \int_{E_{\min}^{(+)}}^{\infty} dE' \frac{A(E')}{E - E'} - (-1)^N \int_{E_{\min}^{(-)}}^{\infty} dE' \frac{B(E')}{E + E'} \quad (2.34)$$

defines an analytical function of E in the complex E plane with the cuts $(-\infty, E_{\min}^{(-)})$ and $[E_{\min}^{(+)}, \infty)$ (see Fig. 1). This equation provides the analytical continuation of the Green function to the complex E plane. According to (2.33), to get the Green function for real E we have to approach the right-hand cut from the upper half-plane and the left-hand cut from the lower half-plane.

In what follows we will be interested in bound states of the system. According to equations (2.24)-(2.34), the bound states correspond to the poles of the function $\mathcal{G}(E)$ on the right-hand real semiaxis. If the interaction between the electron-positron field and the electromagnetic field is switched off, the poles corresponding to bound states are isolated (see Fig. 2). Switching on the interaction between the fields transforms the isolated poles into branch points. This is caused by the fact that due to zero photon mass the bound states are no longer isolated points of the spectrum. Disregarding the instability of excited states, the singularities of the Green function $\mathcal{G}(E)$ are shown in Fig. 3. The poles corresponding to the bound states lie on the upper boundary of the cut starting from the pole corresponding to the ground state. It is natural to assume that $\mathcal{G}(E)$ can be continued analytically under the cut, to the second sheet of the Riemann surface. As a result of this continuation the singularities of $\mathcal{G}(E)$ can be turned down as displayed in Fig. 4.

In fact due to instability of excited states the energies of these states have small imaginary components and, therefore, the related poles lie slightly below the right-hand real semiaxis (Fig. 5). However, in calculations of the energy levels and the transition and scattering amplitudes of non-resonance processes we will neglect the instability of the excited states and, therefore, will assume that the poles lie on the real axis. The imaginary parts of the energies will be taken into account when we will consider the resonance scattering processes.

To formulate the perturbation theory for calculations of the energy levels and the transition and scattering amplitudes we will need to isolate the poles corresponding to the bound states from the related cuts. It can be done by introducing a non-zero photon mass μ which is generally assumed to be larger than the energy shift (or the energy splitting) of the level (levels) under consideration and much smaller than the distance to other levels. The singularities of $\mathcal{G}(E)$ with non-zero photon mass, including one- and two-photon spectra, are shown in Fig. 6. As one can see from this figure, introducing the photon mass makes the poles corresponding to the bound states to be isolated.

In every finite order of perturbation theory the singularities of the Green function $\mathcal{G}(E)$ in the complex E -plane are defined by the unperturbed Hamiltonian. In quantum mechanics this fact easily follows from the expansion of the Green function $(E - H)^{-1} = (E - H_0 - \delta V)^{-1}$ in powers of the perturbation potential δV

$$(E - H)^{-1} = \sum_{n=0}^{\infty} (E - H_0)^{-1} [\delta V (E - H_0)^{-1}]^n. \quad (2.35)$$

As one can see from this equation, to n -th order of perturbation theory the Green function has poles of all orders till $n+1$ at the unperturbed positions of the bound state energies. This fact remains also valid in quantum electrodynamics for $\mathcal{G}(E)$ defined above. It can easily be checked for every specific diagram in first and second order in α . A general proof for an arbitrary diagram is given in Appendix B.

C. Energy shift of a single level

In this section we are interested in the energy shift $\Delta E_a = E_a - E_a^{(0)}$ of a single isolated level a of an N -electron atom due to the perturbative interaction. The unperturbed energy $E_a^{(0)}$ is equal to the sum of the one-electron Dirac-Coulomb energies

$$E_a^{(0)} = \varepsilon_{a_1} + \dots + \varepsilon_{a_N}, \quad (2.36)$$

which are defined by the Dirac equation (2.1). In the simplest case the unperturbed wave function $u_a(\mathbf{x}_1, \dots, \mathbf{x}_N)$ is a one-determinant function

$$u_a(\mathbf{x}_1, \dots, \mathbf{x}_N) = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \psi_{Pa_1}(\mathbf{x}_1) \cdots \psi_{Pa_N}(\mathbf{x}_N), \quad (2.37)$$

where ψ_n are the one-electron Dirac wave functions defined by equation (2.1) and P is the permutation operator. In the general case the unperturbed wave function is a linear combination of the one-determinant functions

$$u_a(\mathbf{x}_1, \dots, \mathbf{x}_N) = \sum_b C_a^b \frac{1}{\sqrt{N!}} \sum_P (-1)^P \psi_{Pb_1}(\mathbf{x}_1) \cdots \psi_{Pb_N}(\mathbf{x}_N). \quad (2.38)$$

We introduce the Green function $g_{aa}(E)$ by

$$\begin{aligned} g_{aa}(E) &= \langle u_a | \mathcal{G}(E) \gamma_1^0 \cdots \gamma_N^0 | u_a \rangle \\ &\equiv \int d\mathbf{x}_1 \cdots d\mathbf{x}_N d\mathbf{x}'_1 \cdots d\mathbf{x}'_N u_a^\dagger(\mathbf{x}'_1, \dots, \mathbf{x}'_N) \\ &\quad \times \mathcal{G}(E, \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) \gamma_1^0 \cdots \gamma_N^0 u_a(\mathbf{x}_1, \dots, \mathbf{x}_N). \end{aligned} \quad (2.39)$$

From the spectral representation for $\mathcal{G}(E)$ (see equations (2.24)-(2.34)) we have

$$g_{aa}(E) = \frac{A_a}{E - E_a} + \text{terms that are regular at } E \sim E_a, \quad (2.40)$$

where

$$\begin{aligned} A_a &= \frac{1}{N!} \int d\mathbf{x}_1 \cdots d\mathbf{x}_N d\mathbf{x}'_1 \cdots d\mathbf{x}'_N u_a^\dagger(\mathbf{x}'_1, \dots, \mathbf{x}'_N) \langle 0 | \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) | a \rangle \\ &\quad \times \langle a | \psi^\dagger(0, \mathbf{x}_N) \cdots \psi^\dagger(0, \mathbf{x}_1) | 0 \rangle u_a(\mathbf{x}_1, \dots, \mathbf{x}_N). \end{aligned} \quad (2.41)$$

We assume here that a non-zero photon mass μ is introduced to isolate the pole corresponding to the bound state a from the related cut. We consider that the photon mass is larger than the energy shift under consideration and much smaller than the distance to other levels. To generate the perturbation series for E_a it is convenient to use a contour integral formalism developed first in operator theory by Szökefalvi-Nagy and Kato [65–70]. Choosing a contour Γ in the complex E plane in a way that it surrounds the pole corresponding to the level a and keeps outside all other singularities (see Fig. 7), we have

$$\frac{1}{2\pi i} \oint_\Gamma dE E g_{aa}(E) = E_a A_a, \quad (2.42)$$

$$\frac{1}{2\pi i} \oint_\Gamma dE g_{aa}(E) = A_a. \quad (2.43)$$

Here we have assumed that the contour Γ is oriented anticlockwise. Dividing the equation (2.42) by (2.43), we obtain

$$E_a = \frac{\frac{1}{2\pi i} \oint_\Gamma dE E g_{aa}(E)}{\frac{1}{2\pi i} \oint_\Gamma dE g_{aa}(E)} \quad (2.44)$$

It is convenient to transform the equation (2.44) to a form that directly yields the energy shift $\Delta E_a = E_a - E_a^{(0)}$. In zeroth order, substituting the operators

$$\psi_{\text{in}}(0, \mathbf{x}) = \sum_{\varepsilon_n > 0} b_n \psi_n(\mathbf{x}) + \sum_{\varepsilon_n < 0} d_n^\dagger \psi_n(\mathbf{x}), \quad (2.45)$$

$$\bar{\psi}_{\text{in}}(0, \mathbf{x}) = \sum_{\varepsilon_n > 0} b_n^\dagger \bar{\psi}_n(\mathbf{x}) + \sum_{\varepsilon_n < 0} d_n \bar{\psi}_n(\mathbf{x}) \quad (2.46)$$

into equations (2.25) and (2.26) instead of $\psi(0, \mathbf{x})$ and $\bar{\psi}(0, \mathbf{x})$, respectively, and considering the states $|n\rangle$ in (2.25) and (2.26) as unperturbed states in the Fock space, from equations (2.24)-(2.26) and (2.39) we find

$$g_{aa}^{(0)} = \frac{1}{E - E_a^{(0)}}. \quad (2.47)$$

This equation can also be derived using the Feynman rules for G (see subsection II(E1)). Denoting $\Delta g_{aa} = g_{aa} - g_{aa}^{(0)}$, from (2.44) we obtain the desired formula [29]

$$\Delta E_a = \frac{\frac{1}{2\pi i} \oint_{\Gamma} dE (E - E_a^{(0)}) \Delta g_{aa}(E)}{1 + \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta g_{aa}(E)}. \quad (2.48)$$

The Green function $\Delta g_{aa}(E)$ is constructed by perturbation theory

$$\Delta g_{aa}(E) = \Delta g_{aa}^{(1)}(E) + \Delta g_{aa}^{(2)}(E) + \dots, \quad (2.49)$$

where the superscript denotes the order in α . If we represent the energy shift as a series in α

$$\Delta E_a = \Delta E_a^{(1)} + \Delta E_a^{(2)} + \dots, \quad (2.50)$$

the formula (2.48) yields

$$\Delta E_a^{(1)} = \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{aa}^{(1)}(E), \quad (2.51)$$

$$\Delta E_a^{(2)} = \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{aa}^{(2)}(E) - \left(\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{aa}^{(1)}(E) \right) \left(\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta g_{aa}^{(1)}(E) \right), \quad (2.52)$$

where $\Delta E \equiv E - E_a^{(0)}$.

Deriving equations (2.44) and (2.48) we have assumed that a non-zero photon mass μ is introduced. This allows taking all the cuts outside the contour Γ as well as regularizing the infrared singularities of individual diagrams. In the Feynman gauge, the photon propagator with non-zero photon mass μ is

$$D_{\rho\sigma}(\omega, \mathbf{x} - \mathbf{y}) = -g_{\rho\sigma} \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{\exp(i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y}))}{\omega^2 - \mathbf{k}^2 - \mu^2 + i0} \quad (2.53)$$

or, after integration,

$$D_{\rho\sigma}(\omega, \mathbf{x} - \mathbf{y}) = g_{\rho\sigma} \frac{\exp(i\sqrt{\omega^2 - \mu^2 + i0} |\mathbf{x} - \mathbf{y}|)}{4\pi |\mathbf{x} - \mathbf{y}|}, \quad (2.54)$$

where $\text{Im}\sqrt{\omega^2 - \mu^2 + i0} > 0$. $D_{\rho\sigma}(\omega, \mathbf{x} - \mathbf{y})$ is an analytical function of ω in the complex ω plane with cuts beginning at the points $\omega = -\mu + i0$ and $\omega = \mu - i0$ (Fig. 8). The related expressions for the photon propagator with non-zero photon mass in other covariant gauges are presented, e.g., in [28].

As was noted in the previous subsection, the singularities of the two-time Green function in the complex E plane are defined by the unperturbed Hamiltonian if it is constructed by perturbation theory. In particular, it means that in n -th order of perturbation theory $g_{aa}(E)$ has poles of all orders till $n + 1$ at the position of the unperturbed energy level under consideration. Therefore, in calculations by perturbation theory it is sufficient to consider the photon mass as a very small parameter which provides a separation of the pole from the related cut. At the end of the calculations after taking into account a whole gauge invariant set of Feynman diagrams we can put $\mu \rightarrow 0$. The possibility of taking the limit $\mu \rightarrow 0$ follows, in particular, from the fact that the contour Γ can be shrunk continuously to the point $E = E_a^{(0)}$ (see Fig. 7).

Generally speaking, the energy shift of an excited level derived by formula (2.48) contains an imaginary component which is caused by its instability. This component defines the width of the spectral line in the Lorentz approximation (see sections III(F,G) for details).

For practical calculations it is convenient to express the Green function $g_{aa}(E)$ in terms of the Fourier transform of the $2N$ -time Green function defined by equation (2.12). By using the identity

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} dx \exp(i\omega x) = \delta(\omega) \quad (2.55)$$

one easily finds (see Appendix C)

$$\begin{aligned}
g_{aa}(E)\delta(E-E') &= \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \\
&\quad \times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\
&\quad \times \langle u_a | G(p_1^0, \dots, p_N^0; p_1'^0, \dots, p_N'^0) \gamma_1^0 \cdots \gamma_N^0 | u_a \rangle,
\end{aligned} \tag{2.56}$$

where

$$\begin{aligned}
&\langle u_a | G(p_1^0, \dots, p_N^0; p_1'^0, \dots, p_N'^0) \gamma_1^0 \cdots \gamma_N^0 | u_a \rangle \\
&\equiv \int d\mathbf{x}_1 \cdots d\mathbf{x}_N d\mathbf{x}'_1 \cdots d\mathbf{x}'_N u_a(\mathbf{x}'_1, \dots, \mathbf{x}'_N) \\
&\quad \times G((p_1^0, \mathbf{x}'_1), \dots, (p_N^0, \mathbf{x}'_N); (p_1^0, \mathbf{x}_1), \dots, (p_N^0, \mathbf{x}_N)) \\
&\quad \times \gamma_1^0 \cdots \gamma_N^0 u_a(\mathbf{x}_1, \dots, \mathbf{x}_N).
\end{aligned} \tag{2.57}$$

According to equation (2.38) the calculation of the matrix elements in (2.56) is reduced to the calculation of the matrix elements between the one-determinant wave functions

$$u_i = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \psi_{P i_1}(\mathbf{x}_1) \cdots \psi_{P i_N}(\mathbf{x}_N), \tag{2.58}$$

$$u_k = \frac{1}{\sqrt{N!}} \sum_P (-1)^P \psi_{P k_1}(\mathbf{x}_1) \cdots \psi_{P k_N}(\mathbf{x}_N). \tag{2.59}$$

To simplify the summation procedure over the permutations in (2.56) which arise from the wave functions as well as from the Green function $G(p_1^0, \dots, p_N^0; p_1'^0, \dots, p_N'^0)$, it is convenient to transform equation (2.56) in the following way. Denoting $\overline{G} = G\gamma_1^0 \cdots \gamma_N^0$, we can write

$$\begin{aligned}
&\overline{G}((p_1^0, \xi'_1), \dots, (p_N^0, \xi'_N); (p_1^0, \xi_1), \dots, (p_N^0, \xi_N)) \\
&= \sum_P (-1)^P \widehat{G}((p_{P1}^0, \xi'_{P1}), \dots, (p_{PN}^0, \xi'_{PN}); (p_1^0, \xi_1), \dots, (p_N^0, \xi_N)),
\end{aligned} \tag{2.60}$$

where $\xi \equiv (\mathbf{x}, \alpha)$ and α is the bispinor index ($\alpha = 1, 2, 3, 4$). Substituting (2.60) in (2.56) and using the symmetry of $\widehat{G}$ with respect to the electron coordinates, for $g_{ik}(E) = \langle u_i | \widehat{G}(E) \gamma_1^0 \cdots \gamma_N^0 | u_k \rangle$ one can obtain (see Appendix D)

$$\begin{aligned}
g_{ik}(E)\delta(E-E') &= \frac{2\pi}{i} \sum_P (-1)^P \psi_{P i_1}^*(\xi'_1) \cdots \psi_{P i_N}^*(\xi'_N) \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \\
&\quad \times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\
&\quad \times \widehat{G}((p_1^0, \xi'_1), \dots, (p_N^0, \xi'_N); (p_1^0, \xi_1), \dots, (p_N^0, \xi_N)) \\
&\quad \times \psi_{k_1}(\xi_1) \cdots \psi_{k_N}(\xi_N),
\end{aligned} \tag{2.61}$$

where repeated variables $\{\xi\}$ imply integration (the integration over $\mathbf{x}$ and the summation over α). In practical calculations by perturbation theory, the formula (2.61) must be only employed for symmetric sets of Feynman diagrams since the symmetry property was used in its derivation.

D. Perturbation theory for degenerate and quasidegenerate levels

In this section we are interested in the atomic levels with energies $E_1, \dots, E_s$ arising from unperturbed degenerate or quasidegenerate levels with energies $E_1^{(0)}, \dots, E_s^{(0)}$. As usual, we assume that the energy shifts of the levels under consideration or their splitting caused by the interaction are much smaller than the distance to other levels. The unperturbed eigenstates form an s -dimensional subspace Ω . We denote the projector on Ω by

$$P^{(0)} = \sum_{k=1}^s P_k^{(0)} = \sum_{k=1}^s u_k u_k^\dagger, \tag{2.62}$$

where $\{u_k\}_{k=1}^s$ are the unperturbed wave functions which, in a general case, are linear combinations of one-determinant functions (see equation (2.38)). We project the Green function $\mathcal{G}(E)$ on the subspace Ω

$$g(E) = P^{(0)} \mathcal{G}(E) \gamma_1^0 \dots \gamma_N^0 P^{(0)}, \quad (2.63)$$

where, as in (2.39), the integration over the electron coordinates is implicit. As in the case of a single level, to isolate the poles of $g(E)$ corresponding to the bound states under consideration, we introduce a non-zero photon mass μ . We assume that the photon mass μ is larger than the energy distance between the levels under consideration and much smaller than the distance to other levels. In this case we can choose a contour Γ in the complex E plane in a way that it surrounds all the poles corresponding to the considered states ($E_1, \dots, E_s$) and keeps outside all other singularities, including the cuts starting from the lower-lying bound states (see Fig. 9). In addition, if we neglect the instability of the states under consideration, the spectral representation (see equations (2.24)-(2.34)) gives

$$g(E) = \sum_{k=1}^s \frac{\varphi_k \varphi_k^\dagger}{E - E_k} + \text{terms that are regular inside of } \Gamma, \quad (2.64)$$

where

$$\varphi_k = P^{(0)} \Phi_k, \quad \varphi_k^\dagger = \Phi_k^\dagger P^{(0)}. \quad (2.65)$$

As in the case of a single level, in zeroth approximation one easily finds

$$g^{(0)}(E) = \sum_{k=1}^s \frac{P_k^{(0)}}{E - E_k^{(0)}}. \quad (2.66)$$

We introduce the operators K and P by

$$K \equiv \frac{1}{2\pi i} \oint_{\Gamma} dE E g(E), \quad (2.67)$$

$$P \equiv \frac{1}{2\pi i} \oint_{\Gamma} dE g(E). \quad (2.68)$$

Using equation (2.64), we obtain

$$K = \sum_{i=1}^s E_i \varphi_i \varphi_i^\dagger, \quad (2.69)$$

$$P = \sum_{i=1}^s \varphi_i \varphi_i^\dagger. \quad (2.70)$$

We note here that, generally speaking, the operator P is not a projector (in particular, $P^2 \neq P$). If the perturbation goes to zero, the vectors $\{\varphi_i\}_{i=1}^s$ approach the correct linearly independent combinations of the vectors $\{u_i\}_{i=1}^s$. Therefore, it is natural to assume that the vectors $\{\varphi_i\}_{i=1}^s$ are also linearly independent. It follows that one can find such vectors $\{v_i\}_{i=1}^s$ that

$$\varphi_i^\dagger v_k = \delta_{ik}. \quad (2.71)$$

Indeed, let

$$\varphi_i = \sum_{j=1}^s a_{ij} u_j, \quad v_k = \sum_{l=1}^s x_{kl} u_l. \quad (2.72)$$

The biorthogonality condition (2.71) gives

$$\sum_{j=1}^s a_{ij} x_{kj} = \delta_{ik}. \quad (2.73)$$

Since the determinant of the matrix $\{a_{ij}\}$ is nonvanishing due to the linear independence of $\{\varphi_i\}_{i=1}^s$, the system (2.73) has a unique solution for any fixed $k = 1, \dots, s$. From (2.69)-(2.71) we have

$$Pv_k = \sum_{i=1}^s \varphi_i \varphi_i^\dagger v_k = \varphi_k, \quad (2.74)$$

$$Kv_k = \sum_{i=1}^s E_i \varphi_i \varphi_i^\dagger v_k = E_k \varphi_k. \quad (2.75)$$

Hence we obtain the equation for v_k , E_k [29]

$$Kv_k = E_k Pv_k. \quad (2.76)$$

According to (2.71) the vectors v_k are normalized by the condition

$$v_{k'}^\dagger Pv_k = \delta_{k'k}. \quad (2.77)$$

The solvability of equation (2.76) yields an equation for the atomic energy levels

$$\det(K - EP) = 0. \quad (2.78)$$

The generalized eigenvalue problem (2.76) with the normalization condition (2.77) can be transformed by the substitution $\psi_k = P^{\frac{1}{2}} v_k$ to the ordinary eigenvalue problem ("Schrödinger-like equation") [33]

$$H\psi_k = E_k \psi_k \quad (2.79)$$

with the ordinary normalization condition

$$\psi_k^\dagger \psi_{k'} = \delta_{kk'}, \quad (2.80)$$

where $H \equiv P^{-\frac{1}{2}}(K)P^{-\frac{1}{2}}$.

The energy levels are determined from the equation

$$\det(H - E) = 0. \quad (2.81)$$

Generally speaking, the energies determined by this equation contain imaginary components which are due to the instability of excited states. In the case when the imaginary components are much smaller than the energy distance between the levels (or the levels have different quantum numbers), they define the widths of the spectral lines in the Lorentz approximation. In the opposite case, when the imaginary components are comparable with the energy distance between the levels which have the same quantum numbers, the spectral line shape depends on the process of the formation of the states under consideration even in the resonance approximation (see sections III(F,G) for details). In what follows, calculating the energy levels we neglect the instability of excited states and assume $H \equiv (H + H^\dagger)/2$ in equations (2.79), (2.81).

The operators K and P are constructed by the formulas (2.67) and (2.68) using perturbation theory

$$K = K^{(0)} + K^{(1)} + K^{(2)} + \dots, \quad (2.82)$$

$$P = P^{(0)} + P^{(1)} + P^{(2)} + \dots, \quad (2.83)$$

where the superscript denotes the order in α . The operator H is

$$H = H^{(0)} + H^{(1)} + H^{(2)} + \dots, \quad (2.84)$$

where

$$H^{(0)} = K^{(0)}, \quad (2.85)$$

$$H^{(1)} = K^{(1)} - \frac{1}{2}P^{(1)}K^{(0)} - \frac{1}{2}K^{(0)}P^{(1)}, \quad (2.86)$$

$$\begin{aligned} H^{(2)} = & K^{(2)} - \frac{1}{2}P^{(2)}K^{(0)} - \frac{1}{2}K^{(0)}P^{(2)} \\ & - \frac{1}{2}P^{(1)}K^{(1)} - \frac{1}{2}K^{(1)}P^{(1)} \\ & + \frac{3}{8}P^{(1)}P^{(1)}K^{(0)} + \frac{3}{8}K^{(0)}P^{(1)}P^{(1)} \\ & + \frac{1}{4}P^{(1)}K^{(0)}P^{(1)}. \end{aligned} \quad (2.87)$$

It is evident that in zeroth order

$$K_{ik}^{(0)} = E_i^{(0)} \delta_{ik}, \quad (2.88)$$

$$P_{ik}^{(0)} = \delta_{ik}, \quad (2.89)$$

$$H_{ik}^{(0)} = E_i^{(0)} \delta_{ik}. \quad (2.90)$$

To derive the equations (2.76)-(2.79) we have introduced a non-zero photon mass μ which was assumed to be larger than the energy distance between the levels under consideration and much smaller than the distance to other levels. At the end of the calculations after taking into account a whole gauge invariant set of Feynman diagrams, we can put $\mu \rightarrow 0$. The possibility of taking this limit in the case of quasidegenerate states follows from the fact that the cuts can be drawn to the related poles by a deformation of the contour Γ as shown in Fig. 10.

As in the case of a single level, for practical calculations we express the Green function $g(E)$ in terms of the Fourier transform of the $2N$ -time Green function

$$\begin{aligned} g(E) \delta(E - E') &= \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \dots dp_N^0 dp_1'^0 \dots dp_N'^0 \delta(E - p_1^0 \dots - p_N^0) \delta(E' - p_1'^0 \dots - p_N'^0) \\ &\quad \times P^{(0)} G(p_1'^0, \dots, p_N'^0; p_1^0, \dots, p_N^0) \gamma_1^0 \dots \gamma_N^0 P^{(0)}, \end{aligned} \quad (2.91)$$

where $G(p_1^0, \dots, p_N^0; p_1^0, \dots, p_N^0)$ is defined by equation (2.12).

E. Practical calculations

In this section we consider some practical applications of the method in the lowest orders of perturbation theory. In what follows we will use the notation

$$I(\omega) = e^2 \alpha^\rho \alpha^\sigma D_{\rho\sigma}(\omega), \quad (2.92)$$

where $\alpha^\rho \equiv \gamma^0 \gamma^\rho = (1, \boldsymbol{\alpha})$. In addition we will employ the following symmetry properties of the photon propagator

$$I(\omega) = I(-\omega), \quad I'(\omega) = -I'(-\omega), \quad (2.93)$$

which, in particular, are valid in the Feynman and Coulomb gauges. Here $I'(\omega) \equiv dI(\omega)/d\omega$.

1. Zeroth order approximation

Let us derive first the formula (2.47) using the Feynman rules for G . According to the equations (2.56)-(2.61) we have

$$\begin{aligned} g_{aa}(E) \delta(E - E') &= \frac{2\pi}{i} \sum_P (-1)^P \int d\mathbf{x}_1 \dots d\mathbf{x}_N d\mathbf{x}_1' \dots d\mathbf{x}_N' \\ &\quad \times \psi_{Pa_1}^\dagger(\mathbf{x}_1') \dots \psi_{Pa_N}^\dagger(\mathbf{x}_N') \int_{-\infty}^{\infty} dp_1^0 \dots dp_N^0 dp_1'^0 \dots dp_N'^0 \\ &\quad \times \delta(E - p_1^0 - \dots - p_N^0) \delta(E' - p_1'^0 - \dots - p_N'^0) \\ &\quad \times \frac{i}{2\pi} S_1(p_1^0, \mathbf{x}_1', \mathbf{x}_1) \delta(p_1'^0 - p_1^0) \dots \frac{i}{2\pi} S_N(p_N^0, \mathbf{x}_N', \mathbf{x}_N) \delta(p_N'^0 - p_N^0) \\ &\quad \times \gamma_1^0 \dots \gamma_N^0 \psi_{a_1}(\mathbf{x}_1) \dots \psi_{a_N}(\mathbf{x}_N) \end{aligned} \quad (2.94)$$

$$\begin{aligned} &= \frac{2\pi}{i} \int_{-\infty}^{\infty} dp_1^0 \dots dp_N^0 dp_1'^0 \dots dp_N'^0 \\ &\quad \times \delta(E - p_1^0 - \dots - p_N^0) \delta(E' - p_1'^0 - \dots - p_N'^0) \\ &\quad \times \frac{i}{2\pi} \frac{1}{p_1^0 - \varepsilon_{a_1} + i0} \delta(p_1'^0 - p_1^0) \dots \frac{i}{2\pi} \frac{1}{p_N^0 - \varepsilon_{a_N} + i0} \delta(p_N'^0 - p_N^0). \end{aligned} \quad (2.95)$$

Integrating over the energies one easily obtains equation (2.47).

2. One-electron atom

Formal expressions for the energy shift in the case of a one-electron atom (or in the case of an atom with one electron over closed shells) can be derived by various methods. In particular, the Dyson-Schwinger equation can be employed for such a derivation. Therefore, the one-electron system is not the best example to demonstrate the advantages of the method under consideration. However, we start with a detailed description of this simple case since it may serve as the simplest introduction to the technique.

Let us consider first a diagram describing the interaction of a one-electron atom with an external potential $\delta V(\mathbf{x})$ to first order in $\delta V(\mathbf{x})$ (Fig. 11). According to equation (2.56) we have

$$\begin{aligned}\Delta g_{aa}^{(1)}(E)\delta(E-E') &= \frac{2\pi}{i} \int d\mathbf{x}' d\mathbf{z} d\mathbf{x} \psi_a^\dagger(\mathbf{x}') \frac{i}{2\pi} \sum_{n_1} \frac{\psi_{n_1}(\mathbf{x}') \bar{\psi}_{n_1}(\mathbf{z})}{E' - \varepsilon_{n_1}(1-i0)} \\ &\quad \times \frac{2\pi}{i} \gamma^0 \delta V(\mathbf{z}) \delta(E' - E) \frac{i}{2\pi} \sum_{n_2} \frac{\psi_{n_2}(\mathbf{z}) \bar{\psi}_{n_2}(\mathbf{x})}{E - \varepsilon_{n_2}(1-i0)} \gamma^0 \psi_a(\mathbf{x}) \\ &= \langle a | \sum_{n_1} \frac{|n_1\rangle\langle n_1|}{E - \varepsilon_{n_1}(1-i0)} \delta V \sum_{n_2} \frac{|n_2\rangle\langle n_2|}{E - \varepsilon_{n_2}(1-i0)} |a\rangle \delta(E' - E) \\ &= \frac{1}{(E - \varepsilon_a)^2} \langle a | \delta V | a \rangle \delta(E' - E).\end{aligned}\tag{2.96}$$

Substituting (2.96) into (2.51), we obtain

$$\begin{aligned}\Delta E_a^{(1)} &= \frac{1}{2\pi i} \oint_{\Gamma} dE (E - \varepsilon_a) \Delta g_{aa}^{(1)}(E) \\ &= \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{\langle a | \delta V | a \rangle}{E - \varepsilon_a} = \langle a | \delta V | a \rangle.\end{aligned}\tag{2.97}$$

To first order in α , the QED corrections are defined by the self energy (SE) and vacuum polarization (VP) diagrams which are shown in Figs. 12 and 13, respectively. Consider first the SE diagram. We find

$$\begin{aligned}\Delta g_{aa}^{(1)}(E)\delta(E-E') &= \frac{2\pi}{i} \int d\mathbf{x}' d\mathbf{y} d\mathbf{z} d\mathbf{x} \psi_a^\dagger(\mathbf{x}') \frac{i}{2\pi} \sum_{n_1} \frac{\psi_{n_1}(\mathbf{x}') \bar{\psi}_{n_1}(\mathbf{y})}{E' - \varepsilon_{n_1}(1-i0)} \frac{i}{2\pi} \int_{-\infty}^{\infty} dp^0 \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \\ &\quad \times e\gamma^\rho \frac{2\pi}{i} \delta(E' - p^0 - \omega) \sum_n \frac{\psi_n(\mathbf{y}) \bar{\psi}_n(\mathbf{z})}{p^0 - \varepsilon_n(1-i0)} D_{\rho\sigma}(\omega, \mathbf{y} - \mathbf{z}) \\ &\quad \times e\gamma^\sigma \frac{2\pi}{i} \delta(p^0 + \omega - E) \frac{i}{2\pi} \sum_{n_2} \frac{\psi_{n_2}(\mathbf{z}) \bar{\psi}_{n_2}(\mathbf{x})}{E - \varepsilon_{n_2}(1-i0)} \gamma^0 \psi_a(\mathbf{x}) \\ &= \frac{1}{(E - \varepsilon_a)^2} e^2 \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \int d\mathbf{y} d\mathbf{z} \psi_a^\dagger(\mathbf{y}) \alpha^\rho \sum_n \frac{\psi_n(\mathbf{y}) \bar{\psi}_n(\mathbf{z})}{E - \omega - \varepsilon_n(1-i0)} \\ &\quad \times D_{\rho\sigma}(\omega, \mathbf{y} - \mathbf{z}) \alpha^\sigma \psi_a(\mathbf{z}) \delta(E' - E),\end{aligned}\tag{2.98}$$

where $\alpha^\rho \equiv \gamma^0 \gamma^\rho = (1, \boldsymbol{\alpha})$. Denoting

$$\langle a | \Sigma(E) | b \rangle = \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_n \frac{\langle an | e^2 \alpha^\rho \alpha^\sigma D_{\rho\sigma}(\omega) | nb \rangle}{E - \omega - \varepsilon_n(1-i0)},\tag{2.99}$$

we get

$$\Delta g_{aa}^{(1)}(E) = \frac{\langle a | \Sigma(E) | a \rangle}{(E - \varepsilon_a)^2}.\tag{2.100}$$

Substituting (2.100) into (2.51), we obtain

$$\begin{aligned}\Delta E_a^{(1)} &= \frac{1}{2\pi i} \oint_{\Gamma} dE (E - \varepsilon_a) \Delta g_{aa}^{(1)}(E) \\ &= \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{\langle a | \Sigma(E) | a \rangle}{E - \varepsilon_a} = \langle a | \Sigma(\varepsilon_a) | a \rangle.\end{aligned}\tag{2.101}$$

Here we have taken into account the fact that, for a non-zero photon mass μ , $\Delta g(E)$ has isolated poles at $E = \varepsilon_a$ in every order of perturbation theory. In the final expression one can put $\mu \rightarrow 0$.

The expression (2.101) suffers from an ultraviolet divergence and has to be considered together with a counterterm diagram (Fig. 14). Taking into account the counterterm results in a replacement

$$\langle a | \Sigma(\varepsilon_a) | a \rangle \rightarrow \langle a | \Sigma_R(\varepsilon_a) | a \rangle = \langle a | (\Sigma(\varepsilon_a) - \gamma^0 \delta m) | a \rangle. \quad (2.102)$$

The corresponding calculation for the VP diagram (Fig. 13) yields

$$\begin{aligned} \Delta g_{aa}^{(1)}(E) \delta(E - E') &= -\frac{2\pi}{i} \int d\mathbf{x} d\mathbf{y} \bar{\psi}_a(\mathbf{x}) \frac{i}{2\pi} \frac{1}{E' - \varepsilon_a} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega' \\ &\quad \times e \gamma^\rho \frac{2\pi}{i} \delta(E' + \omega' - E) D_{\rho\sigma}(\omega', \mathbf{x} - \mathbf{y}) e \frac{2\pi}{i} \delta(\omega') \\ &\quad \times \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \text{Tr} \left[\sum_n \frac{\psi_n(\mathbf{y}) \bar{\psi}_n(\mathbf{y})}{\omega - \varepsilon_n (1 - i0)} \gamma^\sigma \right] \frac{i}{2\pi} \frac{1}{E - \varepsilon_a} \psi_a(\mathbf{x}). \end{aligned} \quad (2.103)$$

Introducing the VP potential by

$$U_{\text{VP}}(\mathbf{x}) = \frac{e^2}{2\pi i} \int d\mathbf{y} \alpha^\rho D_{\rho\sigma}(0, \mathbf{x} - \mathbf{y}) \int_{-\infty}^{\infty} d\omega \text{Tr} \left[\sum_n \frac{\psi_n(\mathbf{y}) \psi_n^\dagger(\mathbf{y})}{\omega - \varepsilon_n (1 - i0)} \alpha^\sigma \right], \quad (2.104)$$

we find

$$\Delta g_{aa}^{(1)}(E) = \frac{\langle a | U_{\text{VP}} | a \rangle}{(E - \varepsilon_a)^2} \quad (2.105)$$

and, therefore,

$$\Delta E_a^{(1)} = \langle a | U_{\text{VP}} | a \rangle. \quad (2.106)$$

In reality, due to a spherical symmetry of the Coulomb potential of the nucleus, only zeroth components of the α matrices contribute to U_{VP} ,

$$U_{\text{VP}}(\mathbf{x}) = \frac{\alpha}{2\pi i} \int d\mathbf{y} \frac{1}{|\mathbf{x} - \mathbf{y}|} \int_{-\infty}^{\infty} d\omega \text{Tr} [G_C(\omega, \mathbf{y}, \mathbf{y})], \quad (2.107)$$

where

$$G_C(\omega, \mathbf{x}, \mathbf{y}) = \sum_n \frac{\psi_n(\mathbf{x}) \psi_n^\dagger(\mathbf{y})}{\omega - \varepsilon_n (1 - i0)} \quad (2.108)$$

is the Dirac-Coulomb Green function. The expression (2.107) is ultraviolet divergent. The charge renormalization makes it finite.

Let us now consider the combined δV -SE corrections described by the Feynman diagrams presented in Fig. 15. For the diagrams "a" and "b" one easily finds

$$\begin{aligned} \Delta g_{aa}^{(2,a+b)}(E) &= \frac{1}{(E - \varepsilon_a)^2} \sum_n \langle a | \delta V | n \rangle \frac{1}{E - \varepsilon_n} \langle n | \Sigma(E) | a \rangle \\ &\quad + \frac{1}{(E - \varepsilon_a)^2} \sum_n \langle a | \Sigma(E) | n \rangle \frac{1}{E - \varepsilon_n} \langle n | \delta V | a \rangle. \end{aligned} \quad (2.109)$$

This contribution is conveniently divided into two parts: *irreducible* ($\varepsilon_n \neq \varepsilon_a$) and *reducible* ($\varepsilon_n = \varepsilon_a$). For the irreducible part one obtains

$$\begin{aligned} \frac{1}{2\pi i} \oint_\Gamma dE (E - \varepsilon_a) \Delta g_{aa}^{(2,a+b,\text{irr})}(E) &= \frac{1}{2\pi i} \oint_\Gamma dE \frac{1}{E - \varepsilon_a} \sum_n^{(n \neq a)} \left[\frac{\langle a | \delta V | n \rangle \langle n | \Sigma(E) | a \rangle}{E - \varepsilon_n} \right. \\ &\quad \left. + \frac{\langle a | \Sigma(E) | n \rangle \langle n | \delta V | a \rangle}{E - \varepsilon_n} \right] \\ &= \sum_n^{(n \neq a)} \left[\frac{\langle a | \delta V | n \rangle \langle n | \Sigma(\varepsilon_a) | a \rangle}{\varepsilon_a - \varepsilon_n} \right. \\ &\quad \left. + \frac{\langle a | \Sigma(\varepsilon_a) | n \rangle \langle n | \delta V | a \rangle}{\varepsilon_a - \varepsilon_n} \right] \end{aligned} \quad (2.110)$$

Here we have taken into account that, due to the spherical symmetry of the Coulomb potential, matrix elements $\langle a|\Sigma(\varepsilon_a)|b\rangle$ are equal to zero if $\varepsilon_a = \varepsilon_b$ and $a \neq b$. The reducible part is

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Gamma} dE (E - \varepsilon_a) \Delta g_{aa}^{(2,a+b,\text{red})}(E) &= \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{1}{(E - \varepsilon_a)^2} \left[\langle a|\delta V|a\rangle \langle a|\Sigma(E)|a\rangle \right. \\ &\quad \left. + \langle a|\Sigma(E)|a\rangle \langle a|\delta V|a\rangle \right] \\ &= 2\langle a|\delta V|a\rangle \langle a|\Sigma'(\varepsilon_a)|a\rangle, \end{aligned} \quad (2.111)$$

where $\Sigma'(\varepsilon_a) \equiv (d\Sigma(E)/dE)_{E=\varepsilon_a}$. The reducible contribution should be considered together with the related contribution from the second term in equation (2.52). Taking into account that

$$\frac{1}{2\pi i} \oint_{\Gamma} dE (E - \varepsilon_a) \Delta g_{aa}^{(1,\delta V)}(E) = \langle a|\delta V|a\rangle \quad (2.112)$$

and

$$\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta g_{aa}^{(1,\text{SE})}(E) = \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{\langle a|\Sigma(E)|a\rangle}{(E - \varepsilon_a)^2} = \langle a|\Sigma'(\varepsilon_a)|a\rangle \quad (2.113)$$

we obtain

$$- \left(\frac{1}{2\pi i} \oint_{\Gamma} dE (E - \varepsilon_a) \Delta g_{aa}^{(1,\delta V)}(E) \right) \left(\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta g_{aa}^{(1,\text{SE})}(E) \right) = -\langle a|\delta V|a\rangle \langle a|\Sigma'(\varepsilon_a)|a\rangle. \quad (2.114)$$

For the total contribution of the diagrams "a" and "b" we find

$$\begin{aligned} \Delta E_a^{(2,a+b)} &= \sum_n^{(n \neq a)} \left[\frac{\langle a|\delta V|n\rangle \langle n|\Sigma(\varepsilon_a)|a\rangle}{\varepsilon_a - \varepsilon_n} + \frac{\langle a|\Sigma(\varepsilon_a)|n\rangle \langle n|\delta V|a\rangle}{\varepsilon_a - \varepsilon_n} \right] \\ &\quad + \langle a|\delta V|a\rangle \langle a|\Sigma'(\varepsilon_a)|a\rangle. \end{aligned} \quad (2.115)$$

For the vertex contribution (the diagram "c") we obtain

$$\begin{aligned} \Delta g_{aa}^{(2,c)}(E) &= \frac{1}{(E - \varepsilon_a)^2} e^2 \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \int d\mathbf{x} d\mathbf{y} d\mathbf{z} \psi_a^\dagger(\mathbf{y}) \alpha^\rho \sum_{n_1} \frac{\psi_{n_1}(\mathbf{y}) \psi_{n_1}^\dagger(\mathbf{x})}{E - \omega - \varepsilon_{n_1}(1 - i0)} \delta V(\mathbf{x}) \\ &\quad \times \sum_{n_2} \frac{\psi_{n_2}(\mathbf{x}) \psi_{n_2}^\dagger(\mathbf{z})}{E - \omega - \varepsilon_{n_2}(1 - i0)} D_{\rho\sigma}(\omega, \mathbf{y} - \mathbf{z}) \alpha^\sigma \psi_a(\mathbf{z}). \end{aligned} \quad (2.116)$$

This diagram is irreducible. A simple evaluation yields

$$\begin{aligned} \Delta E_a^{(2,c)} &= e^2 \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \int d\mathbf{x} d\mathbf{y} d\mathbf{z} \psi_a^\dagger(\mathbf{y}) \alpha^\rho G_C(\varepsilon_a - \omega, \mathbf{y}, \mathbf{x}) \delta V(\mathbf{x}) \\ &\quad \times G_C(\varepsilon_a - \omega, \mathbf{x}, \mathbf{z}) D_{\rho\sigma}(\omega, \mathbf{y} - \mathbf{z}) \alpha^\sigma \psi_a(\mathbf{z}). \end{aligned} \quad (2.117)$$

The related mass-counterterm diagrams (Fig. 16) are accounted for by the replacement $\Sigma \rightarrow \Sigma_R = \Sigma - \gamma^0 \delta m$ in equation (2.115). This replacement makes the irreducible contribution in equation (2.115) to be finite. As to the reducible contribution, its ultraviolet and infrared divergences are cancelled by the corresponding divergences of the vertex contribution given by equation (2.117).

3. Atom with one electron over closed shells

The consideration given above can easily be adopted to the case of an atom with one electron over closed shells by regarding the closed shells as belonging to a redefined vacuum. The redefinition of the vacuum results in replacing $i0$ by $-i0$ in the electron propagator denominators corresponding to the closed shells. In other words, it means replacing the standard Feynman contour of integration over the electron energy C by a new contour C' (Fig. 17). In this formalism the one-electron radiative corrections are incorporated together with the interelectronic-interaction corrections and the total energy of the closed shells is considered as the origin of reference. The difference of the

integrals along C' and C is an integral along the contour C_{int} . It describes the interaction of the valence electron with the closed-shell electrons. Therefore, to find the interelectronic-interaction corrections we have to replace the contour C in the expressions for the one-electron radiative corrections by the contour C_{int} . For example, in the case of one electron over the $(1s)^2$ shell in a lithiumlike ion, the first order interelectronic-interaction corrections are obtained from the formulas for the SE and VP corrections derived above by the replacement

$$\sum_n \frac{|n\rangle\langle n|}{\omega - \varepsilon_n(1-i0)} \rightarrow -\frac{2\pi}{i} \delta(\omega - \varepsilon_{1s}) \sum_c^{(\varepsilon_c = \varepsilon_{1s})} |c\rangle\langle c|. \quad (2.118)$$

As a result of this replacement, one obtains for the interelectronic-interaction correction

$$\Delta E_a^{(1, \text{int})} = \sum_c^{(\varepsilon_c = \varepsilon_{1s})} [\langle ac|I(0)|ac\rangle - \langle ac|I(\varepsilon_a - \varepsilon_{1s})|ca\rangle], \quad (2.119)$$

where $I(\omega)$ is defined by equation (2.92). In Ref. [38] this formalism was employed to derive formal expressions for the interelectronic-interaction corrections to the hyperfine splitting in lithiumlike ions (see section IV(C2)).

4. Two-electron atom

Let us consider now the energy shift of a single level (n) in a two-electron atom. To first order in α , in addition to the one-electron SE and VP contributions (Figs. 18,19) we have to consider the one-photon exchange diagram (Fig. 20). The derivation of the energy shift due to the SE and VP diagrams is easily reduced to the case of a one-electron atom by a simple integration over the energy variable of a disconnected electron propagator. Therefore, below we discuss only the one-photon exchange diagram.

For simplicity, we assume that the unperturbed wave function of the state under consideration is the one-determinant function

$$u_n(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{\sqrt{2}} \sum_P (-1)^P \psi_{Pa}(\mathbf{x}_1) \psi_{Pb}(\mathbf{x}_2). \quad (2.120)$$

The transition to the general case of a many-determinant function (2.38) causes no problem and can be done in the final expression for the energy shift.

According to equation (2.56), for the one-photon exchange diagram we have

$$\begin{aligned} \Delta g_{nn}^{(1)} &= \left(\frac{i}{2\pi}\right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \sum_P (-1)^P \frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \\ &\times \frac{1}{p_1^0 - \varepsilon_a + i0} \frac{1}{E - p_1^0 - \varepsilon_b + i0} \langle PaPb|I(p_1'^0 - p_1^0)|ab\rangle. \end{aligned} \quad (2.121)$$

Formula (2.51) gives

$$\begin{aligned} \Delta E_n^{(1)} &= \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \left(\frac{i}{2\pi}\right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \sum_P (-1)^P \frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} \\ &\times \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \frac{1}{p_1^0 - \varepsilon_a + i0} \frac{1}{E - p_1^0 - \varepsilon_b + i0} \\ &\times \langle PaPb|I(p_1'^0 - p_1^0)|ab\rangle, \end{aligned} \quad (2.122)$$

where, as in (2.51), $\Delta E \equiv E - E_n^{(0)}$. Transforming

$$\frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} = \frac{1}{\Delta E} \left(\frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} + \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \right), \quad (2.123)$$

$$\frac{1}{p_1^0 - \varepsilon_a + i0} \frac{1}{E - p_1^0 - \varepsilon_b + i0} = \frac{1}{\Delta E} \left(\frac{1}{p_1^0 - \varepsilon_a + i0} + \frac{1}{E - p_1^0 - \varepsilon_b + i0} \right), \quad (2.124)$$

we obtain

$$\Delta E_n^{(1)} = \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{1}{\Delta E} \left\{ \left(\frac{i}{2\pi} \right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \sum_P (-1)^P \left(\frac{1}{p_1^0 - \varepsilon_{Pa} + i0} + \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \right) \right. \\ \left. \times \left(\frac{1}{p_1^0 - \varepsilon_a + i0} + \frac{1}{E - p_1^0 - \varepsilon_b + i0} \right) \langle PaPb | I(p_1'^0 - p_1^0) | ab \rangle \right\}. \quad (2.125)$$

The expression in the curly braces of (2.125) is a regular function of E inside the contour Γ , if the photon mass μ is chosen as indicated above. A direct way to check this fact consists in integrating over p_1^0 and $p_1'^0$ by using the apparent expression for the photon propagator given by equation (2.53). It can also be understood by observing that the integrand in this expression is the sum of terms which contain singularities from the electron propagators in p_1^0 ($p_1'^0$) only above or only below the real axis (for real E). Therefore, in each term we can vary E in the complex E plane within the contour Γ , keeping the same order of bypassing the singularities in the p_1^0 ($p_1'^0$) integration by moving slightly the contour of the p_1^0 ($p_1'^0$) integration into the complex plane. (In contrast to that, the contour of the p_1^0 ($p_1'^0$) integration in equation (2.122) is sandwiched between two poles and, therefore, can not be moved to the complex plane.) The branch points of the photon propagators are moved outside the contour Γ due to the non-zero photon mass. However, instead of investigating the analytical properties for every specific diagram, it is more convenient to use the general analytical properties of the Green function $g_{nn}(E)$ in the complex E plane in every order of perturbation theory (see the related discussion above and Appendix B). According to these properties, the function $\Delta g_{nn}^{(1)}(E)$ can not have poles at the point $E = E_n^{(0)}$ of order higher than 2 and, therefore, the expression in the curly braces of (2.125) is a regular function of E at this point. So, we have to calculate the first order residue at the point $E = E_n^{(0)}$. We also stress that we do not need any apparent form for the analytical continuation of the expression in the curly braces to the complex E plane since we calculate it only for real E , at the point $E = E_n^{(0)}$, where the present expression is valid. Calculating the E residue we obtain

$$\Delta E_n^{(1)} = \left(\frac{i}{2\pi} \right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \sum_P (-1)^P \left(\frac{1}{p_1^0 - \varepsilon_{Pa} + i0} + \frac{1}{-(p_1'^0 - \varepsilon_{Pa}) + i0} \right) \\ \times \left(\frac{1}{p_1^0 - \varepsilon_a + i0} + \frac{1}{-(p_1^0 - \varepsilon_a) + i0} \right) \langle PaPb | I(p_1'^0 - p_1^0) | ab \rangle. \quad (2.126)$$

Taking into account the identity

$$\frac{i}{2\pi} \left(\frac{1}{x + i0} + \frac{1}{-x + i0} \right) = \delta(x) \quad (2.127)$$

we find

$$\Delta E_n^{(1)} = \sum_P (-1)^P \langle PaPb | I(\varepsilon_{Pa} - \varepsilon_a) | ab \rangle. \quad (2.128)$$

5. General rules for practical calculations

Let us consider now some general remarks to the derivation given above. In order to perform first the integration over E we have separated the singularity in ΔE by employing the identities (2.123) and (2.124). Another way could consist in transforming the left-hand sides of equations (2.123) and (2.124) by the identity

$$\frac{1}{p^0 - \varepsilon_a + i0} \frac{1}{E - p^0 - \varepsilon_b + i0} = \frac{2\pi}{i} \delta(p^0 - \varepsilon_a) \frac{1}{\Delta E} \\ + \frac{1}{p^0 - \varepsilon_a - i0} \frac{1}{E - p^0 - \varepsilon_b + i0}, \quad (2.129)$$

where we have used equation (2.127). Using this identity allows one to separate contributions singular in $1/\Delta E$ from non-singular ones. The singular contributions result only from the first term in the right-hand side of equation (2.129). It can easily be understood by observing that the second term has both singularities in p^0 above the real axis (for real E). It follows that the contour of the p^0 integration in the expression for the energy shift can be moved slightly into the complex E plane keeping the same order of bypassing the singularities. It means that we can vary E in the complex E plane within the contour Γ and, therefore, the integrand is a regular function of E within this contour. Identities like (2.129) are very useful in calculations of three- and more electron atoms (see section IV(B3)).

We also want to note that in all cases the order of the singularity in $1/\Delta E$ is quite evident from the type of the diagram under consideration. If the diagram is irreducible, the factors $1/\Delta E$ may come only from the initial and final propagators. In this case the second term in the right-hand side of equation (2.129) does not contribute to the energy shift and, therefore, the derivation of the formal expression for the energy shift becomes trivial. For reducible diagrams the factors $1/\Delta E$ arise also from internal electron propagators.

We can formulate the following simple rule for deriving the energy shift from a certain diagram. Using identities like (2.123), (2.124) or (2.129), we separate all singularities in $1/\Delta E$ and then integrate over E assuming that the rest is a regular function of E within the contour Γ . As is discussed above, the order of the singularity is quite evident for every specific diagram and it is a simple task to separate the factor $1/\Delta E$ to the right power. However, even if we separate this factor to a power which is larger or smaller than the real order of the singularity, it is impossible to miss the correct result. In the first case (the power is larger than the real order of the singularity) the result of the calculation remains the same as in the case when we separate the factor $1/\Delta E$ to the right power. In the second case (the power is smaller than the real order of the singularity) we obtain an infinite result ($\sim 1/0$). It means that we should increase the power of the separated singularity and repeat the calculation until we get a finite result.

6. Two-photon exchange diagrams for the ground state of a heliumlike atom

The two-photon exchange diagrams are presented in Fig. 21. Here we derive the energy shift from these diagrams for the case of the ground state of a heliumlike atom. The case of a single excited state of a two-electron atom is considered in detail in [36]. The wave function of the ground state is given by

$$u_1(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{\sqrt{2}} \sum_P (-1)^P \psi_{Pa}(\mathbf{x}_1) \psi_{Pb}(\mathbf{x}_2). \quad (2.130)$$

The unperturbed energy is $E_1^{(0)} = \varepsilon_a + \varepsilon_b$, where $\varepsilon_a = \varepsilon_b$.

Consider first the two-photon ladder diagram (Fig. 21a). For the first term in (2.52) we have

$$\begin{aligned} \Delta E_{\text{lad}}^{(2)} &= \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \sum_P (-1)^P \left(\frac{i}{2\pi}\right)^3 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 d\omega \\ &\times \sum_{n_1 n_2} \langle PaPb | I(p_1'^0 - \omega) | n_1 n_2 \rangle \langle n_1 n_2 | I(\omega - p_1^0) | ab \rangle \\ &\times \frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \frac{1}{\omega - \varepsilon_{n_1}(1 - i0)} \\ &\times \frac{1}{E - \omega - \varepsilon_{n_2}(1 - i0)} \frac{1}{p_1^0 - \varepsilon_a + i0} \frac{1}{E - p_1^0 - \varepsilon_b + i0}. \end{aligned} \quad (2.131)$$

Let us divide this contribution into irreducible ($\varepsilon_{n_1} + \varepsilon_{n_2} \neq \varepsilon_a + \varepsilon_b$) and reducible ($\varepsilon_{n_1} + \varepsilon_{n_2} = \varepsilon_a + \varepsilon_b$) parts

$$\Delta E_{\text{lad}}^{(2)} = \Delta E_{\text{lad}}^{(2, \text{irred})} + \Delta E_{\text{lad}}^{(2, \text{red})}. \quad (2.132)$$

Using the identities (2.123) and (2.124) we obtain for the irreducible part

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{11}^{(2, \text{irred})}(E) &= \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{1}{\Delta E} \left\{ \sum_P (-1)^P \left(\frac{i}{2\pi}\right)^3 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 d\omega \right. \\ &\times \sum_{\substack{n_1, n_2 \\ \varepsilon_{n_1} + \varepsilon_{n_2} \neq \varepsilon_a + \varepsilon_b}} \langle PaPb | I(p_1'^0 - \omega) | n_1 n_2 \rangle \langle n_1 n_2 | I(\omega - p_1^0) | ab \rangle \\ &\times \left(\frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} + \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \right) \\ &\times \frac{1}{\omega - \varepsilon_{n_1}(1 - i0)} \frac{1}{E - \omega - \varepsilon_{n_2}(1 - i0)} \\ &\times \left. \left(\frac{1}{p_1^0 - \varepsilon_a + i0} + \frac{1}{E - p_1^0 - \varepsilon_b + i0} \right) \right\}. \end{aligned} \quad (2.133)$$

The expression in the curly braces of (2.133) is a regular function of E inside the contour Γ if the photon mass μ is chosen as indicated above. (If it were not so, we would get an infinite result; see the related discussion in the previous subsection.) Calculating the E residue we find

$$\begin{aligned} \Delta E_{\text{lad}}^{(2,\text{irred})} &= \sum_P (-1)^P \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_{n_1, n_2}^{\varepsilon_{n_1} + \varepsilon_{n_2} \neq \varepsilon_a + \varepsilon_b} \langle PaPb | I(\varepsilon_{Pa} - \omega) | n_1 n_2 \rangle \\ &\quad \times \langle n_1 n_2 | I(\omega - \varepsilon_a) | ab \rangle \frac{1}{\omega - \varepsilon_{n_1}(1 - i0)} \frac{1}{E_1^{(0)} - \omega - \varepsilon_{n_2}(1 - i0)}. \end{aligned} \quad (2.134)$$

This derivation shows that the energy shift from an irreducible diagram is obtained by evaluation of the "usual S-matrix" element. For the numerical evaluation of (2.134) it is convenient to rotate the contour of the integration in the complex ω plane [71].

For the reducible contribution we have

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{11}^{(2,\text{red})}(E) &= \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{1}{(\Delta E)^2} \left\{ \sum_P (-1)^P \left(\frac{i}{2\pi} \right)^3 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 d\omega \right. \\ &\quad \times \sum_{n_1, n_2}^{\varepsilon_{n_1} + \varepsilon_{n_2} = \varepsilon_a + \varepsilon_b} \langle PaPb | I(p_1'^0 - \omega) | n_1 n_2 \rangle \langle n_1 n_2 | I(\omega - p_1^0) | ab \rangle \\ &\quad \times \left(\frac{1}{p_1'^0 - \varepsilon_{Pa} + i0} + \frac{1}{E - p_1'^0 - \varepsilon_{Pb} + i0} \right) \\ &\quad \times \left(\frac{1}{\omega - \varepsilon_{n_1} + i0} + \frac{1}{E - \omega - \varepsilon_{n_2} + i0} \right) \\ &\quad \left. \times \left(\frac{1}{p_1^0 - \varepsilon_a + i0} + \frac{1}{E - p_1^0 - \varepsilon_b + i0} \right) \right\}. \end{aligned} \quad (2.135)$$

The expression in the curly braces of (2.135) is a regular function within the contour Γ . Calculating the E residue and taking into account that $\varepsilon_a = \varepsilon_b$, we get

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{11}^{(2,\text{red})}(E) &= -\frac{i}{2\pi} \sum_P (-1)^P \sum_{n_1, n_2}^{\varepsilon_{n_1} + \varepsilon_{n_2} = 2\varepsilon_a} \left\{ \int_{-\infty}^{\infty} dp_1'^0 \langle PaPb | I(p_1'^0 - \varepsilon_a) | n_1 n_2 \rangle \right. \\ &\quad \times \langle n_1 n_2 | I(0) | ab \rangle \frac{1}{(\varepsilon_a - p_1'^0 + i0)^2} + \int_{-\infty}^{\infty} dp_1^0 \langle PaPb | I(0) | n_1 n_2 \rangle \\ &\quad \times \langle n_1 n_2 | I(\varepsilon_a - p_1^0) | ab \rangle \frac{1}{(\varepsilon_a - p_1^0 + i0)^2} \\ &\quad + \int_{-\infty}^{\infty} d\omega \langle PaPb | I(\varepsilon_a - \omega) | n_1 n_2 \rangle \\ &\quad \left. \times \langle n_1 n_2 | I(\omega - \varepsilon_a) | ab \rangle \frac{1}{(\varepsilon_a - \omega + i0)^2} \right\}. \end{aligned} \quad (2.136)$$

This contribution should be considered together with the second term in equation (2.52). As was obtained above, the first factor in this term is

$$\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{11}^{(1)}(E) = \sum_P (-1)^P \langle PaPb | I(0) | ab \rangle. \quad (2.137)$$

A simple calculation of the second factor yields

$$\begin{aligned} \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta g_{11}^{(1)}(E) &= \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{1}{(\Delta E)^2} \left\{ \left(\frac{i}{2\pi} \right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \right. \\ &\quad \times \sum_P (-1)^P \left(\frac{1}{p_1'^0 - \varepsilon_a + i0} + \frac{1}{E - p_1'^0 - \varepsilon_a + i0} \right) \\ &\quad \times \left(\frac{1}{p_1^0 - \varepsilon_a + i0} + \frac{1}{E - p_1^0 - \varepsilon_a + i0} \right) \langle PaPb | I(p_1'^0 - p_1^0) | ab \rangle \end{aligned}$$

$$\begin{aligned}
&= -\frac{i}{2\pi} \sum_P (-1)^P \left\{ \int_{-\infty}^{\infty} dp_1^0 \frac{1}{(p_1^0 - \varepsilon_a - i0)^2} \langle PaPb | I(p_1^0 - \varepsilon_a) | ab \rangle \right. \\
&\quad \left. + \int_{-\infty}^{\infty} dp_1^0 \frac{1}{(p_1^0 - \varepsilon_a - i0)^2} \langle PaPb | I(p_1^0 - \varepsilon_a) | ab \rangle \right\}.
\end{aligned} \tag{2.138}$$

For the total reducible contribution we obtain

$$\begin{aligned}
\Delta E_{\text{lad}}^{(2,\text{red})} &= \frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{11}^{(2,\text{red})}(E) - \left(\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta E \Delta g_{11}^{(1)}(E) \right) \left(\frac{1}{2\pi i} \oint_{\Gamma} dE \Delta g_{11}^{(1)}(E) \right) \\
&= - \sum_P (-1)^P \sum_{n_1, n_2}^{\varepsilon_{n_1} + \varepsilon_{n_2} = 2\varepsilon_a} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \langle PaPb | I(\omega - \varepsilon_a) | n_1 n_2 \rangle \\
&\quad \times \langle n_1 n_2 | I(\omega - \varepsilon_a) | ab \rangle \frac{1}{(\omega - \varepsilon_a - i0)^2}.
\end{aligned} \tag{2.139}$$

A similar calculation of the two-photon crossed-ladder diagram (Fig. 21b) gives

$$\begin{aligned}
\Delta E_{\text{cross}}^{(2)} &= \sum_P (-1)^P \sum_{n_1, n_2} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \langle Pan_2 | I(\omega - \varepsilon_a) | n_1 b \rangle \langle n_1 Pb | I(\omega - \varepsilon_a) | an_2 \rangle \\
&\quad \times \frac{1}{\omega - \varepsilon_{n_1}(1 - i0)} \frac{1}{\omega - \varepsilon_{n_2}(1 - i0)}.
\end{aligned} \tag{2.140}$$

The contribution $\Delta E_{\text{lad}}^{(2,\text{red})}$ contains an infrared divergent term which is cancelled by a related term ($\varepsilon_{n_1} = \varepsilon_{n_2} = \varepsilon_a$) from the contribution $\Delta E_{\text{cross}}^{(2)}$. In the individual contributions the infrared singularities are regularized by a non-zero photon mass μ . If the ladder and crossed-ladder contributions are merged by the common ω -integration, the integral is convergent and we can put $\mu \rightarrow 0$ before the integration over ω (see [36] for details). However, to show how the calculation for a non-zero photon mass can be performed, let us calculate the reducible contribution for a finite μ . We have to calculate the integral

$$I_1 = \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \frac{\exp[i\sqrt{\omega^2 - \mu^2 + i0}(r_{12} + r_{34})]}{r_{12}r_{34}} \frac{1}{(\omega - i0)^2}. \tag{2.141}$$

Using the identity

$$\exp[i\sqrt{\omega^2 - \mu^2 + i0} r] = -\frac{2}{\pi} \int_0^{\infty} dk k \frac{\sin(kr)}{(\omega^2 - k^2 - \mu^2 + i0)}, \tag{2.142}$$

we obtain

$$I_1 = -\frac{2}{\pi} \frac{i}{2\pi} \frac{1}{r_{12}r_{34}} \int_{-\infty}^{\infty} d\omega \int_0^{\infty} dk k \frac{\sin(k(r_{12} + r_{34}))}{(\omega^2 - k^2 - \mu^2 + i0)} \frac{1}{(\omega - i0)^2}. \tag{2.143}$$

Decomposing the denominator

$$\omega^2 - k^2 - \mu^2 + i0 = (\omega - \sqrt{k^2 + \mu^2} + i0)(\omega + \sqrt{k^2 + \mu^2} - i0) \tag{2.144}$$

and integrating over ω , we find

$$I_1 = -\frac{1}{\pi} \frac{1}{r_{12}r_{34}} \int_0^{\infty} dk k \frac{\sin(k(r_{12} + r_{34}))}{(k^2 + \mu^2)^{3/2}}. \tag{2.145}$$

According to [72], the last integral is

$$I_1 = -\frac{1}{\pi} \frac{r_{12} + r_{34}}{r_{12}r_{34}} K_0[\mu(r_{12} + r_{34})], \tag{2.146}$$

where

$$K_0(z) = -\log(z/2) \sum_{k=0}^{\infty} \frac{(z/2)^{2k}}{(k!)^2} + \sum_{k=0}^{\infty} \frac{z^{2k}}{2^{2k}(k!)^2} \psi(k+1). \quad (2.147)$$

Considering $\mu \rightarrow 0$ we find

$$\begin{aligned} \Delta E_{\text{ladder}}^{(2,\text{red})} &= -\frac{\alpha^2}{\pi} \sum_P (-1)^P \int d\mathbf{x}_1 \cdots d\mathbf{x}_4 \bar{\psi}_{Pa}(\mathbf{x}_3) \bar{\psi}_{Pb}(\mathbf{x}_4) \gamma_3^\rho \gamma_4^\sigma \\ &\quad \times \sum_{\substack{\varepsilon_{n_1}=\varepsilon_{n_2}=\varepsilon_a \\ n_1, n_2}} \psi_{n_1}(\mathbf{x}_3) \bar{\psi}_{n_1}(\mathbf{x}_1) \psi_{n_2}(\mathbf{x}_4) \bar{\psi}_{n_2}(\mathbf{x}_2) \gamma_1^\lambda \gamma_2^\nu \left(\frac{1}{r_{12}} + \frac{1}{r_{34}} \right) \\ &\quad \times [\log(r_{12} + r_{34}) + \log \mu - \log 2 - \psi(1)] g_{\rho\sigma} g_{\lambda\nu} \psi_a(\mathbf{x}_1) \psi_b(\mathbf{x}_2). \end{aligned} \quad (2.148)$$

The corresponding contribution ($\varepsilon_{n_1} = \varepsilon_{n_2} = \varepsilon_a$) from the crossed-ladder diagram is calculated in the same way. The sum of the reducible contribution from the ladder diagram and the related contribution from the crossed-ladder diagram is [29]

$$\begin{aligned} \Delta E^{(2,\text{infr})} &= -\frac{\alpha^2}{\pi} \sum_P (-1)^P \sum_{\substack{\varepsilon_{n_1}=\varepsilon_{n_2}=\varepsilon_a \\ n_1, n_2}} \int d\mathbf{x}_1 \cdots d\mathbf{x}_4 \bar{\psi}_{Pa}(\mathbf{x}_3) \bar{\psi}_{Pb}(\mathbf{x}_4) \gamma_3^\rho \gamma_4^\sigma \\ &\quad \times \psi_{n_1}(\mathbf{x}_3) \bar{\psi}_{n_1}(\mathbf{x}_1) \psi_{n_2}(\mathbf{x}_4) \bar{\psi}_{n_2}(\mathbf{x}_2) \gamma_1^\lambda \gamma_2^\nu \\ &\quad \times \left[g_{\rho\sigma} g_{\lambda\nu} \left(\frac{1}{r_{12}} + \frac{1}{r_{34}} \right) \log(r_{12} + r_{34}) \right. \\ &\quad \left. - g_{\rho\nu} g_{\sigma\lambda} \left(\frac{1}{r_{14}} + \frac{1}{r_{23}} \right) \log(r_{14} + r_{23}) \right] \psi_a(\mathbf{x}_1) \psi_b(\mathbf{x}_2). \end{aligned} \quad (2.149)$$

The terms containing the factor $\log \mu - \log 2 - \psi(1)$ have cancelled each other. For the numerical evaluation of this contribution, one can use the following representation [36]

$$\begin{aligned} \Delta E^{(2,\text{infr})} &= \frac{1}{\pi} \int_0^\infty \frac{dy}{y} \langle ba|S'(y)|ab \rangle [2\langle ab|S(y)|ab \rangle - \langle aa|S(y)|aa \rangle - \langle bb|S(y)|bb \rangle] \\ &\quad + \frac{1}{\pi} \int_0^\infty \frac{dy}{y} \langle ba|S(y)|ab \rangle [2\langle ab|S'(y)|ab \rangle - \langle aa|S'(y)|aa \rangle - \langle bb|S'(y)|bb \rangle] \\ &\quad - \frac{2}{\pi} \int_0^\infty \frac{dy}{y} \langle ba|S(y)|ab \rangle \langle ba|S'(y)|ab \rangle, \end{aligned} \quad (2.150)$$

where

$$S(y) = \frac{\alpha(1 - \boldsymbol{\alpha}_1 \cdot \boldsymbol{\alpha}_2)}{r_{12}} e^{-yr_{12}}, \quad (2.151)$$

$$S'(y) = -\alpha(1 - \boldsymbol{\alpha}_1 \cdot \boldsymbol{\alpha}_2) e^{-yr_{12}}. \quad (2.152)$$

Other representations of this term can be found in Refs. [71,73].

7. Self-energy and vacuum-polarization screening diagrams

The self-energy and vacuum-polarization screening diagrams are presented in Figs. 22 and 23, respectively. The derivation of the calculation formulas for the energy shift of a single level due to the SE screening diagrams using the TTGF method is described in detail in Ref. [43]. If the unperturbed wave function is given by equation (2.120), the contribution of the SE screening diagrams is written as

$$\begin{aligned} \Delta E &= \Delta E^{\text{irred}} + \Delta E^{\text{red}} + \Delta E^{\text{ver}}, \\ \Delta E^{\text{irred}} &= \sum_P (-1)^P \left\{ \sum_{\substack{\varepsilon_n \neq \varepsilon_a \\ n}} \langle PaPb|I(\Delta)|nb \rangle \frac{1}{\varepsilon_a - \varepsilon_n} \langle n|\Sigma(\varepsilon_a)|a \rangle \right. \\ &\quad \left. + \sum_{\substack{\varepsilon_n \neq \varepsilon_b \\ n}} \langle PaPb|I(\Delta)|an \rangle \frac{1}{\varepsilon_b - \varepsilon_n} \langle n|\Sigma(\varepsilon_b)|b \rangle \right\} \end{aligned} \quad (2.153)$$

$$\begin{aligned}
& + \sum_n^{\varepsilon_n \neq \varepsilon_{Pa}} \langle Pa | \Sigma(\varepsilon_{Pa}) | n \rangle \frac{1}{\varepsilon_{Pa} - \varepsilon_n} \langle nPb | I(\Delta) | ab \rangle \\
& + \sum_n^{\varepsilon_n \neq \varepsilon_{Pb}} \langle Pb | \Sigma(\varepsilon_{Pb}) | n \rangle \frac{1}{\varepsilon_{Pb} - \varepsilon_n} \langle Pan | I(\Delta) | ab \rangle \Big\}, \tag{2.154}
\end{aligned}$$

$$\begin{aligned}
\Delta E^{\text{red}} &= \langle ba | I'(\varepsilon_b - \varepsilon_a) | ab \rangle \Big[\langle a | \Sigma(\varepsilon_a) | a \rangle - \langle b | \Sigma(\varepsilon_b) | b \rangle \Big] \\
& + \sum_P (-1)^P \langle PaPb | I(\Delta) | ab \rangle \Big[\langle a | \Sigma'(\varepsilon_a) | a \rangle + \langle b | \Sigma'(\varepsilon_b) | b \rangle \Big], \tag{2.155}
\end{aligned}$$

$$\begin{aligned}
\Delta E^{\text{ver}} &= \sum_P (-1)^P \sum_{n_1 n_2} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \left[\frac{\langle n_1 Pb | I(\Delta) | n_2 b \rangle \langle Pan_2 | I(\omega) | n_1 a \rangle}{[\varepsilon_{Pa} - \omega - \varepsilon_{n_1}(1-i0)][\varepsilon_a - \omega - \varepsilon_{n_2}(1-i0)]} \right. \\
& \left. + \frac{\langle Pan_1 | I(\Delta) | an_2 \rangle \langle Pbn_2 | I(\omega) | n_1 b \rangle}{[\varepsilon_{Pb} - \omega - \varepsilon_{n_1}(1-i0)][\varepsilon_b - \omega - \varepsilon_{n_2}(1-i0)]} \right], \tag{2.156}
\end{aligned}$$

where $\Delta = \varepsilon_{Pa} - \varepsilon_a$. The corresponding contribution of the VP screening diagrams is [40,44]

$$\Delta E = \Delta E_a^{\text{irred}} + \Delta E_a^{\text{red}} + \Delta E_b, \tag{2.157}$$

$$\begin{aligned}
\Delta E_a^{\text{irred}} &= \sum_P (-1)^P \Big\{ \sum_n^{\varepsilon_n \neq \varepsilon_a} \langle PaPb | I(\Delta) | nb \rangle \frac{1}{\varepsilon_a - \varepsilon_n} \langle n | U_{\text{VP}} | a \rangle \\
& + \sum_n^{\varepsilon_n \neq \varepsilon_b} \langle PaPb | I(\Delta) | an \rangle \frac{1}{\varepsilon_b - \varepsilon_n} \langle n | U_{\text{VP}} | b \rangle \\
& + \sum_n^{\varepsilon_n \neq \varepsilon_{Pa}} \langle Pa | U_{\text{VP}} | n \rangle \frac{1}{\varepsilon_{Pa} - \varepsilon_n} \langle nPb | I(\Delta) | ab \rangle \\
& + \sum_n^{\varepsilon_n \neq \varepsilon_{Pb}} \langle Pb | U_{\text{VP}} | n \rangle \frac{1}{\varepsilon_{Pb} - \varepsilon_n} \langle Pan | I(\Delta) | ab \rangle \Big\}, \tag{2.158}
\end{aligned}$$

$$\Delta E_a^{\text{red}} = \langle ba | I'(\varepsilon_b - \varepsilon_a) | ab \rangle \Big[\langle a | U_{\text{VP}} | a \rangle - \langle b | U_{\text{VP}} | b \rangle \Big], \tag{2.159}$$

$$\Delta E_b = \sum_P (-1)^P \langle PaPb | I_{\text{VP}}(\Delta) | ab \rangle, \tag{2.160}$$

where

$$\begin{aligned}
I_{\text{VP}}(\varepsilon, \mathbf{x}, \mathbf{y}) &= \frac{\alpha^2}{2\pi i} \int_{-\infty}^{\infty} d\omega \int d\mathbf{z}_1 \int d\mathbf{z}_2 \frac{\alpha_{1\mu} \exp(i|\varepsilon||\mathbf{x} - \mathbf{z}_1|)}{|\mathbf{x} - \mathbf{z}_1|} \frac{\alpha_{2\nu} \exp(i|\varepsilon||\mathbf{y} - \mathbf{z}_2|)}{|\mathbf{y} - \mathbf{z}_2|} \\
& \times \text{Tr}[\alpha^\mu G(\omega - \frac{\varepsilon}{2}, \mathbf{z}_1, \mathbf{z}_2) \alpha^\nu G(\omega + \frac{\varepsilon}{2}, \mathbf{z}_2, \mathbf{z}_1)]. \tag{2.161}
\end{aligned}$$

The expressions (2.153)-(2.160) are ultraviolet divergent. The renormalization of these expressions can be performed in the same way as for the first-order SE and VP contributions (see Refs. [39,40,42,43] for details).

8. Quasidegenerate states

Let us now consider some applications of the method to the case of quasidegenerate states of a heliumlike ion. This case arises, for instance, if one is interested in the energies of the $(1s2p_{1/2})_1$ and $(1s2p_{3/2})_1$ states. These states are strongly mixed for low and middle Z and, therefore, must be treated as quasidegenerate. It means that the off-diagonal matrix elements of the energy operator H between these states have to be taken into account. The unperturbed wave functions are written as

$$u_i(\mathbf{x}_1, \mathbf{x}_2) = A_N \sum_{m_{i_1} m_{i_2}} \langle j_{i_1} m_{i_1} j_{i_2} m_{i_2} | JM \rangle \sum_P (-1)^P \psi_{P i_1}(\mathbf{x}_1) \psi_{P i_2}(\mathbf{x}_2), \tag{2.162}$$

where A_N is the normalization factor equal to $1/\sqrt{2}$ for non-equivalent electrons and to $1/2$ for equivalent electrons, J is the total angular momentum, and M is its projection. However, in what follows, to compactify the formulas we will construct the matrix elements of H between the one-determinant wave functions

$$u_i(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{\sqrt{2}} \sum_P (-1)^P \psi_{Pi_1}(\mathbf{x}_1) \psi_{Pi_2}(\mathbf{x}_2). \quad (2.163)$$

The transition to the wave functions defined by equation (2.162) can easily be accomplished in the final formulas.

First we consider the contribution from the one-photon exchange diagram. To derive the formulas for $H_{ik}^{(1)}$ we will assume that $E_i^{(0)} \neq E_k^{(0)}$. However, all the final formulas remain to be valid also for the case $E_i^{(0)} = E_k^{(0)}$ which was considered in detail above. According to the Feynman rules and the definition of $g(E)$, the contribution of the one-photon exchange diagram to $g^{(1)}(E)$ is

$$g_{ik}^{(1)}(E) = \left(\frac{i}{2\pi}\right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \sum_P (-1)^P \frac{1}{p_1'^0 - \varepsilon_{Pi_1} + i0} \frac{1}{E - p_1'^0 - \varepsilon_{Pi_2} + i0} \\ \times \frac{1}{p_1^0 - \varepsilon_{k_1} + i0} \frac{1}{E - p_1^0 - \varepsilon_{k_2} + i0} \langle Pi_1 Pi_2 | I(p_1'^0 - p_1^0) | k_1 k_2 \rangle. \quad (2.164)$$

Using the identities (2.123) and (2.124), we obtain

$$K_{ik}^{(1)} = \frac{1}{2\pi i} \oint_{\Gamma} dE \frac{E}{(E - E_i^{(0)})(E - E_k^{(0)})} \left\{ \left(\frac{i}{2\pi}\right)^2 \int_{-\infty}^{\infty} dp_1^0 dp_1'^0 \sum_P (-1)^P \right. \\ \times \left(\frac{1}{p_1'^0 - \varepsilon_{Pi_1} + i0} + \frac{1}{E - p_1'^0 - \varepsilon_{Pi_2} + i0} \right) \left(\frac{1}{p_1^0 - \varepsilon_{k_1} + i0} + \frac{1}{E - p_1^0 - \varepsilon_{k_2} + i0} \right) \\ \left. \times \langle Pi_1 Pi_2 | I(p_1'^0 - p_1^0) | k_1 k_2 \rangle \right\}. \quad (2.165)$$

The expression in the curly braces of (2.165) is a regular function of E inside the contour Γ , if the photon mass μ is chosen as indicated above. Calculating the E residues and taking into account the identity (2.127), we obtain

$$K_{ik}^{(1)} = \frac{i}{2\pi} \int_{-\infty}^{\infty} dp_1^0 \sum_P (-1)^P \frac{E_i^{(0)} \langle Pi_1 Pi_2 | I(\varepsilon_{Pi_1} - p_1^0) | k_1 k_2 \rangle}{E_i^{(0)} - E_k^{(0)}} \\ \times \left(\frac{1}{p_1^0 - \varepsilon_{k_1} + i0} + \frac{1}{E_i^{(0)} - p_1^0 - \varepsilon_{k_2} + i0} \right) \\ + \frac{i}{2\pi} \int_{-\infty}^{\infty} dp_1'^0 \sum_P (-1)^P \frac{E_k^{(0)} \langle Pi_1 Pi_2 | I(p_1'^0 - \varepsilon_{k_1}) | k_1 k_2 \rangle}{E_k^{(0)} - E_i^{(0)}} \\ \times \left(\frac{1}{p_1'^0 - \varepsilon_{Pi_1} + i0} + \frac{1}{E_k^{(0)} - p_1'^0 - \varepsilon_{Pi_2} + i0} \right). \quad (2.166)$$

In the same way we find

$$P_{ik}^{(1)} = \frac{i}{2\pi} \int_{-\infty}^{\infty} dp_1^0 \sum_P (-1)^P \frac{\langle Pi_1 Pi_2 | I(\varepsilon_{Pi_1} - p_1^0) | k_1 k_2 \rangle}{E_i^{(0)} - E_k^{(0)}} \\ \times \left(\frac{1}{p_1^0 - \varepsilon_{k_1} + i0} + \frac{1}{E_i^{(0)} - p_1^0 - \varepsilon_{k_2} + i0} \right) \\ + \frac{i}{2\pi} \int_{-\infty}^{\infty} dp_1'^0 \sum_P (-1)^P \frac{\langle Pi_1 Pi_2 | I(p_1'^0 - \varepsilon_{k_1}) | k_1 k_2 \rangle}{E_k^{(0)} - E_i^{(0)}} \\ \times \left(\frac{1}{p_1'^0 - \varepsilon_{Pi_1} + i0} + \frac{1}{E_k^{(0)} - p_1'^0 - \varepsilon_{Pi_2} + i0} \right). \quad (2.167)$$

Symmetrizing equations (2.166) and (2.167) with respect to both electrons we transform them to the form

$$\begin{aligned}
K_{ik}^{(1)} = \sum_P (-1)^P \left\{ \frac{1}{2} [\langle Pi_1 Pi_2 | I(\Delta_1) | k_1 k_2 \rangle + \langle Pi_1 Pi_2 | I(\Delta_2) | k_1 k_2 \rangle] \right. \\
- \frac{(E_i^{(0)} + E_k^{(0)})}{2} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \langle Pi_1 Pi_2 | I(\omega) | k_1 k_2 \rangle \\
\left. \times \left[\frac{1}{(\omega + \Delta_1 - i0)(\omega - \Delta_2 - i0)} + \frac{1}{(\omega + \Delta_2 - i0)(\omega - \Delta_1 - i0)} \right] \right\}, \quad (2.168)
\end{aligned}$$

$$\begin{aligned}
P_{ik}^{(1)} = - \sum_P (-1)^P \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \langle Pi_1 Pi_2 | I(\omega) | k_1 k_2 \rangle \\
\times \left[\frac{1}{(\omega + \Delta_1 - i0)(\omega - \Delta_2 - i0)} + \frac{1}{(\omega + \Delta_2 - i0)(\omega - \Delta_1 - i0)} \right], \quad (2.169)
\end{aligned}$$

where $\Delta_1 = \varepsilon_{Pi_1} - \varepsilon_{k_1}$ and $\Delta_2 = \varepsilon_{Pi_2} - \varepsilon_{k_2}$. Substituting (2.168), (2.169) into (2.86), we get [35,74]

$$H_{ik}^{(1)} = \frac{1}{2} \sum_P (-1)^P [\langle Pi_1 Pi_2 | I(\Delta_1) | k_1 k_2 \rangle + \langle Pi_1 Pi_2 | I(\Delta_2) | k_1 k_2 \rangle]. \quad (2.170)$$

Let us now consider the contribution to H from the combined δV - interelectronic interaction diagrams presented in Fig. 24. For simplicity, we will assume that δV is a spherically-symmetric potential. In the case under consideration, the simplest way to derive the formulas for $H_{ik}^{(2)}$ consists in using the fact that these diagrams can be obtained as the first-order (in δV) correction to the one-photon exchange contribution derived above. So, the contribution from these diagrams can be obtained by the following replacements in equation (2.170)

$$|k_1\rangle \rightarrow |k_1\rangle + \delta|k_1\rangle, \quad (2.171)$$

$$|k_2\rangle \rightarrow |k_2\rangle + \delta|k_2\rangle, \quad (2.172)$$

$$|Pi_1\rangle \rightarrow |Pi_1\rangle + \delta|Pi_1\rangle, \quad (2.173)$$

$$|Pi_2\rangle \rightarrow |Pi_2\rangle + \delta|Pi_2\rangle, \quad (2.174)$$

$$I(\varepsilon_a - \varepsilon_b) \rightarrow I(\varepsilon_a + \delta\varepsilon_a - \varepsilon_b - \delta\varepsilon_b), \quad (2.175)$$

where, to first order in δV ,

$$\delta\varepsilon_a = \langle a | \delta V | a \rangle, \quad (2.176)$$

$$\delta|a\rangle = \sum_{n \neq a} \frac{|n\rangle \langle n | \delta V | a \rangle}{\varepsilon_a - \varepsilon_n}. \quad (2.177)$$

Here we have taken into account that, due to the spherical symmetry of δV , $\langle n | \delta V | a \rangle = 0$ if $\varepsilon_n = \varepsilon_a$ and $|n\rangle \neq |a\rangle$. Decomposing the modified expression for the one-photon exchange diagram to the first order in δV , we find that the total correction is the sum of the irreducible and reducible parts,

$$H_{ik}^{(2)} = H_{ik}^{(2,\text{irred})} + H_{ik}^{(2,\text{red})}, \quad (2.178)$$

where

$$\begin{aligned}
H_{ik}^{(2,\text{irred})} = \frac{1}{2} \sum_P (-1)^P [\langle \delta Pi_1 Pi_2 | I(\Delta_1) + I(\Delta_2) | k_1 k_2 \rangle \\
+ \langle Pi_1 \delta Pi_2 | I(\Delta_1) + I(\Delta_2) | k_1 k_2 \rangle \\
+ \langle Pi_1 Pi_2 | I(\Delta_1) + I(\Delta_2) | \delta k_1 k_2 \rangle \\
+ \langle Pi_1 Pi_2 | I(\Delta_1) + I(\Delta_2) | k_1 \delta k_2 \rangle] \quad (2.179)
\end{aligned}$$

and

$$\begin{aligned}
H_{ik}^{(2,\text{red})} = \frac{1}{2} \sum_P (-1)^P \{ [\langle Pi_1 | \delta V | Pi_1 \rangle - \langle k_1 | \delta V | k_1 \rangle] \langle Pi_1 Pi_2 | I'(\Delta_1) | k_1 k_2 \rangle \\
+ [\langle Pi_2 | \delta V | Pi_2 \rangle - \langle k_2 | \delta V | k_2 \rangle] \langle Pi_1 Pi_2 | I'(\Delta_2) | k_1 k_2 \rangle \}. \quad (2.180)
\end{aligned}$$

Equations (2.179) and (2.180) provide the matrix elements between the one-determinant wave functions defined by equation (2.163). To get the matrix elements between the wave functions defined by equation (2.162), we have to multiply these equations by the Clebsch-Gordan coefficients and to sum over the projections of the one-electron angular momenta.

The expression for $H_{ik}^{(2)}$ can also be derived by direct application of the TTGF method. It can easily be performed in the same way as for the one-photon exchange diagram. We note that this derivation yields a formula for $H_{ik}^{(2)}$ which is slightly different from the expression given above. In particular, for the irreducible contribution one finds

$$\begin{aligned} H_{ik}^{(2,\text{irred})} = & \frac{1}{2} \sum_P (-1)^P [\langle \delta P i_1 P i_2 | I(\Delta_1) + I(\Delta_2) | k_1 k_2 \rangle \\ & + \langle P i_1 \delta P i_2 | I(\Delta_1) + I(\Delta_2) | k_1 k_2 \rangle \\ & + \langle P i_1 P i_2 | I(\Delta_1) + I(\Delta_2) | \delta k_1 k_2 \rangle \\ & + \langle P i_1 P i_2 | I(\Delta_1) + I(\Delta_2) | k_1 \delta k_2 \rangle] \\ & + \Delta H_{ik}^{(2,\text{irred})}, \end{aligned} \quad (2.181)$$

where

$$\begin{aligned} \Delta H_{ik}^{(2,\text{irred})} = & \frac{1}{2} (E_i^{(0)} - E_k^{(0)}) \sum_P (-1)^P \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \left\{ \frac{\langle \delta P i_1 P i_2 | I(\omega - \varepsilon_{P i_1}) | k_1 k_2 \rangle}{(\omega - \varepsilon_{k_1} + i0)(E_i^{(0)} - \omega - \varepsilon_{k_2} - i0)} \right. \\ & + \frac{\langle P i_1 \delta P i_2 | I(\omega - \varepsilon_{P i_1}) | k_1 k_2 \rangle}{(\omega - \varepsilon_{k_1} - i0)(E_i^{(0)} - \omega - \varepsilon_{k_2} + i0)} \\ & - \frac{\langle P i_1 P i_2 | I(\omega - \varepsilon_{k_1}) | \delta k_1 k_2 \rangle}{(\omega - \varepsilon_{P i_1} + i0)(E_k^{(0)} - \omega - \varepsilon_{P i_2} - i0)} \\ & - \frac{\langle P i_1 P i_2 | I(\omega - \varepsilon_{k_1}) | k_1 \delta k_2 \rangle}{(\omega - \varepsilon_{P i_1} - i0)(E_k^{(0)} - \omega - \varepsilon_{P i_2} + i0)} \\ & + \sum_{n \neq k_1} \frac{\langle P i_1 P i_2 | I(\omega - \varepsilon_{P i_1}) | n k_2 \rangle \langle n | \delta V | k_1 \rangle}{(\omega - \varepsilon_{k_1} - i0)(E_i^{(0)} - \omega - \varepsilon_{k_2} + i0)(\omega - \varepsilon_n(1 - i0))} \\ & + \sum_{n \neq k_2} \frac{\langle P i_1 P i_2 | I(\omega - \varepsilon_{P i_1}) | k_1 n \rangle \langle n | \delta V | k_2 \rangle}{(\omega - \varepsilon_{k_1} + i0)(E_i^{(0)} - \omega - \varepsilon_{k_2} - i0)(E_i^{(0)} - \omega - \varepsilon_n(1 - i0))} \\ & - \sum_{n \neq P i_1} \frac{\langle P i_1 | \delta V | n \rangle \langle n P i_2 | I(\omega - \varepsilon_{k_1}) | k_1 k_2 \rangle}{(\omega - \varepsilon_{P i_1} - i0)(E_k^{(0)} - \omega - \varepsilon_{P i_2} + i0)(\omega - \varepsilon_n(1 - i0))} \\ & \left. - \sum_{n \neq P i_2} \frac{\langle P i_2 | \delta V | n \rangle \langle P i_1 n | I(\omega - \varepsilon_{k_1}) | k_1 k_2 \rangle}{(\omega - \varepsilon_{P i_1} + i0)(E_k^{(0)} - \omega - \varepsilon_{P i_2} - i0)(E_k^{(0)} - \omega - \varepsilon_n(1 - i0))} \right\}. \end{aligned} \quad (2.182)$$

The term $\Delta H_{ik}^{(2,\text{irred})}$ approaches zero if $E_i^{(0)} \rightarrow E_k^{(0)}$. The expressions (2.179) and (2.181) differ by the term $\Delta H_{ik}^{(2,\text{irred})}$ which can be represented as

$$\Delta H_{ik}^{(2,\text{irred})} = (E_i^{(0)} - E_k^{(0)}) O_{ik}^{(2)}. \quad (2.183)$$

Here $O^{(2)}$ is an operator of second order in the perturbation parameter which we denote by λ_0 (for simplicity, we assume here that δV and the interelectronic-interaction operator $I(\omega)$ are characterized by the same perturbation parameter). This fact can be understood by observing that the integrand in (2.182) is the sum of terms which contain singularities from the external electron propagators only above or below of the real axis and, therefore, integrating over ω cannot result in the appearance of contributions $\sim 1/(E_i^{(0)} - E_k^{(0)})$ that could compensate the factor $(E_i^{(0)} - E_k^{(0)})$. In particular, it means that $O_{ik}^{(2)}$ remains finite when $E_i^{(0)} \rightarrow E_k^{(0)}$. It can be shown that the term $\Delta H_{ik}^{(2,\text{irred})}$ contributes only to third and higher orders in λ_0 and, therefore, can be omitted if we restrict our calculations to second order in λ_0 . Let us prove this fact for the case of two quasidegenerate levels. In this case the energy levels are determined from the equation

$$(E - H_{11})(E - H_{22}) - H_{12}H_{21} = 0 \quad (2.184)$$

which yields

$$E_{1,2} = \frac{H_{11} + H_{22}}{2} \pm \frac{1}{2} \sqrt{(H_{11} - H_{22})^2 + 4H_{12}H_{21}}. \quad (2.185)$$

If $E_1 - E_2 \sim \lambda_0$, the proof of the statement is evident. If $E_1 - E_2 \gg \lambda_0$, the contribution of the second-order off-diagonal matrix elements is given by

$$\Delta E_{1,2} \approx \pm (H_{12}^{(2)} H_{21}^{(1)} + H_{12}^{(1)} H_{21}^{(2)}) / (E_1^{(0)} - E_2^{(0)}). \quad (2.186)$$

From this equation we find that the terms $\sim (E_i^{(0)} - E_k^{(0)}) O_{ik}^{(2)}$ in $H_{ik}^{(2)}$ contribute only to third and higher orders in λ_0 .

Equations (2.178)-(2.180) will give a part of the VP screening contribution, if we replace δV by U_{VP} . The total contribution of the VP screening diagrams for quasidegenerate states is derived in Ref. [44]. Corresponding formulas for the SE screening diagrams are derived in Ref. [75].

F. Nuclear recoil corrections

So far we considered the nucleus as a source of the external Coulomb field V_C . This consideration corresponds to the approximation of an infinite nuclear mass. However, high precision calculations of the energy levels in high- Z few-electron atoms must include also the nuclear recoil corrections to first order in m/M (M is the nuclear mass) and to zeroth order in α (but to all orders in αZ). As was shown in [41], these corrections can be included in calculations of the energy levels by adding an additional term to the standard Hamiltonian of the electron-positron field interacting with the quantized electromagnetic field and with the Coulomb field of the nucleus V_C . In the Coulomb gauge and the Schrödinger representation, this term is given by


$$H_M = \frac{1}{2M} \int d\mathbf{x} \psi^\dagger(\mathbf{x}) (-i\nabla_{\mathbf{x}}) \psi(\mathbf{x}) \int d\mathbf{y} \psi^\dagger(\mathbf{y}) (-i\nabla_{\mathbf{y}}) \psi(\mathbf{y}) - \frac{eZ}{M} \int d\mathbf{x} \psi^\dagger(\mathbf{x}) (-i\nabla_{\mathbf{x}}) \psi(\mathbf{x}) \mathbf{A}(0) + \frac{e^2 Z^2}{2M} \mathbf{A}^2(0). \quad (2.187)$$

H_M taken in the interaction representation must be added to the interaction Hamiltonian. It gives the following additional lines and vertices to the Feynman rules (we assume that the Coulomb gauge and the Furry picture are used).

1. *Coulomb contribution.* An additional line ("Coulomb-recoil" line) appears to be

$$\begin{array}{c} \omega \\ \bullet \cdots \cdots \bullet \\ \mathbf{x} \qquad \qquad \mathbf{y} \end{array} \quad \frac{i}{2\pi} \frac{\delta_{kl}}{M} \int_{-\infty}^{\infty} d\omega.$$


This line joins two vertices each of which corresponds to



$$-2\pi i \gamma^0 \delta(\omega_1 - \omega_2 - \omega_3) \int d\mathbf{x} p_k,$$

where $\mathbf{p} = -i\nabla_{\mathbf{x}}$ and $k = 1, 2, 3$.

2. *One-transverse-photon contribution.* An additional vertex on an electron line appears to be



$$-2\pi i \gamma^0 \delta(\omega_1 - \omega_2 - \omega_3) \frac{eZ}{M} \int d\mathbf{x} p_k ,$$

The transverse photon line attached to this vertex (at the point $\mathbf{x}$) is

$$\begin{array}{c} \omega \\ \bullet \text{---} \text{---} \text{---} \bullet \\ \mathbf{x} \qquad \qquad \mathbf{y} \end{array} \quad \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega D_{kl}(\omega, \mathbf{y}) .$$

At the point $\mathbf{y}$ this line is to be attached to a usual vertex, where we have $-2\pi i e \gamma^0 \alpha_l 2\pi \delta(\omega_1 - \omega_2 - \omega_3) \int d\mathbf{y}$. α_l ($l = 1, 2, 3$) are the usual Dirac matrices.

3. *Two-transverse-photon contribution.* An additional line ("two-transverse-photon-recoil" line) appears to be

$$\begin{array}{c} \omega \\ \text{---} \bullet \text{---} \text{---} \\ \mathbf{x} \qquad \qquad \mathbf{y} \end{array} \quad \frac{i}{2\pi} \frac{e^2 Z^2}{M} \int_{-\infty}^{\infty} d\omega D_{il}(\omega, \mathbf{x}) D_{lk}(\omega, \mathbf{y}) .$$

This line joins usual vertices (see the previous item).

Let us apply this formalism to the case of a single level a in a one-electron atom. To find the Coulomb nuclear recoil correction, we have to calculate the contribution of the diagram shown in Fig. 25. According to the Feynman rules given above we obtain

$$\Delta g_{aa}^{(1)}(E) = \frac{1}{(E - E_a^{(0)})^2} \frac{1}{M} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_n \frac{\langle a | p_i | n \rangle \langle n | p_i | a \rangle}{\omega - \varepsilon_n(1 - i0)} . \quad (2.188)$$

The formula (2.51) gives

$$\Delta E_C = \frac{1}{M} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_n \frac{\langle a | p_i | n \rangle \langle n | p_i | a \rangle}{\omega - \varepsilon_n(1 - i0)} . \quad (2.189)$$

The one-transverse-photon nuclear recoil correction corresponds to the diagrams shown in Fig. 26. A similar calculation yields

$$\begin{aligned} \Delta E_{\text{tr}(1)} = & \frac{4\pi\alpha Z}{M} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_n \left\{ \frac{\langle a | p_i | n \rangle \langle n | \alpha_k D_{ik}(\varepsilon_a - \omega) | a \rangle}{\omega - \varepsilon_n(1 - i0)} \right. \\ & \left. + \frac{\langle a | \alpha_k D_{ik}(\varepsilon_a - \omega) | n \rangle \langle n | p_i | a \rangle}{\omega - \varepsilon_n(1 - i0)} \right\} . \end{aligned} \quad (2.190)$$

The two-transverse-photon nuclear recoil correction is defined by the diagram shown in Fig. 27. We find

$$\begin{aligned} \Delta E_{\text{tr}(2)} = & \frac{(4\pi\alpha Z)^2}{M} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \sum_n \\ & \times \frac{\langle a | \alpha_i D_{il}(\varepsilon_a - \omega) | n \rangle \langle n | \alpha_k D_{lk}(\varepsilon_a - \omega) | a \rangle}{\omega - \varepsilon_n(1 - i0)} . \end{aligned} \quad (2.191)$$

The sum of all the contributions is

$$\begin{aligned} \Delta E = & \frac{1}{M} \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \langle a | [p_i + 4\pi\alpha Z \alpha_l D_{li}(\omega)] \\ & \times G_C(\omega + \varepsilon_a) [p_i + 4\pi\alpha Z \alpha_m D_{mi}(\omega)] | a \rangle , \end{aligned} \quad (2.192)$$

where $G_C(\omega)$ is the Dirac-Coulomb Green function defined above. For practical calculations it is convenient to represent the expression (2.192) by the sum of a lower-order term ΔE_L and a higher-order term ΔE_H :

$$\Delta E = \Delta E_L + \Delta E_H, \quad (2.193)$$

$$\Delta E_L = \frac{1}{2M} \langle a | [p_i^2 - (D_i(0)p_i + p_i D_i(0))] | a \rangle, \quad (2.194)$$

$$\Delta E_H = \frac{i}{2\pi M} \int_{-\infty}^{\infty} d\omega \langle a | \left(D_i(\omega) - \frac{[p_i, V_C]}{\omega + i0} \right) G_C(\omega + \varepsilon_a) \left(D_i(\omega) + \frac{[p_i, V_C]}{\omega + i0} \right) | a \rangle, \quad (2.195)$$

where $D_i(\omega) = -4\pi\alpha Z\alpha_l D_{li}(\omega)$. The term ΔE_L contains all the recoil corrections within the $(\alpha Z)^4 m^2/M$ approximation, while the term ΔE_H contains the contribution of order $(\alpha Z)^5 m^2/M$ and all contributions of higher orders in αZ which are not included in ΔE_L . The formulas (2.193)-(2.195) were first derived by a quasipotential method in Ref. [50] and subsequently rederived by other methods in Refs. [76,77]. The representation (2.192) was found in Ref. [76].

Consider now a two-electron atom. For simplicity, as usual, we assume that the unperturbed wave function is a one-determinant function (2.120). The nuclear recoil correction is the sum of the one-electron and two-electron contributions. Using the Feynman rules and the formula (2.51), one easily finds that the one-electron contribution is equal to the sum of the expressions (2.192) for the a and b states. The two-electron contributions correspond to the diagrams shown in Figs. 28-30. A simple calculation of these diagrams yields

$$\begin{aligned} \Delta E^{(\text{int})} = \frac{1}{M} \sum_P (-1)^P \langle Pa | [p_i + 4\pi\alpha Z\alpha_l D_{li}(\varepsilon_{Pa} - \varepsilon_a)] | a \rangle \\ \times \langle Pb | [p_i + 4\pi\alpha Z\alpha_m D_{mi}(\varepsilon_{Pb} - \varepsilon_b)] | b \rangle. \end{aligned} \quad (2.196)$$

The formula (2.196) was first derived by a quasipotential method in Ref. [51].

III. TRANSITION PROBABILITIES AND CROSS SECTIONS OF SCATTERING PROCESSES

According to the basic principles of quantum field theory [78], the number of the particles scattered into the interval $d\mathbf{p}'_1 \cdots d\mathbf{p}'_r$ for a unit time and in a unit volume is

$$dW_{p \rightarrow p'} = (2\pi)^{3s+1} n_1 \cdots n_s |\tau_{p'p}|^2 \delta(E_p - E_{p'}) d\mathbf{p}'_1 \cdots d\mathbf{p}'_r, \quad (3.1)$$

where p, p' are the initial and final states of the system, respectively; $\tau_{p'p}$ is the amplitude of the process defined by

$$\langle p' | (S - I) | p \rangle = 2\pi i \delta(E_p - E_{p'}) \tau_{p'p}, \quad (3.2)$$

S is the scattering operator, I is the identity operator, s is the number of initial particles, r is the number of final particles; $n_1, \dots, n_s$ are the average numbers of particles per unit volume.

We will consider the scattering of photons and electrons by an atom that is located at the origin of the coordinate frame. The differential cross section is defined by

$$d\sigma = \frac{dW_{p \rightarrow p'}}{j}, \quad (3.3)$$

where j is the current of initial particles (for photons $j = nc$; for electrons $j = nv$, where v is the velocity of the electrons in the nucleus frame). The total cross section can be found by integrating the differential cross section over all final states. The cross section for the elastic scattering is

$$\sigma_{\text{tot}}^{(\text{elast})}(p) = \frac{(2\pi)^4}{v} \int d\mathbf{p}' \delta(E_p - E_{p'}) |\tau_{p'p}|^2. \quad (3.4)$$

The total (elastic plus inelastic) cross section can be found by employing the optical theorem

$$\sigma_{\text{tot}}(p) = \frac{2(2\pi)^3}{v} \text{Im} \tau_{pp}. \quad (3.5)$$

In terms of the amplitude $f_{p'p}$ which is defined so that $d\sigma = |f_{p'p}|^2 d\Omega$, the optical theorem has a well-known form (see, e.g., [79])

$$\sigma_{\text{tot}}(p) = \frac{4\pi}{|\mathbf{p}|} \text{Im} f_{pp}. \quad (3.6)$$

The aim of this section is to derive formulas for the transition and scattering amplitudes for various processes in the framework of QED.

A. Photon emission by an atom

Consider the process of the photon emission by an atom. According to the standard reduction technique (see, e.g., [28,56]), the atomic transition amplitude from state a to state b accompanied by photon emission with momentum $\mathbf{k}_f$ and polarization ϵ_f is

$$S_{\gamma_f, b; a} = \langle b | a_{\text{out}}(k_f, \epsilon_f) | a \rangle = -i Z_3^{-\frac{1}{2}} \int d^4 y \frac{\epsilon_f^{\nu*} \exp(i \mathbf{k}_f \cdot \mathbf{y})}{\sqrt{2 k_f^0 (2\pi)^3}} \langle b | j_\nu(y) | a \rangle. \quad (3.7)$$

Here $j_\nu(y)$ is the electron-positron current operator in the Heisenberg representation, $|a\rangle$ and $|b\rangle$ are the vectors of the initial and final states in the Heisenberg representation, Z_3 is a renormalization constant, $a \cdot b \equiv a_\nu b^\nu$, $\epsilon_f = (0, \epsilon_f)$, $k_f = (k_f^0, \mathbf{k}_f)$, and $k_f^0 \equiv |\mathbf{k}_f|$. Employing the equation (see Appendix A)

$$j^\nu(y) = \exp(i H y^0) j^\nu(0, \mathbf{y}) \exp(-i H y^0), \quad (3.8)$$

we obtain

$$\begin{aligned} S_{\gamma_f, b; a} &= -i Z_3^{-\frac{1}{2}} \int d^4 y \exp[i(E_b + k_f^0 - E_a)y^0] A_f^{\nu*}(\mathbf{y}) \langle b | j_\nu(0, \mathbf{y}) | a \rangle \\ &= -2\pi i Z_3^{-\frac{1}{2}} \delta(E_b + k_f^0 - E_a) \int d\mathbf{y} A_f^{\nu*}(\mathbf{y}) \langle b | j_\nu(0, \mathbf{y}) | a \rangle, \end{aligned} \quad (3.9)$$

where

$$A_f^\nu(\mathbf{x}) = \frac{\epsilon_f^\nu \exp(i \mathbf{k}_f \cdot \mathbf{x})}{\sqrt{2 k_f^0 (2\pi)^3}} \quad (3.10)$$

is the wave function of the emitted photon. Since $|a\rangle$ and $|b\rangle$ are bound states, equation (3.9) as well as the standard reduction technique [28,56] cannot be used for a direct evaluation of the amplitude. The desired calculation formula can be derived within the two-time Green function formalism [30–32].

To formulate the method for a general case, we assume that in zeroth approximation the state a belongs to an s_a -dimensional subspace of unperturbed degenerate states Ω_a and the state b belongs to an s_b -dimensional subspace of unperturbed degenerate states Ω_b . We denote the projectors onto these subspaces by $P_a^{(0)}$ and $P_b^{(0)}$, respectively. We denote the exact states originating from Ω_a by $|n_a\rangle$ and the exact states originating from Ω_b by $|n_b\rangle$. We also assume that on an intermediate stage of the calculations a non-zero photon mass μ is introduced. It is considered to be larger than the energy splitting of the initial and final states under consideration and much smaller than the distance to other levels.

We introduce

$$\begin{aligned} &\mathcal{G}_{\gamma_f}(E', E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) \delta(E' + k^0 - E) \\ &= \frac{1}{2\pi i} \frac{1}{2\pi} \frac{1}{N!} \int_{-\infty}^{\infty} dx^0 dx'^0 \int d^4 y \exp(i E' x'^0 - i E x^0) \exp(i k^0 y^0) \\ &\quad \times A_f^{\nu*}(\mathbf{y}) \langle 0 | T \psi(x'^0, \mathbf{x}'_1) \cdots \psi(x'^0, \mathbf{x}'_N) \\ &\quad \times j_\nu(y) \bar{\psi}(x^0, \mathbf{x}_N) \cdots \bar{\psi}(x^0, \mathbf{x}_1) | 0 \rangle, \end{aligned} \quad (3.11)$$

where, as in the previous section, $\psi(x)$ is the electron-positron field operator in the Heisenberg representation. Let us investigate the singularities of $\mathcal{G}_{\gamma_f}$ in the vicinity of the points $E' \approx E_b^{(0)}$ and $E \approx E_a^{(0)}$. Using the transformation rules

$$\begin{aligned}\psi(x^0, \mathbf{x}) &= \exp(iHy^0)\psi(x^0 - y^0, \mathbf{x})\exp(-iHy^0), \\ j(y^0, \mathbf{y}) &= \exp(iHy^0)j(0, \mathbf{y})\exp(-iHy^0),\end{aligned}\tag{3.12}$$

we obtain

$$\begin{aligned}\mathcal{G}_{\gamma_f}(E', E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &\delta(E' + k^0 - E) \\ &= \frac{1}{2\pi i} \frac{1}{2\pi} \frac{1}{N!} \int_{-\infty}^{\infty} dt dt' \int d^4y \exp(iE't' - iEt) \exp[i(E' + k^0 - E)y^0] \\ &\quad \times A_f^{\nu*}(\mathbf{y}) \langle 0 | T \psi(t', \mathbf{x}'_1) \cdots \psi(t', \mathbf{x}'_N) \\ &\quad \times j_\nu(0, \mathbf{y}) \bar{\psi}(t, \mathbf{x}_N) \cdots \bar{\psi}(t, \mathbf{x}_1) | 0 \rangle \\ &= \frac{1}{2\pi i} \delta(E' + k^0 - E) \frac{1}{N!} \int_{-\infty}^{\infty} dt dt' \int d\mathbf{y} \exp(iE't' - iEt) \\ &\quad \times A_f^{\nu*}(\mathbf{y}) \langle 0 | T \psi(t', \mathbf{x}'_1) \cdots \psi(t', \mathbf{x}'_N) \\ &\quad \times j_\nu(0, \mathbf{y}) \bar{\psi}(t, \mathbf{x}_N) \cdots \bar{\psi}(t, \mathbf{x}_1) | 0 \rangle.\end{aligned}\tag{3.13}$$

Using again the time-shift transformation rules, we obtain

$$\begin{aligned}\mathcal{G}_{\gamma_f}(E', E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{1}{2\pi i} \frac{1}{N!} \int_{-\infty}^{\infty} dt dt' \int d\mathbf{y} \exp(iE't' - iEt) \sum_{n_1, n_2} A_f^{\nu*}(\mathbf{y}) \\ &\quad \times \exp(-iE_{n_1}t') \exp(iE_{n_2}t) \theta(t') \theta(-t) \langle 0 | T \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) | n_1 \rangle \\ &\quad \times \langle n_1 | j_\nu(0, \mathbf{y}) | n_2 \rangle \langle n_2 | \bar{\psi}(0, \mathbf{x}_N) \cdots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle + \dots.\end{aligned}\tag{3.14}$$

Here we have assumed $E_0 = 0$, as in the previous section. Taking into account the identities

$$\begin{aligned}\int_0^\infty dt \exp[i(E' - E_{n_1})t] &= \frac{i}{E' - E_{n_1} + i0}, \\ \int_{-\infty}^0 dt \exp[i(-E + E_{n_2})t] &= \frac{i}{E - E_{n_2} + i0},\end{aligned}\tag{3.15}$$

we find

$$\begin{aligned}\mathcal{G}_{\gamma_f}(E', E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &= \frac{i}{2\pi} \frac{1}{N!} \sum_{n_1, n_2} \int d\mathbf{y} A_f^{\nu*}(\mathbf{y}) \frac{1}{E' - E_{n_1} + i0} \frac{1}{E - E_{n_2} + i0} \\ &\quad \times \langle 0 | T \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) | n_1 \rangle \langle n_1 | j_\nu(0, \mathbf{y}) | n_2 \rangle \\ &\quad \times \langle n_2 | \bar{\psi}(0, \mathbf{x}_N) \cdots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle + \dots.\end{aligned}\tag{3.16}$$

We are interested in the analytical properties of $\mathcal{G}_{\gamma_f}$ as a function of the two complex variables E' and E in the region $E' \approx E_b^{(0)}$, $E \approx E_a^{(0)}$. These properties can be studied using the double spectral representation of this type of Green function which is given in Appendix E (a similar representation was derived in Ref. [64]). As it follows from the spectral representation, the terms which are omitted in equation (3.16) are regular functions of E' or E if $E' \approx E_b^{(0)}$ and $E \approx E_a^{(0)}$. The equation (3.16) and the spectral representation given in Appendix E show that, for a non-zero photon mass μ , the Green function $\mathcal{G}_{\gamma_f}(E', E)$ has isolated poles in the variables E' and E at the points $E' = E_{n_b}$ and $E = E_{n_a}$, respectively. Let us now introduce a Green function $g_{\gamma_f, b; a}(E', E)$ by

$$g_{\gamma_f, b; a}(E', E) = P_b^{(0)} \mathcal{G}_{\gamma_f}(E', E) \gamma_1^0 \cdots \gamma_N^0 P_a^{(0)},\tag{3.17}$$

where, as in (2.39), the integration over the electron coordinates is implicit. According to equation (3.16) (see also Appendix E) the Green function $g_{\gamma_f, b; a}(E', E)$ can be written as

$$\begin{aligned}g_{\gamma_f, b; a}(E', E) &= \frac{i}{2\pi} \sum_{n_a=1}^{s_a} \sum_{n_b=1}^{s_b} \frac{1}{E' - E_{n_b}} \frac{1}{E - E_{n_a}} \varphi_{n_b} \int d\mathbf{y} A_f^{\nu*}(\mathbf{y}) \langle n_b | j_\nu(0, \mathbf{y}) | n_a \rangle \varphi_{n_a}^\dagger \\ &\quad + \text{terms that are regular functions of } E' \text{ or } E \text{ if } E' \approx E_b^{(0)} \\ &\quad \text{and } E \approx E_a^{(0)},\end{aligned}\tag{3.18}$$

where the vectors φ_k are defined by equation (2.65). Let the contours Γ_a and Γ_b surround the poles corresponding to the initial and final levels, respectively, and keep outside other singularities of $g_{\gamma_f, b; a}(E', E)$ including the cuts starting from the lower-lying bound states. Comparing equation (3.18) with equation (3.9) and taking into account the biorthogonality condition (2.71), we obtain the desired formula [30]

$$S_{\gamma_f, b; a} = Z_3^{-1/2} \delta(E_b + k_f^0 - E_a) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE v_b^\dagger g_{\gamma_f, b; a}(E', E) v_a, \quad (3.19)$$

where by a we imply one of the initial states and by b one of the final states under consideration. The vectors v_k are determined from equations (2.76)- (2.77).

In the case of a single initial state (a) and a single final state (b), the vectors v_a and v_b simply become normalization factors. So, for the initial state,

$$v_a^* P_a v_a = v_a^* \frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}(E) v_a = 1 \quad (3.20)$$

and, therefore,

$$|v_a|^2 = \left[\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}(E) \right]^{-1}. \quad (3.21)$$

Choosing

$$v_a = \left[\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}(E) \right]^{-1/2}, \quad v_b = \left[\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}(E) \right]^{-1/2}, \quad (3.22)$$

we obtain

$$\begin{aligned} S_{\gamma_f, b; a} &= Z_3^{-1/2} \delta(E_b + k_f^0 - E_a) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}(E', E) \\ &\times \left[\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}(E) \right]^{-1/2} \left[\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}(E) \right]^{-1/2}. \end{aligned} \quad (3.23)$$

For practical calculations of the transition amplitude, it is convenient to express the Green function $g_{\gamma_f, b; a}(E', E)$ in terms of the Fourier transform of the $2N$ -time Green function,

$$\begin{aligned} g_{\gamma_f, b; a}(E', E) \delta(E' + k^0 - E) &= \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \\ &\times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\ &\times P_b^{(0)} G_{\gamma_f}(p_1^0, \dots, p_N^0; k^0; p_1'^0, \dots, p_N'^0) \gamma_1^0 \cdots \gamma_N^0 P_a^{(0)}, \end{aligned} \quad (3.24)$$

where

$$\begin{aligned} &G_{\gamma_f}((p_1^0, \mathbf{x}'_1), \dots, (p_N^0, \mathbf{x}'_N); k^0; (p_1^0, \mathbf{x}_1), \dots, (p_N^0, \mathbf{x}_N)) \\ &= \frac{2\pi}{i} \frac{1}{(2\pi)^{2N+1}} \int_{-\infty}^{\infty} dx_1^0 \cdots dx_N^0 dx_1'^0 \cdots dx_N'^0 \int d^4 y \\ &\times \exp(ip_1^0 x_1^0 + \dots + ip_N^0 x_N^0 - ip_1'^0 x_1'^0 - \dots - ip_N'^0 x_N'^0 + ik^0 y^0) \\ &\times A_f^{\nu*}(\mathbf{y}) \langle 0 | T \psi(x'_1) \cdots \psi(x'_N) j_\nu(y) \bar{\psi}(x_N) \cdots \bar{\psi}(x_1) | 0 \rangle. \end{aligned} \quad (3.25)$$

The Green function G_{γ_f} is constructed by perturbation theory after the transition in (3.25) to the interaction representation and using the Wick theorem. The Feynman rules for G_{γ_f} differ from those for G considered in the previous section only by the presence of an outgoing photon line that corresponds to the wave function of the emitted photon $A_f^{\nu*}(\mathbf{x})$. The Feynman rule for a vertex in which a real photon is emitted remains the same as for a virtual photon vertex. The energy variable of the emitted photon (k^0) in the expression corresponding to a real photon vertex is considered as a free variable ($k^0 \neq k_f^0 = |\mathbf{k}_f|$) which, due to the energy conservation, can be expressed in terms of the initial (E) and final (E') atomic energy variables.

B. Transition probability in a one-electron atom

To demonstrate the practical ability of the formalism, in this section we derive formulas for the transition probability in a one-electron atom to zeroth and first orders in α . An application of the method for two-electron atoms is considered in [80].

1. Zeroth order approximation

To zeroth order the transition amplitude is described by the diagram shown in Fig. 31. The formula (3.23) gives

$$S_{\gamma_f, b; a}^{(0)} = \delta(E_b + k_f^0 - E_a) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(0)}(E', E), \quad (3.26)$$

where the superscript indicates the order in α . Here we have taken into account that

$$\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}^{(0)}(E) = \frac{1}{2\pi i} \oint_{\Gamma_a} dE \frac{1}{E - \varepsilon_a} = 1 \quad (3.27)$$

and a similar equation exists for the b state. According to the Feynman rules, we have

$$\begin{aligned} G_{\gamma_f}^{(0)}((E', \mathbf{x}'); k^0; (E, \mathbf{x})) &= \int d\mathbf{y} \frac{i}{2\pi} S(E', \mathbf{x}', \mathbf{y}) (-2\pi i e \gamma^\nu) \delta(E' + k^0 - E) \\ &\quad \times A_{f, \nu}^*(\mathbf{y}) \frac{i}{2\pi} S(E, \mathbf{y}, \mathbf{x}). \end{aligned} \quad (3.28)$$

Substituting the expression (3.28) into the definition of $g_{\gamma_f, b; a}(E', E)$ (see equation (3.24)), we obtain

$$g_{\gamma_f, b; a}^{(0)}(E', E) = \frac{i}{2\pi} \frac{1}{E' - \varepsilon_b} \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle \frac{1}{E - \varepsilon_a}. \quad (3.29)$$

The equations (3.26), (3.29) yield

$$S_{\gamma_f, b; a}^{(0)} = -2\pi i \delta(E_b + k_f^0 - E_a) \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle \quad (3.30)$$

or, in accordance with the definition (3.2),

$$\tau_{\gamma_f, b; a}^{(0)} = -\langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle = \langle b | e \boldsymbol{\alpha} \cdot \mathbf{A}_f^* | a \rangle. \quad (3.31)$$

The transition probability to zeroth order is

$$\begin{aligned} dW_{\gamma_f, b; a}^{(0)} &= 2\pi |\tau_{\gamma_f, b; a}^{(0)}|^2 \delta(E_b + k_f^0 - E_a) d\mathbf{k}_f \\ &= 2\pi \frac{e^2}{2k_f^0 (2\pi)^3} |\langle b | \epsilon_f^{\nu*} \alpha_\nu \exp(-i\mathbf{k}_f \cdot \mathbf{x}) | a \rangle|^2 \delta(E_b + k_f^0 - E_a) d\mathbf{k}_f. \end{aligned} \quad (3.32)$$

Integrating over the photon energy, we obtain

$$dW_{\gamma_f, b; a}^{(0)} = \alpha \frac{k_f^0}{2\pi} |\boldsymbol{\epsilon}_f^* \cdot \mathbf{j}_{ba}(\mathbf{k}_f)|^2 d\Omega_f, \quad (3.33)$$

where

$$\mathbf{j}_{ba}(\mathbf{k}_f) = \langle b | \boldsymbol{\alpha} \exp(-i\mathbf{k}_f \cdot \mathbf{x}) | a \rangle. \quad (3.34)$$

2. QED corrections of first order in α

The QED corrections of first order in α are defined by the diagrams shown in Fig. 32. Let us consider the derivation of the formulas for the self-energy (SE) corrections (the diagrams (a)-(c)) in detail. The formula (3.23) gives in the order under consideration

$$S_{\gamma_f, b; a}^{(1)} = \delta(E_b + k_f^0 - E_a) \left\{ \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(1)}(E', E) - \frac{1}{2} \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(0)}(E', E) \left[\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}^{(1)}(E) + \frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}^{(1)}(E) \right] \right\}, \quad (3.35)$$

where $g_{aa}^{(1)}(E)$ and $g_{bb}^{(1)}(E)$ are defined by the first-order self-energy diagram. Here we have omitted a term of first order in α which comes from the factor $Z_3^{-1/2}$ since it has to be combined with the vacuum-polarization (VP) correction. Consider first the diagram (a). According to the Feynman rules, we have

$$G_{\gamma_f}^{(1,a)}((E', \mathbf{x}'); k^0; (E, \mathbf{x})) = \delta(E' + k^0 - E) \int d\mathbf{y} d\mathbf{y}' d\mathbf{z} \frac{i}{2\pi} S(E', \mathbf{x}', \mathbf{y}) \frac{2\pi}{i} \gamma^0 \Sigma(E', \mathbf{y}', \mathbf{y}) \times \frac{i}{2\pi} S(E', \mathbf{y}, \mathbf{z}) A_{f, \nu}^*(\mathbf{z}) (-2\pi i e \gamma^\nu) \frac{i}{2\pi} S(E, \mathbf{z}, \mathbf{x}), \quad (3.36)$$

where

$$\Sigma(E', \mathbf{y}', \mathbf{y}) = e^2 \frac{i}{2\pi} \int d\omega \gamma^0 \gamma^\rho S(E' - \omega, \mathbf{y}', \mathbf{y}) \gamma^\sigma D_{\rho\sigma}(\omega, \mathbf{y}' - \mathbf{y}) \quad (3.37)$$

is the kernel of the self-energy operator and $D_{\rho\sigma}(\omega, \mathbf{y}' - \mathbf{y})$ is the photon propagator for a non-zero photon mass. According to the definition of $g_{\gamma_f, b; a}(E', E)$ (see equation (3.24)), we find

$$g_{\gamma_f, b; a}^{(1,a)}(E', E) = \frac{i}{2\pi} \sum_n \frac{\langle b | \Sigma(E') | n \rangle \langle n | e \alpha^\nu A_{f, \nu}^* | a \rangle}{(E' - \varepsilon_b)(E' - \varepsilon_n)(E - \varepsilon_a)} \quad (3.38)$$

and

$$\oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(1,a)}(E', E) = -2\pi i \left[\sum_n^{n \neq b} \frac{\langle b | \Sigma(\varepsilon_b) | n \rangle \langle n | e \alpha^\nu A_{f, \nu}^* | a \rangle}{\varepsilon_b - \varepsilon_n} + \langle b | \Sigma'(\varepsilon_b) | b \rangle \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle \right], \quad (3.39)$$

where $\Sigma'(\varepsilon_b) \equiv \frac{d\Sigma(\varepsilon)}{d\varepsilon} \Big|_{\varepsilon=\varepsilon_b}$. A similar calculation of the diagram (b) gives

$$\oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(1,b)}(E', E) = -2\pi i \left[\sum_n^{n \neq a} \frac{\langle b | e \alpha^\nu A_{f, \nu}^* | n \rangle \langle n | \Sigma(\varepsilon_a) | a \rangle}{\varepsilon_a - \varepsilon_n} + \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle \langle a | \Sigma'(\varepsilon_a) | a \rangle \right]. \quad (3.40)$$

The second ("reducible") terms in equations (3.39) and (3.40) have to be combined with the second term in equation (3.35). Taking into account that

$$\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}^{(1)}(E) = \langle a | \Sigma'(\varepsilon_a) | a \rangle, \quad (3.41)$$

$$\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}^{(1)}(E) = \langle b | \Sigma'(\varepsilon_b) | b \rangle, \quad (3.42)$$

we obtain

$$-\frac{1}{2} \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(0)}(E', E) \left[\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}^{(1)}(E) + \frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}^{(1)}(E) \right] = 2\pi i \left[\frac{1}{2} \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle (\langle b | \Sigma'(\varepsilon_b) | b \rangle + \langle a | \Sigma'(\varepsilon_a) | a \rangle) \right]. \quad (3.43)$$

For the diagram (c) we find

$$g_{\gamma_f, b; a}^{(1, c)}(E', E) = \frac{i}{2\pi} \frac{1}{E' - \varepsilon_b} \int d\mathbf{x} d\mathbf{y} d\mathbf{z} \bar{\psi}_b(\mathbf{x}) \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \gamma^\rho S(E' - \omega, \mathbf{x}, \mathbf{z}) \\ \times e A_{f, \nu}^*(\mathbf{z}) \gamma^\nu S(E - \omega, \mathbf{z}, \mathbf{y}) \gamma^\sigma e^2 D_{\rho\sigma}(\omega, \mathbf{x} - \mathbf{y}) \psi_a(\mathbf{y}) \frac{1}{E - \varepsilon_a}. \quad (3.44)$$

Substituting (3.44) into (3.35), we obtain

$$\oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; a}^{(1, c)}(E', E) = -2\pi i \int d\mathbf{z} e A_{f, \nu}^*(\mathbf{z}) \Lambda^\nu(\varepsilon_b, \varepsilon_a, \mathbf{z}), \quad (3.45)$$

where

$$\Lambda^\nu(\varepsilon_b, \varepsilon_a, \mathbf{z}) = e^2 \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \int d\mathbf{x} d\mathbf{y} \bar{\psi}_b(\mathbf{x}) \gamma^\rho S(\varepsilon_b - \omega, \mathbf{x}, \mathbf{z}) \gamma^\nu S(\varepsilon_a - \omega, \mathbf{z}, \mathbf{y}) \\ \times \gamma^\sigma D_{\rho\sigma}(\omega, \mathbf{x} - \mathbf{y}) \psi_a(\mathbf{y}). \quad (3.46)$$

Summing all the first order SE contributions derived above and adding the contribution of the mass-counterterm diagrams (Fig. 33), we find

$$S_{\gamma_f, b; a}^{(1, \text{SE})} = -2\pi i \delta(E_b + k_f^0 - E_a) \left[\sum_n^{n \neq b} \frac{\langle b | \Sigma(\varepsilon_b) - \beta \delta m | n \rangle \langle n | e \alpha^\nu A_{f, \nu}^* | a \rangle}{\varepsilon_b - \varepsilon_n} \right. \\ + \sum_n^{n \neq a} \frac{\langle b | e \alpha^\nu A_{f, \nu}^* | n \rangle \langle n | \Sigma(\varepsilon_a) - \beta \delta m | a \rangle}{\varepsilon_a - \varepsilon_n} \\ + \int d\mathbf{z} e A_{f, \nu}^*(\mathbf{z}) \Lambda^\nu(\varepsilon_b, \varepsilon_a, \mathbf{z}) \\ \left. + \frac{1}{2} \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle (\langle b | \Sigma'(\varepsilon_b) | b \rangle + \langle a | \Sigma'(\varepsilon_a) | a \rangle) \right]. \quad (3.47)$$

A similar calculation of the vacuum-polarization diagrams (Fig. 32, diagrams (d)-(f)) gives

$$S_{\gamma_f, b; a}^{(1, \text{VP})} = -2\pi i \delta(E_b + k_f^0 - E_a) \left[\sum_n^{n \neq b} \frac{\langle b | U_{\text{VP}} | n \rangle \langle n | e \alpha^\nu A_{f, \nu}^* | a \rangle}{\varepsilon_b - \varepsilon_n} \right. \\ + \sum_n^{n \neq a} \frac{\langle b | e \alpha^\nu A_{f, \nu}^* | n \rangle \langle n | U_{\text{VP}} | a \rangle}{\varepsilon_a - \varepsilon_n} \\ \left. + \int d\mathbf{z} e A_{f, \nu}^*(\mathbf{z}) Q^\nu(k_f^0, \mathbf{z}) + (Z_3^{-1/2} - 1) \langle b | e \alpha^\nu A_{f, \nu}^* | a \rangle \right], \quad (3.48)$$

where

$$U_{\text{VP}}(\mathbf{x}) = \frac{\alpha}{2\pi i} \int d\mathbf{y} \frac{1}{|\mathbf{x} - \mathbf{y}|} \int_{-\infty}^{\infty} d\omega \text{Tr}[S(\omega, \mathbf{y}, \mathbf{y}) \gamma^0] \quad (3.49)$$

is the VP potential and

$$Q^\nu(k^0, \mathbf{z}) = -e^2 \int d\mathbf{x} d\mathbf{y} \bar{\psi}_b(\mathbf{x}) \gamma^\rho \psi_a(\mathbf{x}) D_{\rho\sigma}(k^0, \mathbf{x} - \mathbf{y}) \\ \times \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \text{Tr}[\gamma^\sigma S(\omega, \mathbf{y}, \mathbf{z}) \gamma^\nu S(\omega + k^0, \mathbf{z}, \mathbf{y})]. \quad (3.50)$$

Some individual terms in equations (3.47) and (3.48) contain ultraviolet divergences. These divergences arise solely from the zero- and one-potential terms in the expansion of the electron propagators in powers of the Coulomb potential. Using the standard expressions for the divergent parts of the zero- and one-potential SE terms (see, e.g., [28]) and

the Ward identity ($Z_1 = Z_2$) one easily finds that the ultraviolet divergences cancel each other in equation (3.47). As to equation (3.48), the divergent parts incorporate into the charge renormalization factor ($e = Z_3^{1/2} e_0$).

An alternative approach to the renormalization problem consists in using the renormalized field operators $\psi_R = Z_2^{-1/2} \psi$, $A_R = Z_3^{-1/2} A$, the renormalized electron charge $e = e_0 + \delta e = Z_1^{-1} Z_2 Z_3^{1/2} e_0$, and, respectively, the renormalized Green functions from the very beginning. In that approach, additional counterterms arise for the Feynman rules.

The vertex and reducible contributions to the SE corrections (third and fourth terms in equation (3.47)) contain infrared divergences which cancel each other when being considered together.

In addition to the QED corrections derived in this subsection, we must take into account the contribution originating from changing the photon energy in the zeroth-order transition probability (3.33) due to the QED correction to the energies of the bound states a and b . It follows that the total QED correction to the transition probability of first order in α is given by

$$dW_{\gamma_f, b; a}^{(1)} = 2\pi(k_f^0)^2 2\text{Re} \left\{ \tau_{\gamma_f, b; a}^{(0)*} \tau_{\gamma_f, b; a}^{(1)} \right\} d\Omega_f + \left[dW_{\gamma_f, b; a}^{(0)} \Big|_{k_f^0 = E_a - E_b} - dW_{\gamma_f, b; a}^{(0)} \Big|_{k_f^0 = \varepsilon_a - \varepsilon_b} \right]. \quad (3.51)$$

Here $\tau_{\gamma_f, b; a}^{(1)} = \tau_{\gamma_f, b; a}^{(1, \text{SE})} + \tau_{\gamma_f, b; a}^{(1, \text{VP})}$ is the QED correction given by equations (3.47) and (3.48) in accordance with the definition (3.2). E_a , E_b and ε_a , ε_b are the energies of the bound states a and b with and without the QED correction, respectively.

C. Radiative recombination of an electron with an atom

In calculations of processes which contain a free electron in the initial or final or in both states we consider that the interaction with the Coulomb field of the nucleus $V_C(\mathbf{x})$ is included in the source $\tilde{j}(x)$ which leads to a scattering [56],

$$(i \not{\partial} - m)\psi(x) = \tilde{j}(x). \quad (3.52)$$

It means that, after the transition to the "in" operators, the unperturbed Hamiltonian does not contain the interaction with the Coulomb potential. As a result, the Feynman rules contain free-electron propagators, instead of the bound-electron ones, and the vertices corresponding to the interaction of electrons with $V_C(\mathbf{x})$ appear. Since we consider the case of a strong Coulomb field, we will sum up infinite sequences of Feynman diagrams describing the interaction of electrons with the Coulomb potential. As a result of this summation, the free-electron propagators are replaced by the bound-electron propagators,

$$[p^0 - H(1 - i0)]^{-1} = [p^0 - H_0(1 - i0)]^{-1} + [p^0 - H_0(1 - i0)]^{-1} V_C [p^0 - H_0(1 - i0)]^{-1} + \dots, \quad (3.53)$$

where $H = H_0 + V_C$ and $H_0 = -i\boldsymbol{\alpha} \cdot \boldsymbol{\nabla} + \beta m$, and the free-electron wave functions are replaced by the wave functions in the Coulomb field. For instance, the wave function of an incident electron with momentum $\mathbf{p}_i$ and polarization μ_i is

$$\begin{aligned} \psi_{p_i \mu_i (+)} &= U_{p_i \mu_i} + [p_i^0 - H_0(1 - i0)]^{-1} V_C U_{p_i \mu_i} \\ &\quad + [p_i^0 - H_0(1 - i0)]^{-1} V_C [p_i^0 - H_0(1 - i0)]^{-1} U_{p_i \mu_i} + \dots \\ &= U_{p_i \mu_i} + [p_i^0 - H(1 - i0)]^{-1} V_C U_{p_i \mu_i}, \end{aligned} \quad (3.54)$$

where $p_i = (p_i^0, \mathbf{p}_i)$, $p_i^0 = \sqrt{\mathbf{p}_i^2 + m^2}$,

$$U_{p_i \mu_i} = \frac{u(p_i, \mu_i) \exp(i\mathbf{p}_i \cdot \mathbf{x})}{\sqrt{(p_i^0/m)(2\pi)^3}} \quad (3.55)$$

is the free wave function of the incident electron, and $u(p_i, \mu_i)$ is normalized by the condition $\bar{u}u = 1$.

Consider the process of the radiative recombination of an electron with momentum $\mathbf{p}_i$ and polarization μ_i with an $(N-1)$ -electron atom $X^{(Z-N+1)+}$ in a state a . As a result of the process, the N -electron atom $X^{(Z-N)+}$ in a state b and a photon with momentum $\mathbf{k}_f$ and polarization $\epsilon_f = (0, \boldsymbol{\epsilon}_f)$ arise,

$$e^-(p_i, \mu_i) + X^{(Z-N+1)+}(a) \rightarrow \gamma(k_f, \epsilon_f) + X^{(Z-N)+}(b), \quad (3.56)$$

where $k_f = (k_f^0, \mathbf{k}_f)$ and $k_f^0 = |\mathbf{k}_f|$. In this section we will assume that we consider a non-resonant process. It means that the total initial energy ($p_i^0 + E_a$) is not close to any excited-state energy of the N -electron atom. As to the resonance recombination, a detailed description of this process is given in [37] (see also the next sections of the present paper).

According to the standard reduction technique [28,56], the amplitude of the process is

$$\begin{aligned} S_{\gamma_f, b; e_i, a} &= \langle b | a_{\text{out}}(k_f, \epsilon_f) b_{\text{in}}^\dagger(p_i, \mu_i) | a \rangle \\ &= (-iZ_3^{-\frac{1}{2}})(-iZ_2^{-\frac{1}{2}}) \int d^4y d^4z \frac{\epsilon_f^{\nu*} \exp(ik_f \cdot y)}{\sqrt{2k_f^0(2\pi)^3}} \\ &\quad \times \langle b | T j_\nu(y) \bar{\psi}(z) | a \rangle (-i \overleftarrow{\not{\partial}}_z - m) \frac{u(p_i, \mu_i) \exp(-ip_i \cdot z)}{\sqrt{\frac{p_i^0}{m}(2\pi)^3}}, \end{aligned} \quad (3.57)$$

where $|a\rangle, |b\rangle$ are the vectors of the initial and final states in the Heisenberg representation. Taking into account that

$$\bar{\psi}(z)(-i \overleftarrow{\not{\partial}}_z - m) = e \bar{\psi}(z) A(z) \equiv \bar{\eta}(z) \quad (3.58)$$

we obtain

$$\begin{aligned} S_{\gamma_f, b; e_i, a} &= 2\pi \delta(E_b + k_f^0 - E_a - p_i^0) (-iZ_3^{-\frac{1}{2}})(-iZ_2^{-\frac{1}{2}}) \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) \\ &\quad \times \left\{ \int_0^\infty d\tau \langle b | j_\nu(\tau, \mathbf{y}) \bar{\eta}(0, \mathbf{z}) | a \rangle \exp(ik_f^0 \tau) \right. \\ &\quad + \int_0^\infty d\tau \langle b | \bar{\eta}(\tau, \mathbf{z}) j_\nu(0, \mathbf{y}) | a \rangle \exp(-ip_i^0 \tau) \\ &\quad \left. - i \langle b | [\psi^\dagger(0, \mathbf{z}), j_\nu(0, \mathbf{y})] | a \rangle \right\} U_{p_i \mu_i}(\mathbf{z}), \end{aligned} \quad (3.59)$$

where $A_f^\nu(\mathbf{y})$ is the wave function of the emitted photon defined by equation (3.10) and $U_{p_i \mu_i}(\mathbf{z})$ is the free wave function of the incident electron defined by equation (3.55). To construct the perturbation theory for the amplitude of the process, we introduce the Fourier transform of the corresponding two-time Green function,

$$\begin{aligned} \mathcal{G}_{\gamma_f; e_i}(E', E, p^0; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_{N-1}) &\delta(E' + k^0 - E - p^0) \\ &= \left(\frac{i}{2\pi}\right)^2 \frac{1}{\sqrt{N!}} \frac{1}{\sqrt{(N-1)!}} \int_{-\infty}^\infty dx^0 dx'^0 \exp(iE' x'^0 - iE x^0) \\ &\quad \times \int d^4y d^4z \exp(ik^0 y^0 - ip^0 z^0) A_f^{\nu*}(\mathbf{y}) \\ &\quad \times \langle 0 | T \psi(x'^0, \mathbf{x}'_1) \dots \psi(x'^0, \mathbf{x}'_N) j_\nu(y) \bar{\psi}(z) \bar{\psi}(x^0, \mathbf{x}_{N-1}) \dots \bar{\psi}(x^0, \mathbf{x}_1) | 0 \rangle \\ &\quad \times (-i \overleftarrow{\not{\partial}}_z - m) U_{p_i \mu_i}(\mathbf{z}). \end{aligned} \quad (3.60)$$

As in the derivation of the formulas for the transition amplitudes, we will assume that in zeroth approximation the initial and final states are degenerate in energy with some other states and we will use the same notations for these states as in section III(A). As usual, we also assume that a non-zero photon mass μ is introduced. The spectral representation of $\mathcal{G}_{\gamma_f; e_i}(E', E, p^0)$ can be derived in the same way as for the Green function describing the transition amplitude (see section III(A) and Appendix E). It gives

$$\begin{aligned} \mathcal{G}_{\gamma_f; e_i}(E', E, p^0; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_{N-1}) &= \frac{1}{2\pi} \frac{1}{\sqrt{N!}} \frac{1}{\sqrt{(N-1)!}} \sum_{n_a=1}^{s_a} \sum_{n_b=1}^{s_b} \frac{1}{E' - E_{n_b}} \frac{1}{E - E_{n_a}} \\ &\quad \times \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) \langle 0 | T \psi(0, \mathbf{x}'_1) \dots \psi(0, \mathbf{x}'_N) | n_b \rangle \\ &\quad \times \left\{ \int_0^\infty d\tau \langle n_b | j_\nu(\tau, \mathbf{y}) \bar{\eta}(0, \mathbf{z}) | n_a \rangle \exp(i(E - E_{n_b} + p^0)\tau) \right. \end{aligned}$$

$$\begin{aligned}
& + \int_0^\infty d\tau \langle n_b | \bar{\eta}(\tau, \mathbf{z}) j_\nu(0, \mathbf{y}) | n_a \rangle \exp(i(E' - E_{n_b} - p^0)\tau) \\
& - i \langle n_b | [\psi^\dagger(0, \mathbf{z}), j_\nu(0, \mathbf{y})] | n_a \rangle \{ \langle n_a | \bar{\psi}(0, \mathbf{x}_{N-1}) \cdots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle U_{p_i \mu_i}(\mathbf{z}) \\
& + \text{terms that are regular functions of } E' \text{ or } E \text{ if } E' \approx E_b^{(0)} \\
& \text{and } E \approx E_a^{(0)}.
\end{aligned} \tag{3.61}$$

We introduce the Green function $g(E', E, p^0)$ by

$$g_{\gamma_f, b; e_i, a}(E', E, p^0) = P_b^{(0)} \mathcal{G}_{\gamma_f; e_i}(E', E, p^0) \gamma_1^0 \cdots \gamma_{N-1}^0 P_a^{(0)}. \tag{3.62}$$

From equation (3.61) we have

$$\begin{aligned}
g_{\gamma_f, b; e_i, a}(E', E, p^0) &= \frac{1}{2\pi} \sum_{n_a=1}^{s_a} \sum_{n_b=1}^{s_b} \frac{\varphi_{n_b}}{E' - E_{n_b}} \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) \\
&\times \left\{ \int_0^\infty d\tau \langle n_b | j_\nu(\tau, \mathbf{y}) \bar{\eta}(0, \mathbf{z}) | n_a \rangle \exp(i(E - E_{n_b} + p^0)\tau) \right. \\
&+ \int_0^\infty d\tau \langle n_b | \bar{\eta}(\tau, \mathbf{z}) j_\nu(0, \mathbf{y}) | n_a \rangle \exp(i(E' - E_{n_b} - p^0)\tau) \\
&- i \langle n_b | [\psi^\dagger(0, \mathbf{z}), j_\nu(0, \mathbf{y})] | n_a \rangle \} U_{p_i \mu_i}(\mathbf{z}) \frac{\varphi_{n_a}^\dagger}{E - E_{n_a}} \\
&+ \text{terms that are regular functions of } E' \text{ or } E \text{ if } E' \approx E_b^{(0)} \\
&\text{and } E \approx E_a^{(0)},
\end{aligned} \tag{3.63}$$

where φ_{n_a} and φ_{n_b} are the wave functions of the initial and final states as defined above. Let the contours Γ_a and Γ_b surround the poles corresponding to the initial and final levels, respectively, and keep outside other singularities of $g_{\gamma_f, b; e_i, a}(E', E, p^0)$. This can be performed if the photon mass μ is chosen as indicated in section III(A). Taking into account the biorthogonality condition (2.71) and comparing (3.63) with (3.59) we obtain the desired formula [37]

$$S_{\gamma_f, b; e_i, a} = (Z_2 Z_3)^{-\frac{1}{2}} \delta(E_b + k_f^0 - E_a - p_i^0) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE v_b^\dagger g_{\gamma_f, b; e_i, a}(E', E, p_i^0) v_a, \tag{3.64}$$

where by a we imply one of the initial states and by b one of the final states under consideration. In the case of a single initial state (a) and a single final state (b), it yields

$$\begin{aligned}
S_{\gamma_f, b; e_i, a} &= (Z_2 Z_3)^{-\frac{1}{2}} \delta(E_b + k_f^0 - E_a - p_i^0) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}(E', E, p_i^0) \\
&\times \left[\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}(E) \right]^{-1/2} \left[\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}(E) \right]^{-1/2}.
\end{aligned} \tag{3.65}$$

For practical calculations by perturbation theory, it is convenient to express the Green function $g_{\gamma_f, b; e_i, a}$ in terms of the Fourier transform of the $(2N-1)$ -time Green function,

$$\begin{aligned}
& g_{\gamma_f, b; e_i, a}(E', E, p^0) \delta(E' + k^0 - E - p^0) \\
&= \frac{1}{\sqrt{N!(N-1)!}} \int_{-\infty}^\infty dp_1^0 \cdots dp_{N-1}^0 dp_1'^0 \cdots dp_N'^0 \delta(E - p_1^0 \cdots - p_{N-1}^0) \delta(E' - p_1'^0 \cdots - p_N'^0) \\
&\times P_b^{(0)} G_{\gamma_f; e_i}(p_1^0, \dots, p_N^0; k^0, p^0; p_1^0, \dots, p_{N-1}^0) \gamma_1^0 \cdots \gamma_{N-1}^0 P_a^{(0)},
\end{aligned} \tag{3.66}$$

where

$$\begin{aligned}
& G_{\gamma_f; e_i}((p_1^0, \mathbf{x}_1'), \dots, (p_N^0, \mathbf{x}_N'); k^0, p^0; (p_1^0, \mathbf{x}_1), \dots, (p_{N-1}^0, \mathbf{x}_{N-1})) \\
&= \left(\frac{2\pi}{i} \right)^2 \frac{1}{(2\pi)^{2N+1}} \int_{-\infty}^\infty dx_1^0 \cdots dx_{N-1}^0 dx_1'^0 \cdots dx_N'^0 \\
&\times \exp(ip_1'^0 x_1'^0 + \cdots + ip_N'^0 x_N'^0 - ip_1^0 x_1^0 - \cdots - ip_{N-1}^0 x_{N-1}^0)
\end{aligned}$$

$$\begin{aligned}
& \times \int d^4 y d^4 z \exp(ik^0 y^0 - ip^0 z^0) A_f^{\nu*}(\mathbf{y}) \\
& \times \langle 0 | T \psi(x'_1) \cdots \psi(x'_N) j_\nu(y) \bar{\psi}(z) \bar{\psi}(x_{N-1}) \cdots \bar{\psi}(x_1) | 0 \rangle \\
& \times (-i \overleftarrow{\not{\partial}}_z - m) U_{p_i \mu_i}(\mathbf{z}).
\end{aligned} \tag{3.67}$$

The Green function G_{γ_f, e_i} is constructed using the Wick theorem after the transition in (3.67) to the interaction representation. Since we have not included the interaction with V_C in the unperturbed Hamiltonian, the Feynman rules contain the free electron propagators and the vertices corresponding to the interaction with V_C . Summing over all the insertions of the vertices with V_C in the electron lines we replace the free electron propagators and wave functions by the electron propagators and wave functions in the Coulomb field, according to equations (3.53) and (3.54). The free wave function of the incident electron $U_{p_i \mu_i}(\mathbf{x})$ is replaced by the wave function $\psi_{p_i \mu_i(+)}(\mathbf{x})$ that can be determined from the equation

$$\psi_{p_i, \mu_i(+)} = U_{p_i \mu_i} + (p_i^0 - H_0(1 - i0))^{-1} V_C \psi_{p_i \mu_i(+)} . \tag{3.68}$$

The wave function $\psi_{p_i \mu_i(+)}(\mathbf{x})$ is a solution of the Dirac equation with the Coulomb potential $V_C(\mathbf{x})$ which satisfies the boundary condition [81]

$$\psi_{p \mu(+)}(\mathbf{x}) \sim U_{p \mu}(\mathbf{x}) + G^+(\mathbf{n}) \frac{\exp(i|\mathbf{p}||\mathbf{x}|)}{|\mathbf{x}|}, \quad |\mathbf{x}| \rightarrow \infty, \tag{3.69}$$

where $G^+(\mathbf{n})$ is a bispinor depending on $\mathbf{n} \equiv \mathbf{x}/|\mathbf{x}|$. The apparent expressions for $\psi_{p \mu(+)}(\mathbf{x})$ are given, e.g., in [82]. Thus, we again revert to the Furry picture. The Feynman rules for $G_{\gamma_f; e_i}$ differ from those for G_{γ_f} only by a replacement of one of the incoming electron propagators $\frac{i}{2\pi} S(\omega, \mathbf{x}, \mathbf{y})$ by the wave function $\psi_{p_i \mu_i(+)}(\mathbf{x})$.

In calculations by perturbation theory a problem appears which is caused by the fact that in zeroth approximation the energy of the $(N-1)$ -electron atom may coincide with the difference of the N -electron energy and the one-electron energy. As a result, some of the terms which are omitted in equation (3.61) are singular functions of E' and E if $E' \approx E_b^{(0)}$ and $E \approx E_a^{(0)}$. A special analysis of the complete spectral representation of the Green function $\mathcal{G}_{\gamma_f, e_i}$ shows that to eliminate these terms in calculations by formula (3.64) the following prescription should be used: integrate first over E and keep the point $E = E_a^{(0)} + (E' - E_b^{(0)})$ outside the contour Γ_a .

In Refs. [45,46] this method was used to derive formulas for the QED corrections to the radiative recombination of an electron with a bare nucleus and for the interelectronic-interaction corrections to the radiative recombination of an electron with a high- Z heliumlike atom. These formulas are presented in sections IV(D1), IV(D2). Here we consider another application of the method.

D. Radiative recombination of an electron with a high- Z hydrogenlike atom

As a practical application of the method, in this section we derive formulas for the radiative recombination of an electron with a high- Z hydrogenlike ion to zeroth and first orders in $1/Z$. We will assume that the final state of the heliumlike ion is described by a one-determinant wave function

$$u_b(\mathbf{x}_1, \mathbf{x}_2) = \frac{1}{\sqrt{2}} \sum_P (-1)^P \psi_{P b_1}(\mathbf{x}_1) \psi_{P b_2}(\mathbf{x}_2) \tag{3.70}$$

and the one-electron state $|b_1\rangle$ coincides with the initial state $|a\rangle$ of the hydrogenlike ion.

1. Zeroth order approximation

To zeroth order, the radiative recombination amplitude is described by the diagram shown in Fig. 34. The formula (3.65) gives

$$S_{\gamma_f, b; e_i, a}^{(0)} = \delta(E_b + k_f^0 - E_a - p_i^0) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(0)}(E', E, p_i^0), \tag{3.71}$$

where the superscript indicates the order in α . According to the definition (3.66) and the Feynman rules, we have

$$\begin{aligned}
& g_{\gamma_f, b; e_i, a}^{(0)}(E', E, p_i^0) \delta(E' + k^0 - E - p_i^0) \\
&= \frac{i}{2\pi} \sum_P (-1)^P \int_{-\infty}^{\infty} dp_1'^0 dp_2'^0 \delta(E' - p_1'^0 - p_2'^0) \frac{1}{p_2'^0 - \varepsilon_{Pb_2} + i0} \\
&\times \langle Pb_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \delta(p_2'^0 + k^0 - p_i^0) \frac{1}{p_1'^0 - \varepsilon_a + i0} \delta(p_1'^0 - E) \langle Pb_1 | a \rangle,
\end{aligned} \tag{3.72}$$

where $|p\rangle \equiv |p_i, \mu_i\rangle$ denotes the state vector of the incident electron. We obtain

$$g_{\gamma_f, b; e_i, a}^{(0)}(E', E, p_i^0) = \frac{i}{2\pi} \frac{1}{E' - E - \varepsilon_{b_2}} \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \frac{1}{E - \varepsilon_a}. \tag{3.73}$$

Following to the rule for the integration over E and E' formulated at the end of section III(C), we find

$$\oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(0)}(E', E, p_i^0) = -2\pi i \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle. \tag{3.74}$$

It yields

$$S_{\gamma_f, b; e_i, a}^{(0)} = -2\pi i \delta(E_b + k_f^0 - E_a - p_i^0) \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \tag{3.75}$$

or, according to the definition (3.2),

$$\tau_{\gamma_f, b; e_i, a}^{(0)} = -\langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle. \tag{3.76}$$

The corresponding cross section is

$$\frac{d\sigma^{(0)}}{d\Omega_f} = \frac{(2\pi)^4}{v_i} \mathbf{k}_f^2 |\tau_{\gamma_f, b; e_i, a}^{(0)}|^2, \tag{3.77}$$

where v_i is the velocity of the incident electron in the frame of the nucleus.

2. Interelectronic-interaction corrections of first order in $1/Z$

The interelectronic-interaction corrections of first order in $1/Z$ are defined by the diagrams shown in Fig. 35. In the order under consideration, the formula (3.65) gives

$$\begin{aligned}
S_{\gamma_f, b; e_i, a}^{(1)} &= \delta(E_b + k_f^0 - E_a - p_i^0) \left[\oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(1)}(E', E, p_i^0) \right. \\
&\quad \left. - \frac{1}{2} \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(0)}(E', E, p_i^0) \frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}^{(1)}(E) \right],
\end{aligned} \tag{3.78}$$

where $g_{bb}^{(1)}(E)$ is defined by the first-order interelectronic-interaction diagram (see Fig. 20). According to the definition (3.66) and the Feynman rules, we have

$$\begin{aligned}
& g_{\gamma_f, b; e_i, a}^{(1)}(E', E, p_i^0) \delta(E' + k^0 - E - p_i^0) \\
&= \left(\frac{i}{2\pi} \right)^2 \sum_P (-1)^P \int_{-\infty}^{\infty} dp_1'^0 dp_2'^0 \delta(E' - p_1'^0 - p_2'^0) \\
&\times \frac{1}{p_1'^0 - \varepsilon_{Pb_1} + i0} \frac{1}{p_2'^0 - \varepsilon_{Pb_2} + i0} \frac{1}{E - \varepsilon_a + i0} \int_{-\infty}^{\infty} dq^0 d\omega \\
&\times \sum_n \left[\langle Pb_1 Pb_2 | I(\omega) | an \rangle \delta(p_2'^0 + \omega - q^0) \delta(p_1'^0 - \omega - E) \right. \\
&\times \frac{1}{q^0 - \varepsilon_n(1 - i0)} \langle n | e\alpha_\nu A_f^{\nu*} | p \rangle \delta(q^0 + k^0 - p_i^0) \\
&\left. + \langle Pb_1 Pb_2 | I(\omega) | np \rangle \delta(p_2'^0 + \omega - p_i^0) \delta(p_1'^0 - \omega - q^0) \right]
\end{aligned}$$

$$\begin{aligned}
& \times \frac{1}{q^0 - \varepsilon_n(1 - i0)} \langle n | e\alpha_\nu A_f^{\nu*} | a \rangle \delta(q^0 + k^0 - E) \\
& + \langle Pb_2 | e\alpha_\nu A_f^{\nu*} | n \rangle \frac{1}{q^0 - \varepsilon_n(1 - i0)} \delta(p_2^0 + k^0 - q^0) \\
& \times \langle Pb_1 n | I(\omega) | ap \rangle \delta(q^0 + \omega - p_i^0) \delta(p_1^0 - \omega - E) \\
& + \langle Pb_1 | e\alpha_\nu A_f^{\nu*} | n \rangle \frac{1}{q^0 - \varepsilon_n(1 - i0)} \delta(p_1^0 + k^0 - q^0) \\
& \times \langle n Pb_2 | I(\omega) | ap \rangle \delta(p_2^0 + \omega - p_i^0) \delta(q^0 - \omega - E) \Big]. \tag{3.79}
\end{aligned}$$

We obtain

$$\begin{aligned}
g_{\gamma_f, b; e_i, a}^{(1)}(E', E, p_i^0) &= \left(\frac{i}{2\pi}\right)^2 \frac{1}{E' - E_b^{(0)}} \frac{1}{E - \varepsilon_a} \sum_P (-1)^P \sum_n \int_{-\infty}^{\infty} dp_1^0 \\
&\times \left(\frac{1}{p_1^0 - \varepsilon_{Pb_1} + i0} + \frac{1}{E' - p_1^0 - \varepsilon_{Pb_2} + i0} \right) \\
&\times \left[\langle Pb_1 Pb_2 | I(p_1^0 - E) | an \rangle \frac{1}{E' - E - \varepsilon_n(1 - i0)} \langle n | e\alpha_\nu A_f^{\nu*} | p \rangle \right. \\
&+ \langle Pb_2 | e\alpha_\nu A_f^{\nu*} | n \rangle \frac{1}{E + p_i^0 - p_1^0 - \varepsilon_n(1 - i0)} \\
&\times \langle Pb_1 n | I(p_1^0 - E) | ap \rangle \Big] \\
&+ \left(\frac{i}{2\pi}\right)^2 \frac{1}{E' - E_b^{(0)}} \frac{1}{E - \varepsilon_a} \sum_P (-1)^P \sum_n \int_{-\infty}^{\infty} dp_2^0 \\
&\times \left(\frac{1}{p_2^0 - \varepsilon_{Pb_2} + i0} + \frac{1}{E' - p_2^0 - \varepsilon_{Pb_1} + i0} \right) \\
&\times \left[\langle Pb_1 Pb_2 | I(p_i^0 - p_2^0) | np \rangle \frac{1}{E' - p_i^0 - \varepsilon_n(1 - i0)} \langle n | e\alpha_\nu A_f^{\nu*} | a \rangle \right. \\
&+ \langle Pb_1 | e\alpha_\nu A_f^{\nu*} | n \rangle \frac{1}{E + p_i^0 - p_2^0 - \varepsilon_n(1 - i0)} \\
&\times \langle n Pb_2 | I(p_i^0 - p_2^0) | ap \rangle \Big]. \tag{3.80}
\end{aligned}$$

The first term in equation (3.80) is conveniently divided into an irreducible ($\varepsilon_n \neq E_b^{(0)} - \varepsilon_a = \varepsilon_{b_2}$) and a reducible ($\varepsilon_n = \varepsilon_{b_2}$) part. Since we consider the case of a single final state described by the one-determinant wave function (3.70) with $|b_1\rangle = |a\rangle$, the condition $\varepsilon_n = \varepsilon_{b_2}$ means $|n\rangle = |b_2\rangle$ (otherwise $\langle Pb_1 Pb_2 | I(\omega) | an \rangle = 0$). Substituting (3.80) into (3.78), integrating over E and E' according to the rule formulated at the end of section III(C), and using the identity (2.127), we find for the irreducible contribution

$$\begin{aligned}
S_{\gamma_f, b; e_i, a}^{(1, \text{irred})} &= \delta(E_b + k_f^0 - E_a - p_i^0) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(1, \text{irred})}(E', E, p_i^0) \\
&= -2\pi i \delta(E_b + k_f^0 - E_a - p_i^0) \sum_P (-1)^P \\
&\times \left[\sum_n^{n \neq b_2} \langle Pb_1 Pb_2 | I(\varepsilon_{Pb_1} - \varepsilon_a) | an \rangle \frac{1}{\varepsilon_{b_2} - \varepsilon_n} \langle n | e\alpha_\nu A_f^{\nu*} | p \rangle \right. \\
&+ \sum_n \langle Pb_2 | e\alpha_\nu A_f^{\nu*} | n \rangle \frac{1}{\varepsilon_a + p_i^0 - \varepsilon_{Pb_1} - \varepsilon_n} \langle Pb_1 n | I(\varepsilon_{Pb_1} - \varepsilon_a) | ap \rangle \\
&+ \sum_n \langle Pb_1 Pb_2 | I(p_i^0 - \varepsilon_{Pb_2}) | np \rangle \frac{1}{E_b^{(0)} - p_i^0 - \varepsilon_n} \langle n | e\alpha_\nu A_f^{\nu*} | a \rangle \\
&\left. + \sum_n \langle Pb_1 | e\alpha_\nu A_f^{\nu*} | n \rangle \frac{1}{\varepsilon_a + p_i^0 - \varepsilon_{Pb_2} - \varepsilon_n} \langle n Pb_2 | I(p_i^0 - \varepsilon_{Pb_2}) | ap \rangle \right]. \tag{3.81}
\end{aligned}$$

For the reducible part we have

$$\begin{aligned}
& \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(1, \text{red})}(E', E, p_i^0) \\
&= 2\pi i \left(\frac{i}{2\pi}\right)^2 \oint_{\Gamma_b} dE' \sum_P (-1)^P \frac{1}{(E' - E_b^{(0)})^2} \int_{-\infty}^{\infty} dp_1'^0 \left(\frac{1}{p_1'^0 - \varepsilon_{Pb_1} + i0} \right. \\
&+ \left. \frac{1}{E' - p_1'^0 - \varepsilon_{Pb_2} + i0} \right) \langle Pb_1 Pb_2 | I(p_1'^0 - \varepsilon_a) | ab_2 \rangle \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \\
&= - \sum_P (-1)^P \int_{-\infty}^{\infty} d\omega \frac{1}{(\omega - \varepsilon_{Pb_1} - i0)^2} \langle Pb_1 Pb_2 | I(\omega - \varepsilon_a) | b_1 b_2 \rangle \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle.
\end{aligned} \tag{3.82}$$

The reducible contribution must be considered together with the second term in formula (3.78). Taking into account that

$$\begin{aligned}
\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}^{(1)}(E) &= -\frac{i}{2\pi} \left[2 \int_{-\infty}^{\infty} dp_1'^0 \frac{1}{(p_1'^0 - \varepsilon_{b_1} - i0)^2} \langle b_1 b_2 | I(p_1'^0 - \varepsilon_{b_1}) | b_1 b_2 \rangle \right. \\
&- \int_{-\infty}^{\infty} dp_1'^0 \frac{1}{(p_1'^0 - \varepsilon_{b_2} - i0)^2} \langle b_2 b_1 | I(p_1'^0 - \varepsilon_{b_1}) | b_1 b_2 \rangle \\
&- \left. \int_{-\infty}^{\infty} dp_1^0 \frac{1}{(p_1^0 - \varepsilon_{b_1} - i0)^2} \langle b_2 b_1 | I(p_1^0 - \varepsilon_{b_2}) | b_1 b_2 \rangle \right],
\end{aligned} \tag{3.83}$$

we find

$$\begin{aligned}
& -\frac{1}{2} \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; e_i, a}^{(0)}(E', E, p_i^0) \frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}^{(1)}(E) \\
&= \frac{1}{2} \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \left[2 \int_{-\infty}^{\infty} d\omega \frac{\langle b_1 b_2 | I(\omega) | b_1 b_2 \rangle}{(\omega - i0)^2} \right. \\
&- \left. \int_{-\infty}^{\infty} d\omega \langle b_2 b_1 | I(\omega) | b_1 b_2 \rangle \left(\frac{1}{(\omega - \Delta_b - i0)^2} + \frac{1}{(\omega + \Delta_b - i0)^2} \right) \right],
\end{aligned} \tag{3.84}$$

where $\Delta_b \equiv \varepsilon_{b_2} - \varepsilon_{b_1}$. Summing (3.82) and (3.84), we obtain for the total reducible contribution

$$\begin{aligned}
S_{\gamma_f, b; e_i, a}^{(1, \text{red})} &= -2\pi i \delta(E_b + k_f^0 - E_a - p_i^0) \frac{1}{2} \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \langle b_2 b_1 | I(\omega) | b_1 b_2 \rangle \\
&\times \left[\frac{1}{(\omega + \Delta_b + i0)^2} - \frac{1}{(\omega + \Delta_b - i0)^2} \right].
\end{aligned} \tag{3.85}$$

Using the identity

$$\frac{1}{(\omega + i0)^2} - \frac{1}{(\omega - i0)^2} = -\frac{2\pi}{i} \frac{d}{d\omega} \delta(\omega) \tag{3.86}$$

and integrating by parts, we obtain

$$S_{\gamma_f, b; e_i, a}^{(1, \text{red})} = 2\pi i \delta(E_b + k_f^0 - E_a - p_i^0) \frac{1}{2} \langle b_2 | e\alpha_\nu A_f^{\nu*} | p \rangle \langle b_2 b_1 | I'(\Delta_b) | b_1 b_2 \rangle. \tag{3.87}$$

In addition to the interelectronic-interaction correction derived in this subsection, we must take into account the contribution originating from changing the photon energy in the zeroth-order cross section (3.77) due to the interelectronic-interaction correction to the energy of the bound state b . It follows that the total interelectronic-interaction correction to the cross section of first order in $1/Z$ is given by

$$\frac{d\sigma^{(1)}}{d\Omega_f} = \frac{(2\pi)^4}{v_i} \mathbf{k}_f^2 2\text{Re} \left\{ \tau_{\gamma_f, b; e_i, a}^{(0)*} \tau_{\gamma_f, b; e_i, a}^{(1)} \right\} + \left[\frac{d\sigma^{(0)}}{d\Omega_f} \Big|_{k_f^0 = p_i^0 + \varepsilon_a - E_b} - \frac{d\sigma^{(0)}}{d\Omega_f} \Big|_{k_f^0 = p_i^0 + \varepsilon_a - E_b^{(0)}} \right]. \tag{3.88}$$

Here $\tau_{\gamma_f, b; e_i, a}^{(1)} = \tau_{\gamma_f, b; e_i, a}^{(1, \text{irred})} + \tau_{\gamma_f, b; e_i, a}^{(1, \text{red})}$ is the interelectronic-interaction correction given by equations (3.81) and (3.87) in accordance with the definition (3.2). E_b and $E_b^{(0)}$ are the energies of the bound state b with and without the interelectronic-interaction correction, respectively.

E. Photon scattering by an atom

Let us consider the scattering of a photon with momentum $\mathbf{k}_i$ and polarization $\epsilon_i = (0, \epsilon_i)$ by an atom which is initially in a state a . As a result of the scattering, a photon with momentum $\mathbf{k}_f$ and polarization $\epsilon_f = (0, \epsilon_f)$ is emitted and the atom is left in a state b . In this section we consider the non-resonant photon scattering. This means that the total initial energy of the system ($E_a + k_i^0$) is not close to any excited-state energy of the atom. The resonant photon scattering will be considered in detail in the next section.

According to the standard reduction technique (see, e.g., [28]), the amplitude of the process is

$$\begin{aligned} S_{\gamma_f, b; \gamma_i, a} &= \langle b | a_{\text{out}}(k_f, \epsilon_f) a_{\text{in}}^\dagger(k_i, \epsilon_i) | a \rangle \\ &= \text{Disconnected term} - Z_3^{-1} \int d^4 y d^4 z \frac{\epsilon_f^\nu \exp(ik_f \cdot y)}{\sqrt{2k_f^0 (2\pi)^3}} \\ &\quad \times \langle b | T j_\nu(y) j_\rho(z) | a \rangle \frac{\epsilon_i^\rho \exp(-ik_i \cdot z)}{\sqrt{2k_i^0 (2\pi)^3}}, \end{aligned} \quad (3.89)$$

where $|a\rangle$ and $|b\rangle$ are the vectors of the initial and final states in the Heisenberg representation; $k_f = (k_f^0, \mathbf{k}_f)$ and $k_i = (k_i^0, \mathbf{k}_i)$. The first term on the right-hand side of equation (3.89) corresponds to a non-scattering process, i.e., the photon does not interact with the atom. We are interested in the second term which describes the photon scattering by the atom. Let us designate this term by $S_{\gamma_f, b; \gamma_i, a}^{\text{scat}}$.

To derive the calculation formula for $S_{\gamma_f, b; \gamma_i, a}^{\text{scat}}$, in the scattering amplitude we isolate a term which describes the scattering of the photon by the Coulomb potential V_C . With this in mind, we write

$$\langle b | T j(y) j(z) | a \rangle = \langle b | [T j(y) j(z) - \langle 0 | T j(y) j(z) | 0 \rangle] | a \rangle + \delta_{ab} \langle 0 | T j(y) j(z) | 0 \rangle. \quad (3.90)$$

The second term on the right-hand side of this equation corresponds to the photon scattering by the Coulomb field and the first term describes the photon scattering by the electrons of the atom. To the first term, only diagrams contribute where the incident photon is connected to at least one external electron line. We denote this term by $S_{\gamma_f, b; \gamma_i, a}^{\text{con}}$. We have

$$\begin{aligned} S_{\gamma_f, b; \gamma_i, a}^{\text{con}} &= -Z_3^{-1} \int d^4 y d^4 z \frac{\epsilon_f^\nu \exp(ik_f \cdot y)}{\sqrt{2k_f^0 (2\pi)^3}} \langle b | [T j_\nu(y) j_\rho(z) \\ &\quad - \langle 0 | T j_\nu(y) j_\rho(z) | 0 \rangle] | a \rangle \frac{\epsilon_i^\rho \exp(-ik_i \cdot z)}{\sqrt{2k_i^0 (2\pi)^3}} \\ &= -Z_3^{-1} 2\pi \delta(E_b + k_f^0 - E_a - k_i^0) \int_{-\infty}^{\infty} dt \exp(ik_f^0 t) \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) \\ &\quad \times \langle b | [T j_\nu(t, \mathbf{y}) j_\rho(0, \mathbf{z}) - \langle 0 | T j_\nu(t, \mathbf{y}) j_\rho(0, \mathbf{z}) | 0 \rangle] | a \rangle A_i^\rho(\mathbf{z}). \end{aligned} \quad (3.91)$$

Here the second term is zero if $a \neq b$.

To construct the perturbation theory for the amplitude of the process, we introduce the Fourier transform of the corresponding two-time Green function

$$\begin{aligned} \mathcal{G}_{\gamma_f; \gamma_i}^{\text{con}}(E', E, k'^0; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &\delta(E' + k'^0 - E - k^0) \\ &= \left(\frac{i}{2\pi}\right)^2 \frac{1}{N!} \int_{-\infty}^{\infty} dx^0 dx'^0 \int d^4 y d^4 z \exp(iE' x'^0 + ik'^0 y^0 - iE x^0 - ik^0 z^0) \\ &\quad \times A_f^{\nu*}(\mathbf{y}) A_i^\rho(\mathbf{z}) \langle 0 | T \psi(x'^0, \mathbf{x}'_1) \cdots \psi(x'^0, \mathbf{x}'_N) [j_\nu(y) j_\rho(z) \\ &\quad - \langle 0 | T j_\nu(y) j_\rho(z) | 0 \rangle] \bar{\psi}(x^0, \mathbf{x}_N) \cdots \bar{\psi}(x^0, \mathbf{x}_1) | 0 \rangle. \end{aligned} \quad (3.92)$$

Introducing $t' = x'^0 - z^0$, $t = x^0 - z^0$, and $\tau = y^0 - z^0$ and integrating over z^0 , we obtain

$$\begin{aligned} \mathcal{G}_{\gamma_f; \gamma_i}^{\text{con}}(E', E, k'^0; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) &\delta(E' + k'^0 - E - k^0) \\ &= -\delta(E' + k'^0 - E - k^0) \frac{1}{2\pi} \frac{1}{N!} \int_{-\infty}^{\infty} dt dt' d\tau \exp(iE' t' + ik'^0 \tau - iE \tau) \\ &\quad \times \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) A_i^\rho(\mathbf{z}) \langle 0 | T \psi(t', \mathbf{x}'_1) \cdots \psi(t', \mathbf{x}'_N) [j_\nu(\tau, \mathbf{y}) j_\rho(0, \mathbf{z}) \\ &\quad - \langle 0 | T j_\nu(\tau, \mathbf{y}) j_\rho(0, \mathbf{z}) | 0 \rangle] \bar{\psi}(t, \mathbf{x}_N) \cdots \bar{\psi}(t, \mathbf{x}_1) | 0 \rangle. \end{aligned} \quad (3.93)$$

Integrating over t and t' we find

$$\begin{aligned}
& \mathcal{G}_{\gamma_f; \gamma_i}^{\text{con}}(E', E, k'^0; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) \\
&= -\frac{1}{2\pi} \frac{1}{N!} \sum_{n, m} \frac{i}{E' - E_n + i0} \frac{i}{E - E_m + i0} \int_{-\infty}^{\infty} d\tau \exp(ik'^0 \tau) \\
&\quad \times \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) A_i^{\rho}(\mathbf{z}) \{ \theta(\tau) \exp[i\tau(E' - E_n)] \langle 0 | T \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) | n \rangle \\
&\quad \times \langle n | [j_{\nu}(\tau, \mathbf{y}) j_{\rho}(0, \mathbf{z}) - \langle 0 | T j_{\nu}(\tau, \mathbf{y}) j_{\rho}(0, \mathbf{z}) | 0 \rangle] | m \rangle \\
&\quad \times \langle m | \bar{\psi}(0, \mathbf{x}_N) \cdots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle + \theta(-\tau) \exp[i\tau(E_m - E)] \\
&\quad \times \langle 0 | T \psi(0, \mathbf{x}'_1) \cdots \psi(0, \mathbf{x}'_N) | n \rangle \langle n | [j_{\rho}(0, \mathbf{z}) j_{\nu}(\tau, \mathbf{y}) \\
&\quad - \langle 0 | T j_{\nu}(\tau, \mathbf{y}) j_{\rho}(0, \mathbf{z}) | 0 \rangle] | m \rangle \langle m | \bar{\psi}(0, \mathbf{x}_N) \cdots \bar{\psi}(0, \mathbf{x}_1) | 0 \rangle + \cdots \} . \tag{3.94}
\end{aligned}$$

It can be shown that the terms which are omitted in equation (3.94) are regular functions of E' or E if $E' \approx E_b$ and $E \approx E_a$. As in the previous sections of this paper, we assume that in the zeroth approximation the initial and final states of the atom are degenerate in energy with some other states and use the same notations for these states as above. As usual, we also assume that a non-zero photon mass μ is introduced. We define the Green function $g_{\gamma_f, b; \gamma_i, a}^{\text{con}}(E', E, k'^0)$ by

$$g_{\gamma_f, b; \gamma_i, a}^{\text{con}}(E', E, k'^0) = P_b^{(0)} \mathcal{G}_{\gamma_f; \gamma_i}^{\text{con}}(E', E, k'^0) \gamma_1^0 \cdots \gamma_N^0 P_a^{(0)} . \tag{3.95}$$

From equation (3.94) we have

$$\begin{aligned}
g_{\gamma_f, b; \gamma_i, a}^{\text{con}}(E', E, k'^0) &= \frac{1}{2\pi} \sum_{n_a, n_b} \frac{\varphi_{n_b}}{E' - E_{n_b}} \int d\mathbf{y} d\mathbf{z} A_f^{\nu*}(\mathbf{y}) A_i^{\rho}(\mathbf{z}) \int_{-\infty}^{\infty} d\tau \exp(ik'^0 \tau) \\
&\quad \times \{ \theta(\tau) \exp[i\tau(E' - E_{n_b})] \langle n_b | [j_{\nu}(\tau, \mathbf{y}) j_{\rho}(0, \mathbf{z}) \\
&\quad - \langle 0 | T j_{\nu}(\tau, \mathbf{y}) j_{\rho}(0, \mathbf{z}) | 0 \rangle] | n_a \rangle + \theta(-\tau) \exp[i\tau(E_{n_a} - E)] \\
&\quad \times \langle n_b | [j_{\rho}(0, \mathbf{z}) j_{\nu}(\tau, \mathbf{y}) - \langle 0 | T j_{\nu}(\tau, \mathbf{y}) j_{\rho}(0, \mathbf{z}) | 0 \rangle] | n_a \rangle \} \frac{\varphi_{n_a}^{\dagger}}{E - E_{n_a}} \\
&\quad + \text{terms that are regular functions of } E' \text{ or } E \text{ if } E' \approx E_b^{(0)} \\
&\quad \text{and } E \approx E_a^{(0)} . \tag{3.96}
\end{aligned}$$

Taking into account the biorthogonality condition (2.71) and comparing (3.91) with (3.96), we obtain the desired formula [30]

$$S_{\gamma_f, b; \gamma_i, a}^{\text{con}} = Z_3^{-1} \delta(E_b + k_f^0 - E_a - k_i^0) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE v_b^{\dagger} g_{\gamma_f, b; \gamma_i, a}^{\text{con}}(E', E, k_f^0) v_a , \tag{3.97}$$

where we imply by a one of the initial states and by b one of the final states under consideration. In the case of a single initial state (a) and a single final state (b) it yields

$$\begin{aligned}
S_{\gamma_f, b; \gamma_i, a}^{\text{con}} &= Z_3^{-1} \delta(E_b + k_f^0 - E_a - k_i^0) \oint_{\Gamma_b} dE' \oint_{\Gamma_a} dE g_{\gamma_f, b; \gamma_i, a}^{\text{con}}(E', E, k_f^0) \\
&\quad \times \left[\frac{1}{2\pi i} \oint_{\Gamma_b} dE g_{bb}(E) \right]^{-1/2} \left[\frac{1}{2\pi i} \oint_{\Gamma_a} dE g_{aa}(E) \right]^{-1/2} . \tag{3.98}
\end{aligned}$$

The disconnected term describing the scattering of the photon by the Coulomb field is calculated by the formula

$$S_{\gamma_f, b; \gamma_i, a}^{\text{discon}} = -Z_3^{-1} \delta_{ab} \int d^4 y d^4 z \frac{\epsilon_f^{\nu} \exp(ik_f \cdot y)}{\sqrt{2k_f^0 (2\pi)^3}} \langle 0 | T j_{\nu}(y) j_{\rho}(z) | 0 \rangle \frac{\epsilon_i^{\rho} \exp(-ik_i \cdot z)}{\sqrt{2k_i^0 (2\pi)^3}} . \tag{3.99}$$

For practical calculations by perturbation theory it is convenient to express the Green function $g_{\gamma_f, b; \gamma_i, a}^{\text{con}}$ in terms of the Fourier transform of the $2N$ -time Green function,

$$\begin{aligned}
& g_{\gamma_f, b; \gamma_i, a}^{\text{con}}(E', E, k'^0) \delta(E' + k'^0 - E - k^0) \\
&= \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \delta(E - p_1^0 \cdots - p_N^0) \delta(E' - p_1'^0 \cdots - p_N'^0) \\
&\times P_b^{(0)} G_{\gamma_f; \gamma_i}^{\text{con}}(p_1^0, \dots, p_N^0; k'^0, k^0; p_1^0, \dots, p_N^0) \gamma_1^0 \cdots \gamma_N^0 P_a^{(0)}, \tag{3.100}
\end{aligned}$$

where

$$\begin{aligned}
& G_{\gamma_f; \gamma_i}^{\text{con}}((p_1^0, \mathbf{x}'_1), \dots, (p_N^0, \mathbf{x}'_N); k'^0, k^0; (p_1^0, \mathbf{x}_1), \dots, (p_N^0, \mathbf{x}_N)) \\
&= -\frac{1}{(2\pi)^{2N}} \int_{-\infty}^{\infty} dx_1^0 \cdots dx_N^0 dx_1'^0 \cdots dx_N'^0 \\
&\times \exp(ip_1'^0 x_1'^0 + \cdots + ip_N'^0 x_N'^0 - ip_1^0 x_1^0 - \cdots - ip_N^0 x_N^0) \\
&\times \int d^4 y d^4 z \exp(ik'^0 y^0 - ik^0 z^0) A_f^{\nu*}(\mathbf{y}) \\
&\times \langle 0 | T \psi(x'_1) \cdots \psi(x'_N) [j_\nu(y) j_\rho(z) - \langle 0 | T j_\nu(y) j_\rho(z) | 0 \rangle] \\
&\times \bar{\psi}(x_N) \cdots \bar{\psi}(x_1) | 0 \rangle A_i^\rho(\mathbf{z}). \tag{3.101}
\end{aligned}$$

The Green function $G_{\gamma_f, \gamma_i}^{\text{con}}$ is constructed using the Wick theorem after the transition in (3.101) to the interaction representation. The Feynman rules for $G_{\gamma_f, \gamma_i}^{\text{con}}$ differ from those for G_{γ_f} only by the presence of the incoming photon line which corresponds to the incident photon wave function $A_i^\rho(\mathbf{x})$.

F. Resonance scattering: Spectral line shape

The formulas for the scattering amplitudes derived above allow one to perform calculations by perturbation theory in the case where the total initial energy of the system is not close to the energy of an intermediate quasistationary state. This is the so-called non-resonant scattering. In the case of resonance scattering, when the initial energy of the system is close to an intermediate-state energy, the direct calculation by perturbation theory according to the formulas derived above leads to some singularities in the scattering amplitude. It means that those formulas cannot be directly applied to the resonance processes. In this section we formulate a method which allows one to calculate the resonance-scattering amplitudes. This method was worked out in Ref. [33]. Another approach, which is limited to the one-electron atom case, was previously developed in Ref. [83]. An attempt to describe a decay process within QED was undertaken in Ref. [84].

The photon scattering by an atom that is initially in its ground state a is now considered for the case of resonance $E_a + k_i^0 \sim E_d (d = 1, \dots, s)$, where E_a is the ground state energy of the atom, k_i^0 is the incident photon energy, and $E_d (d = 1, \dots, s)$ are the energies of intermediate atomic states which in zeroth order approximation are equal to the unperturbed energy $E_d^{(0)}$ of a degenerate level. We consider that, as a result of the scattering, a photon of energy $k_f^0 = k_i^0$ is emitted and the atom returns to its ground state a . The calculation of the photon scattering amplitude by the formula derived above leads to a singularity which is caused by the fact that in any finite order of perturbation theory one of the energy denominators of an intermediate Green function is close to zero. Therefore, in the calculation of the Green function $g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k'^0)$ we have to go beyond the finite-order approximation. With this in mind, let us represent $g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k'^0)$ as

$$\begin{aligned}
g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k'^0) &= \frac{i}{2\pi} g_a(E') R_{\gamma_f}^{(-)}(E', k'^0, E + k^0) \frac{i}{2\pi} g_d(E + k^0) R_{\gamma_i}^{(+)}(E + k^0, k^0, E) \frac{i}{2\pi} g_a(E) \\
&+ \Delta g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k'^0), \tag{3.102}
\end{aligned}$$

where g_a are g_d are the Green functions defined by equation (2.63) with the projectors $P_a^{(0)}$ and $P_d^{(0)}$, respectively, $k^0 = k'^0 + E' - E$, and $\Delta g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k'^0)$ is a part of the Green function $g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k'^0)$ which is regular at $E + k^0 \sim E_d (d = 1, \dots, s)$. The operators $R_{\gamma_f}^{(-)}$ and $R_{\gamma_i}^{(+)}$ are constructed by perturbation theory from equation (3.102) which must be considered as their definition. Taking into account equation (2.64) and using the formula (3.97), we obtain

$$\begin{aligned}
S_{\gamma_f, a; \gamma_i, a}^{\text{con}} &= Z_3^{-1} \delta(k_f^0 - k_i^0) \varphi_a^\dagger R_{\gamma_f}^{(-)}(E_a, k_f^0, E_a + k_i^0) \frac{i}{2\pi} g_d(E_a + k_i^0) R_{\gamma_i}^{(+)}(E_a + k_i^0, k_i^0, E_a) \varphi_a \\
&+ Z_3^{-1} \delta(k_f^0 - k_i^0) \oint_{\Gamma_a} dE' \oint_{\Gamma_a} dE v_a^\dagger \Delta g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k_f^0) v_a. \tag{3.103}
\end{aligned}$$

Consider now how the intermediate Green function $g_d(E_a + k_i^0)$ can be calculated. Let us introduce a quasipotential $V_d(E)$ by

$$g_d(E) = g_d^{(0)}(E) + g_d^{(0)}(E)V_d(E)g_d(E), \quad (3.104)$$

where $g_d^{(0)} = P_d^{(0)}/(E - E_d^{(0)})$. The quasipotential $V_d(E)$ is constructed by perturbation theory according to equation (3.104) which must be considered as its definition. This equation yields

$$\begin{aligned} V_d(E) &= [g_d^{(0)}(E)]^{-1} - [g_d(E)]^{-1} = [g_d^{(0)}(E)]^{-1} - [g_d^{(0)}(E) + g_d^{(1)}(E) + \dots]^{-1} \\ &= [g_d^{(0)}(E)]^{-1}g_d^{(1)}(E)[g_d^{(0)}(E)]^{-1} + \dots \end{aligned} \quad (3.105)$$

If the quasipotential $V_d(E)$ is constructed from equation (3.105) to a finite order of perturbation theory, the Green function g_d is determined by

$$g_d(E) = [E - E_d^{(0)} - V_d(E)]^{-1}. \quad (3.106)$$

The Green function $g_d(E)$ has poles on the second sheet of the Riemann surface, slightly below the right-hand real semiaxis (see Fig. 5), and has no singularities for real E when $E \sim E_d^{(0)}$. It means, in particular, that if we take the quasipotential at least to the lowest order approximation ($V(E) \approx V(E_d^{(0)})$), the Green function $g_d(E)$ calculated by equation (3.106) has no singularities at $E \sim E_d$ ($d = 1, \dots, s$), and, therefore, the calculation of the resonance-scattering amplitude by equation (3.103) will be correct. The calculation of $g_d(E)$ by equation (3.106) effectively corresponds to summing an infinite subsequence of Feynman diagrams.

For the calculation of $g_d(E)$ to the lowest order approximation it is convenient to introduce an operator $\mathcal{H}$ by

$$\mathcal{H} \equiv E_d^{(0)} + V_d(E_d^{(0)}). \quad (3.107)$$

The operator $\mathcal{H}$ is not Hermitian and has complex eigenvalues. We assume that $\mathcal{H}$ is a simple matrix, i.e., its eigenvectors form a complete basis in the space of the unperturbed d -states. We denote its eigenvalues by $\mathcal{E}_d = E_d - i\Gamma_d/2$, the right eigenvectors by $|d_R\rangle$, and the left eigenvectors by $\langle d_L|$. It means

$$\mathcal{H}|d_R\rangle = \mathcal{E}_d|d_R\rangle, \quad \langle d_L|\mathcal{H} = \langle d_L|\mathcal{E}_d. \quad (3.108)$$

It is convenient to normalize the vectors $|d_R\rangle$, $\langle d_L|$ by the condition

$$\langle d_L|d'_R\rangle = \delta_{dd'}. \quad (3.109)$$

They satisfy the completeness condition

$$\sum_{d=1}^s |d_R\rangle\langle d_L| = I. \quad (3.110)$$

For $g_d(E)$ we obtain

$$g_d(E) \approx (E - \mathcal{H})^{-1} = \sum_{d=1}^s \frac{|d_R\rangle\langle d_L|}{E - \mathcal{E}_d}. \quad (3.111)$$

In fact, due to T -invariance, $\mathcal{H}_{ik} = \mathcal{H}_{ki}$. For this reason the components of the vector $\langle d_L|$ can be chosen to be equal to the corresponding components of the vector $|d_R\rangle$. In other words, the components of the vector $|d_R\rangle$ are equal to the complex conjugated components of the vector $\langle d_L|$. If the d states have different quantum numbers, such as the total angular momentum or the parity, one finds $|d_R\rangle = |d_L\rangle \equiv |d\rangle$.

Substituting the lowest order approximation for $g_d(E)$ given by equation (3.111) into (3.103), we obtain in the resonance approximation

$$S_{\gamma_f, a; \gamma_i, a}^{\text{con}} \approx \frac{i}{2\pi} \delta(k_f^0 - k_i^0) \sum_{d=1}^s \frac{\langle a|R_{\gamma_f}^{(-)}|d_R\rangle\langle d_L|R_{\gamma_i}^{(+)}|a\rangle}{E_a + k_i^0 - E_d + i\Gamma_d/2}. \quad (3.112)$$

In the resonance approximation, it is sufficient to evaluate the operators $R_{\gamma_f}^{(-)}$, $R_{\gamma_i}^{(+)}$ to the lowest order of perturbation theory. They are determined directly from equation (3.102).

To demonstrate how the method works we consider below the resonance photon scattering on a one-electron atom. A more general case of a few-electron atom is considered in [33].

G. Resonance photon scattering by a one-electron atom

To study the resonance photon scattering by a one-electron atom, in the lowest order we must consider the diagram shown in Fig. 36. The contribution of this diagram to the Green function G_{γ_f, γ_i} is

$$\begin{aligned} G_{\gamma_f, \gamma_i}^{\text{con}}((E', \mathbf{x}'); k^0, k^0; (E, \mathbf{x})) &= \int d\mathbf{y} d\mathbf{z} \int_{-\infty}^{\infty} dp^0 \frac{i}{2\pi} S(E', \mathbf{x}', \mathbf{y}) \frac{2\pi}{i} e\gamma_\nu \delta(E' + k^0 - p^0) \\ &\quad \times A_f^{\nu*}(\mathbf{y}) \frac{i}{2\pi} S(p^0, \mathbf{y}, \mathbf{z}) \frac{2\pi}{i} e\gamma_\rho \delta(p^0 - k^0 - E) \\ &\quad \times A_i^\rho(\mathbf{z}) \frac{i}{2\pi} S(E, \mathbf{z}, \mathbf{x}). \end{aligned} \quad (3.113)$$

For $g_{\gamma_f, a; \gamma_i, a}^{\text{con}}$ we obtain

$$\begin{aligned} g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k^0) &= \frac{i}{2\pi} \frac{|a\rangle\langle a|}{E' - \varepsilon_a} \frac{2\pi}{i} e\alpha_\nu A_f^{\nu*} \\ &\quad \times \frac{i}{2\pi} \sum_n \frac{|n\rangle\langle n|}{E + k^0 - \varepsilon_n(1 - i0)} \frac{2\pi}{i} e\alpha_\rho A_i^\rho \frac{i}{2\pi} \frac{|a\rangle\langle a|}{E - \varepsilon_a}. \end{aligned} \quad (3.114)$$

We consider that $E_a + k_i^0 \sim E_d$ and, therefore, represent $g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k^0)$ as the sum of two terms

$$\begin{aligned} g_{\gamma_f, a; \gamma_i, a}^{\text{con}}(E', E, k^0) &= \frac{i}{2\pi} g_a^{(0)}(E') \frac{2\pi}{i} e\alpha_\nu A_f^{\nu*} \frac{i}{2\pi} g_d^{(0)}(E + k^0) \\ &\quad \times \frac{2\pi}{i} e\alpha_\rho A_i^\rho \frac{i}{2\pi} g_a^{(0)}(E) + \frac{i}{2\pi} g_a^{(0)}(E') \frac{2\pi}{i} e\alpha_\nu A_f^{\nu*} \\ &\quad \times \sum_{n, \varepsilon_n \neq \varepsilon_d} \frac{|n\rangle\langle n|}{E + k^0 - \varepsilon_n(1 - i0)} e\alpha_\rho A_i^\rho g_a^{(0)}(E). \end{aligned} \quad (3.115)$$

Comparing this equation with equation (3.102), we derive

$$R_{\gamma_f}^{(-)}(\mathbf{y}) = \frac{2\pi}{i} e\alpha_\nu A_f^{\nu*}(\mathbf{y}), \quad R_{\gamma_i}^{(+)}(\mathbf{z}) = \frac{2\pi}{i} e\alpha_\rho A_i^\rho(\mathbf{z}). \quad (3.116)$$

Let us derive now the quasipotential $V_d(E)$. To the lowest order of perturbation theory it is defined by the SE and VP diagrams (see Figs. 12,13). As was derived above (see section II(E)), the contribution of these diagrams to $g_d(E)$ is

$$g_d^{(1)}(E) = g_d^{(0)}(\Sigma_{\text{SE}}(E) + U_{\text{VP}})g_d^{(0)}. \quad (3.117)$$

Therefore,

$$V_d^{(1)}(E) = [g_d^{(0)}(E)]^{-1} g_d^{(1)}(E) [g_d^{(0)}(E)]^{-1} = P_d^{(0)}(\Sigma_{\text{SE}}(E) + U_{\text{VP}})P_d^{(0)}. \quad (3.118)$$

The operator $P_d^{(0)}\Sigma_{\text{SE}}(E)P_d^{(0)}$ contains a non-Hermitian part which is responsible for the imaginary part of the energy. The operator

$$\mathcal{H} = \varepsilon_d + V_d^{(1)}(\varepsilon_a + k_i^0) \quad (3.119)$$

acts in the s -dimensional space of the unperturbed states. In reality, due to the fact that the operators Σ_{SE} and U_{VP} do not mix states with different quantum numbers and among the degenerate one-electron states there are no states with the same quantum numbers, in the case under consideration the right eigenvectors of $\mathcal{H}$ coincide with the left eigenvectors, $|d_R\rangle = |d_L\rangle \equiv |d\rangle$. However, to keep a general form of the equations, below we will use the right and left eigenvectors. In the resonance approximation, the amplitude of the process is defined by equation (3.112), where the operators $R_{\gamma_f}^{(-)}$ and $R_{\gamma_i}^{(+)}$ are given by equations (3.116), $E_a = \varepsilon_a + \langle a | \Sigma_{\text{SE}}(\varepsilon_a) + U_{\text{VP}} | a \rangle$ is the ground state energy including the QED corrections of first order in α , E_d and $-\Gamma_d/2$ are the real and imaginary parts of an eigenvalue of $\mathcal{H}$.

For the differential cross section in the resonance approximation, we obtain

$$\begin{aligned}
d\sigma = (2\pi)^4 \delta(k_f^0 - k_i^0) & \left[\sum_{d=1}^s \frac{|\langle a | e\alpha_\nu A_f^{\nu*} | d_R \rangle \langle d_L | e\alpha_\rho A_i^\rho | a \rangle|^2}{(E_a + k_i^0 - E_d)^2 + \Gamma_d^2/4} \right. \\
& + 2\text{Re} \sum_{d < d'} \frac{\langle a | e\alpha_\nu A_f^{\nu*} | d_R \rangle \langle d_L | e\alpha_\rho A_i^\rho | a \rangle^2}{E_a + k_i^0 - E_d + i\Gamma_d/2} \\
& \left. \times \frac{\langle a | e\alpha_\nu A_f^{\nu*} | d'_R \rangle^* \langle d'_L | e\alpha_\rho A_i^\rho | a \rangle^*}{E_a + k_i^0 - E_{d'} - i\Gamma_{d'}/2} \right] d\mathbf{k}_f.
\end{aligned} \tag{3.120}$$

For the total cross section, using the optical theorem, we find

$$\begin{aligned}
\sigma_{\text{tot}} = 2(2\pi)^3 \sum_{d=1}^s & \left[\frac{\text{Re}(\langle a | e\alpha_\nu A_i^{\nu*} | d_R \rangle \langle d_L | e\alpha_\rho A_i^\rho | a \rangle) (\Gamma_d/2)}{(E_a + k_i^0 - E_d)^2 + \Gamma_d^2/4} \right. \\
& \left. + \frac{\text{Im}(\langle a | e\alpha_\nu A_i^{\nu*} | d_R \rangle \langle d_L | e\alpha_\rho A_i^\rho | a \rangle) (E_d - E_a - k_i^0)}{(E_a + k_i^0 - E_d)^2 + \Gamma_d^2/4} \right].
\end{aligned} \tag{3.121}$$

Let us discuss, for simplicity, the case $s = 2$. Only if the d levels have the same quantum numbers, the second term on the right-hand side of equation (3.121) is not equal to zero. In the opposite case, which takes place in the process under consideration, $|d_R\rangle = |d_L\rangle$ and, therefore,

$$\text{Im}(\langle a | e\alpha_\nu A_i^{\nu*} | d \rangle \langle d | e\alpha_\rho A_i^\rho | a \rangle) = 0. \tag{3.122}$$

It follows that the total cross section given by equation (3.121) is the sum of Lorentz-type terms. As to the differential cross section, the second term in equation (3.120) is not equal to zero even if the states $d = 1, 2$ have different quantum numbers.

Levels close to each other with identical quantum numbers can appear among doubly excited states of high- Z few-electron atoms [85]. As an example, we can consider the $(2s, 2s)_0$ and $(2p_{1/2}, 2p_{1/2})_0$ states of a heliumlike ion which can arise in the process of recombination of an electron with a hydrogenlike ion. A detailed theory of this process was given in Ref. [37]. The related numerical calculations were presented in Refs. [34,86]. The results obtained in these papers are discussed in section IV(D3).

IV. NUMERICAL EVALUATIONS OF QED AND INTERELECTRONIC-INTERACTION CORRECTIONS IN HEAVY IONS

A. Methods of numerical evaluations and renormalization procedure

In the previous sections we have demonstrated how formulas for the energy shifts and the transition and scattering amplitudes can be derived from the first principles of QED. These formulas usually contain infinite summations over intermediate electron states (summations over the bound states and integrations over the continuum). These sums are generally evaluated by using analytical expressions for the Dirac-Coulomb Green function [4,23,87–91] or by using relativistic finite basis set methods [92–97]. In some cases the summation can be performed analytically by employing the generalized virial relations for the Dirac equation [98].

Calculations of most QED corrections require the application of a renormalization procedure. To first order in α , one has to renormalize the self-energy and vacuum-polarization diagrams (Figs. 12, 13).

The renormalized expression for the SE correction is given by equation (2.102) which implies using the same covariant regularization for both terms on the right-hand side. For the numerical evaluation of this correction, it is convenient to analytically isolate the ultraviolet divergence in the $\langle a | \Sigma(\varepsilon_a) | a \rangle$ term and cancel it by the counterterm. This can be performed by expanding the Dirac-Coulomb Green function in terms of the free Dirac Green function,

$$\begin{aligned}
[\omega - H(1 - i0)]^{-1} &= [\omega - H_0(1 - i0)]^{-1} + [\omega - H_0(1 - i0)]^{-1} V_C [\omega - H_0(1 - i0)]^{-1} \\
&+ [\omega - H_0(1 - i0)]^{-1} V_C [\omega - H(1 - i0)]^{-1} V_C [\omega - H_0(1 - i0)]^{-1},
\end{aligned} \tag{4.1}$$

where $H_0 = \boldsymbol{\alpha} \cdot \mathbf{p} + \beta m$ is the free Dirac Hamiltonian. The three terms in equation (4.1) inserted into $\langle a | \Sigma(\varepsilon_a) | a \rangle$ divide the SE correction into zero-potential, one-potential, and many-potential terms, respectively. The ultraviolet divergences in the zero- and one-potential terms and in the counterterm can be cancelled analytically (see Refs. [99–101] for details). As to the many-potential term, it can easily be shown that it does not contain any ultraviolet

divergences. For an overview of other methods of carrying out mass renormalization in numerical calculations, we refer to [4].

The vacuum-polarization correction is determined by equation (2.106) with the VP potential defined by (2.107). The expression (2.107) is ultraviolet divergent. The simplest way to renormalize this expression is to expand the vacuum loop in powers of the Coulomb potential by employing equation (4.1). According to the Furry theorem, contributions of diagrams with odd numbers of vertices in a vacuum loop (with free Dirac propagators) are equal to zero. Therefore, the first non-zero contribution results from second term in the expansion (4.1). Only this contribution, which is called the Uehling term, is ultraviolet divergent. This term becomes finite by charge renormalization. The renormalized expression for the Uehling potential is given by

$$U_{\text{Uehl}}(r) = -\alpha Z \frac{2\alpha}{3\pi} \int_0^\infty dr' 4\pi r' \rho(r') \int_1^\infty dt \left(1 + \frac{1}{2t^2}\right) \frac{\sqrt{t^2 - 1}}{t^2} \times \frac{[\exp(-2m|r - r'|t) - \exp(-2m(r + r')t)]}{4mrt}, \quad (4.2)$$

where $|e|Z\rho(r)$ is the density of the nuclear charge distribution ($\int \rho(r)dr = 1$). The higher order (in V_C) terms are finite and their sum is called the Wichmann-Kroll correction [87]. However, the regularization is still required in the second non-zero term due to a spurious gauge dependent piece of the light-by-light scattering contribution. As was shown in [90,91,102], in the calculation of the vacuum polarization charge density based on the partial wave expansion of the Dirac-Coulomb Green function, the spurious term does not contribute if the sum over the angular momentum quantum number κ is restricted to a finite number of terms ($|\kappa| \leq K$). Thus the Wichmann-Kroll contribution can be calculated by summing up the partial differences between the full contribution and the Uehling term.

The renormalization procedures described above can be adopted for calculations of the self-energy and vacuum-polarization screening diagrams [39,40,42,43,103,104] as well as for the QED corrections to the hyperfine splitting and the bound-electron g factor [6,105–111].

B. Energy levels in heavy ions

1. Hydrogenlike ions

The relativistic energies of a hydrogenlike ion are determined by the Dirac equation (2.1). For the point-nucleus case, the Dirac equation can be solved analytically and the binding energy is given by

$$E_{nj} - mc^2 = -\frac{(\alpha Z)^2}{2\nu^2} \frac{2}{1 + (\alpha Z/\nu)^2 + \sqrt{1 + (\alpha Z/\nu)^2}} mc^2, \quad (4.3)$$

where $\nu = n + \sqrt{(j + 1/2)^2 - (\alpha Z)^2} - (j + 1/2)$, n is the principal quantum number, and j is the total angular momentum. To obtain the binding energy to higher accuracy, QED and nuclear effects must be taken into account.

The finite nuclear size correction is calculated by solving the Dirac equation with the potential of an extended nucleus and by taking the difference between the energies for the extended and point nucleus models. This can be performed numerically (see, e.g., [112,113]) or, with a good accuracy, analytically [114]. With a relative accuracy of $\sim 0.2\%$ for $Z = 1 - 100$, this correction is given by the following approximate formulas [114]

$$\Delta E_{ns} = \frac{(\alpha Z)^2}{10n} [1 + (\alpha Z)^2 f_{ns}(\alpha Z)] \left(2 \frac{\alpha Z}{n} \frac{R}{(\hbar/mc)}\right)^{2\gamma} mc^2, \quad (4.4)$$

$$\Delta E_{np_{1/2}} = \frac{(\alpha Z)^4}{40} \frac{n^2 - 1}{n^3} [1 + (\alpha Z)^2 f_{np_{1/2}}(\alpha Z)] \left(2 \frac{\alpha Z}{n} \frac{R}{(\hbar/mc)}\right)^{2\gamma} mc^2, \quad (4.5)$$

where $\gamma = \sqrt{1 - (\alpha Z)^2}$,

$$\begin{aligned} f_{1s}(\alpha Z) &= 1.380 - 0.162\alpha Z + 1.612(\alpha Z)^2, \\ f_{2s}(\alpha Z) &= 1.508 + 0.215\alpha Z + 1.332(\alpha Z)^2, \\ f_{2p_{1/2}}(\alpha Z) &= 1.615 + 4.319\alpha Z - 9.152(\alpha Z)^2 + 11.87(\alpha Z)^3, \end{aligned}$$

and R is an effective radius of the nuclear charge distribution defined by

$$R = \left\{ \frac{5}{3} \langle r^2 \rangle \left[1 - \frac{3}{4} (\alpha Z)^2 \left(\frac{3}{25} \frac{\langle r^4 \rangle}{\langle r^2 \rangle^2} - \frac{1}{7} \right) \right] \right\}^{1/2}. \quad (4.6)$$

For the Fermi model of the nuclear charge distribution,

$$\rho(r) = \frac{N}{1 + \exp[(r - c)/a]}, \quad (4.7)$$

it is possible to obtain with a very high precision

$$N = \frac{3}{4\pi c^3} \left(1 + \frac{\pi^2 a^2}{c^2} \right)^{-1}, \quad (4.8)$$

$$\langle r^2 \rangle = \frac{3}{5} c^2 + \frac{7}{5} \pi^2 a^2, \quad (4.9)$$

$$\langle r^4 \rangle = \frac{3}{7} c^4 + \frac{18}{7} \pi^2 a^2 c^2 + \frac{31}{7} \pi^4 a^4. \quad (4.10)$$

The expectation values for powers of r for a great variety of nuclear charge distribution models are given in Ref. [113].

Next the QED corrections of first order in α should be taken into account. To this order, the QED correction is defined by the self-energy and vacuum-polarization diagrams (Figs. 12,13). The energy shift from the self-energy diagram (Fig. 12) combined with the related mass counterterm (Fig. 14) is determined by equation (2.102). The self-energy correction for heavy ions was first evaluated by Desederio and Johnson [115] who employed a method suggested by Brown, Langer, and Schaefer [116]. Later, a more efficient method was developed by Mohr [89] who calculated this correction to a very high accuracy in a wide interval of Z . The method of the potential expansion of the bound-electron propagator for the calculation of the SE correction to all orders in αZ was developed by Snyderman [100] and numerically realized first by Blundell and Snyderman [117]. A very efficient procedure for the self-energy calculations which is closely related to the methods of Snyderman and Mohr was developed by Yerokhin and Shabaev [101]. An approach in which the ultraviolet divergences are removed by subtractions in coordinate space was worked out by Indelicato and Mohr [118]. The method of the partial-wave renormalization for the calculation of the first-order SE correction was developed by Persson, Lindgren, and Salomonson [119] and by Quiney and Grant [120]. To date, the most accurate calculations of the SE correction to all orders in αZ were performed by Mohr [89,121] and by Indelicato and Mohr [122] for the point nucleus case and by Mohr and Soff [123] for the extended nucleus case. This correction was comprehensively tabulated by Beier and co-workers [124] for finite nuclear radii. The highest accuracy for low- Z atoms was gained by Jentschura et al. [125,126].

The VP correction (Fig. 13) is the sum of the Uehling and Wichmann-Kroll contributions. The first contribution can easily be calculated using the expression (4.2) for the Uehling potential. Calculations of the Wichmann-Kroll contribution to all orders in αZ were performed first by Soff and Mohr [91] for the extended nucleus case and by Manakov et al. [127] for the point nucleus case. A comprehensive tabulation of this correction for extended nuclei was presented in Ref. [128]. The most accurate calculations for some specific ions were accomplished by Persson et al. [129].

The QED corrections of second order in α have not yet been calculated completely. Most VP-VP and SE-VP diagrams can be evaluated by the methods developed for the first-order SE and VP corrections (see [4,130] and references therein). The most difficult task consists in the evaluation of the SE-SE contribution. The simplest part of this contribution, the loop-after-loop diagram, was calculated by Mitrushenkov and co-workers [131] and by Mallampalli and Sapirstein [132] for high- Z ions. These numerical calculations were extended to low- Z atoms by Mallampalli and Sapirstein [133] and by Yerokhin [134]. Recently they were confirmed by analytical calculations of Yerokhin [135]. As to the residual SE-SE contribution, a specific part of it was evaluated by Mallampalli and Sapirstein [132] and an estimate of the complete SE-SE contribution is in progress [136]. For the current status of the corresponding calculations for low- Z atoms, we refer to [137,138].

The calculations of the corrections discussed above are based on quantum electrodynamics within the external field approximation. It means that in these calculations the nucleus is considered only as a source of the external Coulomb field V_C . The first step beyond this approximation consists in evaluating the nuclear recoil correction. This correction is given by the sum of the lower-order term (2.194) and the higher-order term (2.195). For the point nucleus case, an analytical calculation of the lower-order term employing the virial relations for the Dirac equation yields [50]

$$\Delta E_L = \frac{m^2 - \varepsilon_a^2}{2M}. \quad (4.11)$$

The higher-order term was numerically evaluated to all orders in αZ for point nuclei in Refs. [139,140]. The corresponding calculations for extended nuclei were carried out in Refs. [141,142]. In the case of hydrogen, the highest accuracy was gained in Ref. [143].

Finally, the nuclear polarization correction should be taken into account. This correction results from diagrams describing the interaction of the electron with the nucleus where the intermediate states of the nucleus are excited. It was evaluated by Plunien and Soff [144] and by Nefiodov et al. [145].

In Table 1 we present the individual contributions to the ground-state Lamb shift in $^{238}\text{U}^{91+}$. The uncertainty of the Dirac binding energy results from the uncertainty of the Rydberg constant [146]. As can be deduced from the table, the present status of the experimental precision on the ground-state Lamb shift in hydrogenlike uranium [147–149] provides a test of QED in first order in α on the level of about 5%.

2. Heliumlike ions

In heavy heliumlike ions, in addition to the one-electron contributions considered in the previous subsection, the two-electron corrections have to be taken into account. To lowest order in α , this correction is defined by the one-photon exchange diagram (Fig. 20). The calculation of this diagram causes no problem. To second order in α , we should account for the two-photon exchange diagrams (Fig. 21), the self-energy screening diagrams (Fig. 22), and the vacuum-polarization screening diagrams (Fig. 23). For the ground state of a heliumlike ion, the contribution of the two-photon exchange diagrams is defined by equations (2.134), (2.139), and (2.140). The corresponding expressions for the case of a single excited state were obtained in Ref. [36]. The derivation of the related formulas for the case of degenerate and quasidegenerate states by the TTGF method also causes no difficulties. The self-energy and vacuum-polarization screening contributions are given by expressions (2.153)–(2.160) (the renormalization of these expressions is considered in detail in Refs. [39,40,42,43]). For the case of quasidegenerate states, the corresponding formulas are derived by the TTGF method in Refs. [44,75].

The two-photon exchange contribution is conveniently divided into two parts. The first part is the one which can be derived from the Breit equation. For the ground state, it is well determined by the lowest-order terms of the αZ -expansion series [150],

$$\Delta E^{(\text{Breit})} = \alpha^2 [-0.15766638 - 0.6356(\alpha Z)^2] m. \quad (4.12)$$

The second part is the remaining one. Evaluated in Refs. [71,73], it was found to be much smaller than the first part. The self-energy and vacuum-polarization screening diagrams were evaluated in Refs. [39,40,103,104]. The related calculations for excited states of heliumlike ions were performed for the vacuum-polarization screening diagrams [44] and, in the case of non-mixed states, for the two-photon exchange diagrams [151].

Today, the theoretical uncertainty of the ground-state energy in heavy heliumlike ions is completely defined by the uncertainty of the one-electron contribution. In this connection, a direct measurement of the two-electron contribution to the ground-state energy in heliumlike ions performed in [152] turns out to be rather important. In Table 2 we present the individual two-electron contributions to the ground state energy in heliumlike bismuth. The two-photon exchange contribution is divided into two parts as described above. The three- and more photon contribution is evaluated within the Breit approximation by summing the Z^{-1} expansion terms for the ground state energy beginning from Z^{-3} [39]. For the zeroth order in αZ , the coefficients of this expansion are taken from Ref. [153] and for the second order in αZ from Ref. [150]. The uncertainty of this contribution results from QED corrections yet uncalculated. As one can see from the table, to test the second-order QED effects that result from the theory beyond the Breit approximation, the experimental precision has to be improved by an order of magnitude.

3. Lithiumlike ions

To date, the highest experimental accuracy was reached in Lamb-shift experiments of lithiumlike ions [154–157]. In these ions, in addition to the one- and two-electron contributions, the three-electron corrections have to be calculated. To second order in α , the three-electron contribution is determined by six diagrams that describe two-photon exchange involving all three electrons. One of these diagrams is shown in Fig. 37. In the case of one electron over the closed $(1s)^2$ shell, the unperturbed wave function is a one-determinant function,

$$u(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) = \frac{1}{\sqrt{3!}} \sum_P (-1)^P \psi_{Pa}(\mathbf{x}_1) \psi_{Pb}(\mathbf{x}_2) \psi_{Pv}(\mathbf{x}_3), \quad (4.13)$$

where a and b denote the core states with opposite signs of the angular momentum projection, v indicates the valence state. It can be shown that the derivation of the expressions for the one- and two-electron corrections in

three-electron atoms is easily reduced to the derivation in one- and two-electron atoms, respectively. As to the three-electron correction of second order in α , it can be derived by the TTGF method using the identity (2.129) and the general rules formulated in section II(E5). Such a derivation for the irreducible part yields [158]

$$\Delta E^{\text{irred}} = \sum_{PQ} (-1)^{P+Q} \sum_n' \frac{I_{PbPvnQv}(\varepsilon_{Qv} - \varepsilon_{Pv}) - I_{PanQaQb}(\varepsilon_{Pa} - \varepsilon_{Qa})}{\varepsilon_{Qa} + \varepsilon_{Qb} - \varepsilon_{Pa} - \varepsilon_n}, \quad (4.14)$$

where P and Q denote the permutations over the outgoing and incoming electron states, respectively; $I_{abcd}(\omega) \equiv \langle ab|I(\omega)|cd\rangle$. The prime at the sum indicates that terms with vanishing denominator have to be omitted in the summation. The reducible part of the three-electron contribution is [158]

$$\begin{aligned} \Delta E^{\text{red}} = & \sum_{\mu_a} [I'_{vaav}(\Delta)(I_{ab;ab} - I_{bv;bv}) \\ & + \frac{1}{2} I'_{bv\bar{v}a}(\Delta) I_{a\bar{v};bv} \frac{1}{2} I'_{a\bar{v}vb}(\Delta) I_{vb;\bar{v}a}], \end{aligned} \quad (4.15)$$

where $I_{ab;cd} = I_{abcd}(\varepsilon_b - \varepsilon_d) - I_{bacd}(\varepsilon_a - \varepsilon_d)$, $\Delta = \varepsilon_v - \varepsilon_a$, μ_a is the angular momentum projection of the a electron, $\mu_b = -\mu_a$, $\bar{v}$ is the valence electron with the opposite sign of the angular momentum projection. In derivation of equation (4.15) by the TTGF method, some terms containing the $\bar{v}$ electron have been cancelled with the corresponding terms from the reducible two-electron contribution of the two-photon exchange diagrams (see Ref. [158] for details). The accurate QED calculations of all the two- and three-electron corrections to the $2p_{1/2} - 2s$ transition energy up to second order in α were performed in Refs. [42,43,158]. Approximate evaluations of these corrections were previously considered in Refs. [159–164].

In Table 3 the individual contributions to the $2p_{1/2} - 2s$ transition energy in lithiumlike uranium are presented. The total theoretical value of the transition energy, $280.46(9) \pm 0.20$ eV, is in agreement with the related experimental value, $280.59(10)$ eV [154]. As can be seen from the table, the first-order QED contribution is -42.93 eV while the total second-order QED contribution beyond the Breit approximation amounts to 1.33 ± 0.20 eV. Comparing these values with the total theoretical and experimental uncertainties indicates that the present status of the theory for lithiumlike uranium provides a test of the QED effects of first order in α on the level of about 0.5% and of the QED effects of second order in α , which result from the theory beyond the Breit approximation, on the level of about 15%.

C. Hyperfine splitting and bound-electron g factor

1. Hyperfine splitting in hydrogenlike ions

The ground-state hyperfine splitting of a hydrogenlike ion is conveniently written as [166]

$$\begin{aligned} \Delta E_\mu = & \frac{4}{3} \alpha (\alpha Z)^3 \frac{\mu}{\mu_N} \frac{m}{m_p} \frac{2I+1}{2I} mc^2 \\ & \times \{A(\alpha Z)(1-\delta)(1-\varepsilon) + x_{\text{rad}}\}. \end{aligned} \quad (4.16)$$

Here m_p is the proton mass, μ is the nuclear magnetic moment, μ_N is the nuclear magneton, and I is the nuclear spin. $A(\alpha Z)$ denotes the relativistic factor

$$A(\alpha Z) = \frac{1}{\gamma(2\gamma-1)} = 1 + \frac{3}{2}(\alpha Z)^2 + \frac{17}{8}(\alpha Z)^4 + \dots, \quad (4.17)$$

where $\gamma = \sqrt{1 - (\alpha Z)^2}$. δ is the nuclear charge distribution correction, ε is the nuclear magnetization distribution correction (the Bohr-Weisskopf correction), and x_{rad} is the QED correction. The formulas for the first-order QED corrections to the hyperfine splitting are derived in the same way as formulas (2.115), (2.117) for the δV -SE corrections. For instance, the SE correction is simply determined by equations (2.115), (2.117), if δV is replaced by the hyperfine interaction operator

$$H_{\text{hfs}} = \frac{|e|}{4\pi} \frac{(\boldsymbol{\alpha} \cdot [\boldsymbol{\mu} \times \mathbf{r}])}{r^3} \quad (4.18)$$

and the electron wave functions are replaced by the wave functions of the whole (electron plus nucleus) atomic system. The most complete calculations of the QED and nuclear corrections to the hyperfine splitting were presented

in [107,110] (see also a recent review in [167]). Table 4 shows the individual contributions to the hyperfine splitting in hydrogenlike ions. The total theoretical values are compared with the experimental results obtained in [168–171]. The uncertainty of the theoretical predictions is mainly determined by the uncertainty of the Bohr-Weisskopf correction which was evaluated within the single-particle nuclear model [107]. This uncertainty should be considered only as an estimate of the order of magnitude of the real error. Except for Ho, the nuclear magnetic moments are taken from Ref. [172]. For Ho the value recommended in [173] is used. In case of ^{207}Pb , in Ref. [172] two values of the nuclear magnetic moment are given. One ($\mu = 0.592583(9)\mu_N$) was measured by the nuclear magnetic resonance (NMR) method [174] while another ($\mu = 0.58219(2)\mu_N$) results from an optical pumping (OP) experiment [175]. As was shown in [176], the OP value turns out to be very close to that obtained by NMR if it is corrected for an atomic effect (see the related discussion in [3]). Therefore, in Table 4 the NMR value for the nuclear magnetic moment of lead is used.

Taking into account that the theoretical uncertainties indicated in Table 4 should be considered only as an order of magnitude of the real errors, one can deduce that the total theoretical values are in reasonable agreement with the experimental ones. However, remeasurements of the nuclear magnetic moments by employing modern experimental technique and calculations of the Bohr-Weisskopf effect within many-particle nuclear models are required to promote investigations of the hyperfine splitting in hydrogenlike ions.

2. Hyperfine splitting in lithiumlike ions

The energy difference between the ground-state hyperfine splitting components of a lithiumlike ion is conveniently written as [38]

$$\Delta E_{(1s)^2 2s} = \frac{1}{6}\alpha(\alpha Z)^3 \frac{m}{m_p} \frac{\mu}{\mu_N} \frac{2I+1}{2I} mc^2 \left\{ [A^{(2s)}(\alpha Z)(1 - \delta^{(2s)})(1 - \varepsilon^{(2s)}) + x_{\text{rad}}^{(2s)}] + \frac{1}{Z}B(\alpha Z) + \frac{1}{Z^2}C(\alpha Z) + \dots \right\}. \quad (4.19)$$

Here $A^{(2s)}(\alpha Z)$ denotes the one-electron relativistic factor for the 2s state,

$$A^{(2s)}(\alpha Z) = \frac{2[2(1+\gamma) + \sqrt{2(1+\gamma)}]}{(1+\gamma)^2\gamma(4\gamma^2-1)} = 1 + \frac{17}{8}(\alpha Z)^2 + \frac{449}{128}(\alpha Z)^4 + \dots, \quad (4.20)$$

$\delta^{(2s)}$ is the one-electron nuclear charge distribution correction, $\varepsilon^{(2s)}$ is the one-electron nuclear magnetization distribution correction, and $x_{\text{rad}}^{(2s)}$ is the one-electron QED correction. The terms $B(\alpha Z)/Z$ and $C(\alpha Z)/Z^2$ describe the interelectronic-interaction corrections of first and second orders in $1/Z$, respectively. The first-order interelectronic-interaction correction is conveniently derived using the TTGF method with the closed $(1s)^2$ shell regarded as belonging to the vacuum (see section II(E3)). Such a derivation yields [38]

$$\begin{aligned} \Delta E_{FM_F I j} = & \sum_{M_I, m} \sum_{M'_I, m'} C_{IM'_I j m'}^{FM_F} C_{IM_I j m}^{FM_F} \chi_{IM_I} \sum_{\mu_c} \\ & \times \left\{ \sum_P (-1)^P \sum_{n \atop \varepsilon_n \neq \varepsilon_v}^{\varepsilon_n \neq \varepsilon_v} \frac{\langle P v' P c | I(\Delta_{Pcc}) | n c \rangle \langle n | H_{\text{hfs}} | v \rangle}{\varepsilon_v - \varepsilon_n} \right. \\ & + \sum_P (-1)^P \sum_{n \atop \varepsilon_n \neq \varepsilon_v}^{\varepsilon_n \neq \varepsilon_v} \frac{\langle v' | H_{\text{hfs}} | n \rangle \langle n c | I(\Delta_{Pcc}) | P v P c \rangle}{\varepsilon_v - \varepsilon_n} \\ & + \sum_P (-1)^P \sum_{n \atop \varepsilon_n \neq \varepsilon_c}^{\varepsilon_n \neq \varepsilon_c} \frac{\langle P v' P c | I(\Delta_{Pv'v}) | v n \rangle \langle n | H_{\text{hfs}} | c \rangle}{\varepsilon_c - \varepsilon_n} \\ & + \sum_P (-1)^P \sum_{n \atop \varepsilon_n \neq \varepsilon_c}^{\varepsilon_n \neq \varepsilon_c} \frac{\langle c | H_{\text{hfs}} | n \rangle \langle v' n | I(\Delta_{Pvv'}) | P v P c \rangle}{\varepsilon_c - \varepsilon_n} \\ & - \langle v' | H_{\text{hfs}} | v \rangle \langle c v | I'(\Delta_{vc}) | v c \rangle \\ & \left. + \sum_{\mu_{c'}} \langle c | H_{\text{hfs}} | c' \rangle \langle v' c' | I'(\Delta_{vc}) | c v \rangle \right\} \chi_{IM_I}, \quad (4.21) \end{aligned}$$

where F and M_F are the total angular momentum of the atom and its projection, v and v' are the valence states of electron with quantum numbers (jm) and $(j'm')$, respectively; $C_{IM_I jm}^{FM_F}$ is the Clebsch-Gordan coefficient, χ_{IM_I} is the nuclear wave function, c and c' denote the core states, μ_c indicates the angular momentum projection of the core electron, and $\Delta_{ab} = \varepsilon_a - \varepsilon_b$. Calculations of the nuclear, QED, and interelectronic-interaction corrections to the hyperfine splitting in heavy lithiumlike ions were performed in Refs. [38,109,167,177–183]. As for hydrogenlike ions, the uncertainty of the theoretical values is mainly determined by the uncertainty of the Bohr-Weisskopf effect evaluated within the single particle nuclear model. However, in Refs. [109,177] it was found that this uncertainty can be considerably reduced if the experimental value of the 1s hyperfine splitting in the corresponding hydrogenlike ion is known. The basic idea of this method is the following. It can be shown that, with a good precision, the ratio of the 2s–Bohr-Weisskopf correction to the 1s–Bohr-Weisskopf correction is a function of the atomic structure only and does not depend on the nuclear structure,

$$\frac{\varepsilon^{(2s)}}{\varepsilon^{(1s)}} = f(\alpha Z). \quad (4.22)$$

For $Z=83$, $f(\alpha Z)=1.078$ and, therefore, $\varepsilon^{(2s)} = 1.078 \varepsilon^{(1s)}$. The 1s–Bohr-Weisskopf correction can be found by the equation

$$\varepsilon^{(1s)} = \frac{\Delta E_{\text{Dirac}}^{(1s)} + \Delta E_{\text{QED}}^{(1s)} - \Delta E_{\text{exp}}^{(1s)}}{\Delta E_{\text{Dirac}}^{(1s)}}, \quad (4.23)$$

where $\Delta E_{\text{Dirac}}^{(1s)}$ is the relativistic value of the 1s hyperfine splitting including the nuclear charge distribution correction, $\Delta E_{\text{QED}}^{(1s)}$ is the 1s–QED correction, and $\Delta E_{\text{exp}}^{(1s)}$ is the experimental value of the 1s hyperfine splitting. For $Z=83$, this method predicts the ground-state hyperfine splitting in lithiumlike bismuth to be 0.7971(2) eV [179] (for comparison, the direct evaluation based on the single-particle nuclear model gives 0.800(4) eV). Recently, this value was confirmed by Sapirstein and Cheng [183] who obtained 0.79715(13) eV. Both theoretical values agree with the experimental one of 0.820(26) eV [155].

3. Bound-electron g factor

The bound-electron g factor in a hydrogenlike ion is defined by

$$g^{(e)} = -\frac{\langle JM_J | \mu_z^{(e)} | JM_J \rangle}{\mu_B M_J}, \quad (4.24)$$

where $\mu^{(e)}$ is the operator of the magnetic moment of electron, μ_B is the Bohr magneton, J is the total angular momentum of the electron, and M_J is its projection. For the ground state, a simple relativistic calculation based on the Dirac equation yields [184]

$$g_0 = 2 - (4/3)(1 - \sqrt{1 - (\alpha Z)^2}) \quad (4.25)$$

The QED and nuclear effects give some corrections to this value:

$$g^{(e)} = g_0 + \Delta g_{\text{QED}} + \Delta g_{\text{NS}}. \quad (4.26)$$

Calculation of the nuclear size correction causes no difficulties. For low- Z atoms, this correction can be found by a simple analytical formula [185]. The QED correction of first order in α was evaluated without expansion in αZ in Refs. [106,108,111] (see also [6]).

Direct measurements of the bound-electron g factor in hydrogenlike ions are presently being performed by a GSI - Universität Mainz collaboration [186–189]. To date, the experimental result obtained for hydrogenlike carbon (C^{5+}) [189] amounts to $g_{\text{exp}} = 2.001041596(5)$ and agrees with the theoretical predictions of Refs. [6,111], $g_{\text{theo}} = 2.001041591(7)$, and of Ref. [190], $g_{\text{theo}} = 2.001041590(2)$. The collaboration plans to extend these measurements to heavy ions.

Another possibility for investigations of the bound-electron g factor was recently proposed in [191]. In this work it was shown that the transition probability between the ground-state hyperfine splitting components of a hydrogenlike ion, including the first-order QED and nuclear corrections, is given by

$$w = \frac{\alpha}{3} \frac{\omega^3}{m^2} \frac{I}{2I+1} \left[g^{(e)} - g_I^{(n)} \frac{m}{m_p} \right]^2, \quad (4.27)$$

where ω is the transition frequency, $g^{(e)}$ is the bound-electron g factor defined above, and $g_I^{(n)}$ is the nuclear g factor (both g factors are defined to be positive). Formula (4.27) allows a simple calculation of the QED and nuclear corrections to the transition probability using the corresponding corrections to the bound-electron g factor. In [191] it was found that in the experimentally interesting cases of Pb and Bi, the QED and nuclear corrections increase the transition probability by about 0.3%. In [192] the lifetime of the upper hyperfine splitting component in $^{209}\text{Bi}^{82+}$ was measured to be $\tau_{\text{exp}} = 397.5(1.5) \mu\text{s}$. This result is in good agreement with the theoretical predictions of Ref. [191], $\tau_{\text{theo}} = 399.01(19) \mu\text{s}$, and of Ref. [193], $\tau_{\text{theo}} = 398.89 \mu\text{s}$. Using formula (4.27) and the experimental values of the hyperfine splitting and the transition probability in $^{209}\text{Bi}^{82+}$ [168,192], the experimental value of the bound-electron g factor in $^{209}\text{Bi}^{82+}$ is found to be 1.7343(33). The corresponding theoretical value is 1.7310. The individual contributions to the bound-electron g factor in $^{209}\text{Bi}^{82+}$ are given in Table 5. From this table, it is clear that the QED correction has to be included in order to obtain agreement between theory and experiment.

D. Radiative recombination of an electron with a heavy ion

In an energetic collision between a high- Z ion and a low- Z target atom, an electron may be captured by the projectile, while a simultaneously emitted photon carries away the excess energy and momentum. This process is denoted as radiative electron capture (REC). Since a loosely bound target electron can be considered as quasi-free, this process is essentially equivalent to radiative recombination (RR) or its time-reversed analogon, the photoelectric effect. The relativistic theory of REC was considered in detail in [82,194,195], and the results of this theory are in excellent agreement with experiments [196,197].

A systematic QED theory of the RR process is described in detail in sections III(D,E) of the present paper. In Refs. [34,37,86], this theory was employed to study the process of resonance recombination of an electron with a heavy hydrogenlike ion in the case of resonance with doubly excited $(2s, 2s)_0$, $(2p_{1/2}, 2p_{1/2})_0$, $(2s, 2p_{1/2})_{0,1}$ states of the corresponding heliumlike ion. Later, this theory was used to evaluate QED corrections to radiative recombination of an electron with a bare nucleus [45] and interelectronic-interaction corrections to radiative recombination of an electron with a heavy heliumlike ion [46]. The results of these investigations are briefly discussed below. More details can be found in the original papers [34,37,45,46,86].

1. QED corrections to the radiative recombination of an electron with a bare nucleus

We consider the radiative recombination of an electron with momentum $\mathbf{p}_i$ and polarization μ_i with a bare nucleus that is placed at the origin of the coordinate frame. This corresponds to the projectile system if we study the radiative recombination of a free target electron with a bare heavy projectile. To zeroth order in α , the cross section is

$$\frac{d\sigma^{(0)}}{d\Omega_f} = \frac{(2\pi)^4}{v_i} \mathbf{k}_f^2 |\tau^{(0)}|^2, \quad (4.28)$$

where

$$\tau^{(0)} = -\langle a | e\alpha^\nu A_{f,\nu}^* | p \rangle = \langle a | e\boldsymbol{\alpha} \cdot \mathbf{A}_f | p \rangle, \quad (4.29)$$

$|p\rangle \equiv |p_i, \mu_i\rangle$ is the wave function of the incident electron defined by equation (3.68), $p_i = (p_i^0, \mathbf{p}_i)$, $p_i^0 = \sqrt{\mathbf{p}_i^2 + m^2}$ is the energy of the incident electron, a is the final state of the one-electron atom, $k_f = (k_f^0, \mathbf{k}_f)$ with k_f^0 and $\mathbf{k}_f$ being the photon energy and momentum, respectively, v_i is the velocity of the incident electron in the nucleus frame.

The QED corrections of first order in α are defined by diagrams similar to those shown in Fig. 32. The direct calculation by the TTGF method yields for the self-energy correction to the amplitude of the process

$$\begin{aligned} \tau_{\text{SE}}^{(1)} = & - \left[\sum_n^{n \neq a} \frac{\langle a | \Sigma(\varepsilon_a) - \beta \delta m | n \rangle \langle n | e\alpha^\nu A_{f,\nu}^* | p \rangle}{\varepsilon_a - \varepsilon_n} \right. \\ & \left. + \frac{1}{2} \langle a | \Sigma'(\varepsilon_a) | a \rangle \langle a | e\alpha^\nu A_{f,\nu}^* | p \rangle \right] \end{aligned}$$

$$\begin{aligned}
& + \sum_n \frac{\langle a | e \alpha^\nu A_{f,\nu}^* | n \rangle \langle n | \Sigma(p_i^0) - \beta \delta m | p \rangle}{p_i^0 - \varepsilon_n(1 - i0)} \\
& + \int d\mathbf{z} e A_{f,\nu}^*(\mathbf{z}) \Lambda^\nu(\varepsilon_a, p_i^0, \mathbf{z}) + (Z_2^{-1/2} - 1) \langle a | e \alpha^\nu A_{f,\nu}^* | p \rangle \Big], \tag{4.30}
\end{aligned}$$

where the mass counterterm has been added and

$$\begin{aligned}
\Lambda^\nu(\varepsilon, p^0, \mathbf{z}) = e^2 \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \int d\mathbf{x} d\mathbf{y} \bar{\psi}_a(\mathbf{x}) \gamma^\rho S(\varepsilon - \omega, \mathbf{x}, \mathbf{z}) \gamma^\nu S(p^0 - \omega, \mathbf{z}, \mathbf{y}) \\
\times \gamma^\sigma D_{\rho\sigma}(\omega, \mathbf{x} - \mathbf{y}) \psi_{p_i \mu_i(+)}(\mathbf{y}). \tag{4.31}
\end{aligned}$$

A similar calculation of the VP correction gives

$$\begin{aligned}
\tau_{\text{VP}}^{(1)} = & - \left[\sum_n^{\neq a} \frac{\langle a | U_{\text{VP}} | n \rangle \langle n | e \alpha^\nu A_{f,\nu}^* | p \rangle}{\varepsilon_a - \varepsilon_n} \right. \\
& + \sum_n \frac{\langle a | e \alpha^\nu A_{f,\nu}^* | n \rangle \langle n | U_{\text{VP}} | p \rangle}{p_i^0 - \varepsilon_n(1 - i0)} \\
& \left. + \int d\mathbf{z} e A_{f,\nu}^*(\mathbf{z}) Q^\nu(k_f^0, \mathbf{z}) + (Z_3^{-1/2} - 1) \langle a | e \alpha^\nu A_{f,\nu}^* | p \rangle \right]. \tag{4.32}
\end{aligned}$$

Here $U_{\text{VP}}(\mathbf{x})$ is the vacuum-polarization potential defined by equation (2.107) and

$$\begin{aligned}
Q^\nu(k^0, \mathbf{z}) = & -e^2 \int d\mathbf{x} d\mathbf{y} \bar{\psi}_a(\mathbf{x}) \gamma^\rho \psi_{p_i \mu_i(+)}(\mathbf{x}) D_{\rho\sigma}(k^0, \mathbf{x} - \mathbf{y}) \\
& \times \frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega \text{Tr}[\gamma^\sigma S(\omega, \mathbf{y}, \mathbf{z}) \gamma^\nu S(\omega + k^0, \mathbf{z}, \mathbf{y})]. \tag{4.33}
\end{aligned}$$

In addition to these corrections, we have to take into account a contribution originating from changing the photon energy in the zeroth-order cross section (4.28) due to the QED correction to the energy of the bound state a . It follows that the total QED correction of first order in α to the cross section is given by

$$\frac{d\sigma_{\text{QED}}^{(1)}}{d\Omega_f} = \frac{(2\pi)^4}{v_i} \mathbf{k}_f^2 2\text{Re} \left\{ \tau^{(0)*} \tau_{\text{QED}}^{(1)} \right\} + \left[\frac{d\sigma^{(0)}}{d\Omega_f} \Big|_{k_f^0 = p_i^0 - E_a} - \frac{d\sigma^{(0)}}{d\Omega_f} \Big|_{k_f^0 = p_i^0 - \varepsilon_a} \right]. \tag{4.34}$$

Here $\tau_{\text{QED}}^{(1)} = \tau_{\text{SE}}^{(1)} + \tau_{\text{VP}}^{(1)}$ is the QED correction given by equations (4.30) and (4.32). E_a and ε_a are the energies of the bound state a with and without the QED correction, respectively.

The expressions (4.30) and (4.32) contain ultraviolet and infrared divergences. While the ultraviolet divergences can be eliminated by the standard renormalization procedure (see the related discussion in section III(B2)), the infrared divergences are more difficult to remove.

The infrared-divergent part of τ results from the region of small momenta of the virtual photon and is regularized by a non-zero photon mass μ . An evaluation of this part yields

$$\tau_{\text{infr}} = \tau^{(0)} \frac{\alpha}{\pi} \left[-\log(\mu/m) - \frac{1}{2} \log(\mu/m) \sqrt{1 + \frac{m^2}{\mathbf{p}_i^2}} \log \left(\frac{\sqrt{\mathbf{p}_i^2 + m^2} - |\mathbf{p}_i|}{\sqrt{\mathbf{p}_i^2 + m^2} + |\mathbf{p}_i|} \right) \right]. \tag{4.35}$$

The related contribution to the cross section is

$$\frac{d\sigma_{\text{infr}}}{d\Omega_f} = \frac{d\sigma^{(0)}}{d\Omega_f} \frac{\alpha}{\pi} \left[-2 \log(\mu/m) - \log(\mu/m) \sqrt{1 + \frac{m^2}{\mathbf{p}_i^2}} \log \left(\frac{\sqrt{\mathbf{p}_i^2 + m^2} - |\mathbf{p}_i|}{\sqrt{\mathbf{p}_i^2 + m^2} + |\mathbf{p}_i|} \right) \right], \tag{4.36}$$

where $\frac{d\sigma^{(0)}}{d\Omega_f}$ is the cross section in the zeroth-order approximation defined by equation (4.28). To cancel the infrared divergent contribution (4.36), we have to take into account that any experiment has a finite energy resolution ΔE . It means that any numbers of photons of the total energy less than ΔE can be emitted in the process. It follows that to find the total cross section in the order under consideration, we must include diagrams in which one photon of energy $k^0 = \sqrt{\mathbf{k}^2 + \mu^2} \leq \Delta E$ is emitted along with the emission of the photon with the energy $k_f^0 \approx p_i^0 - \varepsilon_a$ (we

assume $\Delta E \ll k_f^0$). These diagrams are shown in Fig. 38. Assuming that the energy resolution is sufficiently high ($\Delta E \ll k_f^0, m$), we retain only those contributions from the diagrams shown in Fig. 38 which dominate at $\Delta E \rightarrow 0$. Using the standard technique and omitting terms which approach zero at $\mu \rightarrow 0$, we obtain for this contribution

$$\begin{aligned} \frac{d\sigma_\gamma(\Delta E)}{d\Omega_f} &= \frac{d\sigma^{(0)}}{d\Omega_f} \frac{\alpha}{\pi} \left[1 - 2 \log 2 - 2 \log (\Delta E/\mu) - \sqrt{1 + \frac{m^2}{\mathbf{p}_i^2}} F(|\mathbf{p}_i|/p_i^0) \right. \\ &\quad \left. - \left(\frac{1}{2} + \log (\Delta E/\mu) \right) \sqrt{1 + \frac{m^2}{\mathbf{p}_i^2}} \log \left(\frac{\sqrt{\mathbf{p}_i^2 + m^2} - |\mathbf{p}_i|}{\sqrt{\mathbf{p}_i^2 + m^2} + |\mathbf{p}_i|} \right) \right], \end{aligned} \quad (4.37)$$

where

$$F(a) = \int_0^\infty dx \frac{x}{x^2 + 1} \log \left[\frac{(1+a)(\sqrt{x^2 + 1} - ax)}{(1-a)(\sqrt{x^2 + 1} + ax)} \right]. \quad (4.38)$$

We can see that the infrared divergent parts in the equations (4.36) and (4.37) cancel each other,

$$\begin{aligned} \frac{d\sigma_{\text{infr}}}{d\Omega_f} + \frac{d\sigma_\gamma(\Delta E)}{d\Omega_f} &= \frac{d\sigma^{(0)}}{d\Omega_f} \frac{\alpha}{\pi} \left[1 - 2 \log 2 - 2 \log (\Delta E/m) - \sqrt{1 + \frac{m^2}{\mathbf{p}_i^2}} F(|\mathbf{p}_i|/p_i^0) \right. \\ &\quad \left. - \left(\frac{1}{2} + \log (\Delta E/m) \right) \sqrt{1 + \frac{m^2}{\mathbf{p}_i^2}} \log \left(\frac{\sqrt{\mathbf{p}_i^2 + m^2} - |\mathbf{p}_i|}{\sqrt{\mathbf{p}_i^2 + m^2} + |\mathbf{p}_i|} \right) \right]. \end{aligned} \quad (4.39)$$

According to this equation, at a fixed energy of the incident electron, the QED correction depends on the photon-energy resolution ΔE and becomes infinite when $\Delta E \rightarrow 0$. It means that the validity of this equation is restricted by the condition $\frac{\alpha}{\pi} |\log (\Delta E/m)| \ll 1$. For extension of the theory beyond this limit it is necessary to include the radiative corrections of higher orders in α (see, e.g., [198,199]). It results in an "exponentiation" of the radiative corrections and removes the singularity for $\Delta E \rightarrow 0$.

In the derivation of the formulas (4.37), (4.39) it has been assumed that the incident electrons have a fixed energy. These formulas remain also valid in the case when the energy spread of the incident electrons is much smaller than the energy interval ΔE in which the photons are detected. However, this is not the case for the present REC experiments [196,197], where the energy spread of a quasi-free target electron is much larger than the finite photon-energy resolution. In that case, the QED correction to the total RR cross section depends on the form of the energy distribution of the target electron. Since the form of this distribution is not well determined, the only way to study the QED effects in REC processes is to investigate the cross section into a photon-energy interval which is chosen to be much larger than the effective energy spread of the quasi-free target electrons and much smaller than the energy of the emitted photon.

The Uehling part of the VP correction to the RR cross section and a part of the SE correction were numerically evaluated in Ref. [45]. Expressed in terms of the unperturbed cross section, the individual QED corrections to the total cross section for the radiative recombination into the K-shell of bare uranium are presented in the Table 6. The correction $\sigma_{\text{en}}^{(1)}$ results from changing the bound-state energy. It is determined by the term in the square brackets of equation (4.34). The correction $\sigma_{\text{bw}}^{(1)}$ corresponds to the irreducible part of the diagrams describing the first-order QED effect on the bound-state electron wave function. The correction $\sigma_{\text{cw}}^{(1)}$ results from the diagrams describing the QED effect on the continuum-state wave function.

As in bound-state QED, the Uehling approximation may be expected to account for a dominant part of the VP correction. As to the self-energy correction, we expect that the terms calculated in Ref. [45] give a reasonable estimate of the order of magnitude of the self-energy correction which is beyond the correction depending on the photon-energy interval ΔE (see equation (4.39)). The relative value of the last correction, which we denote by $\delta(\Delta E)$, is defined by

$$\delta(\Delta E) = \frac{\alpha}{\pi} \left[-2 + \frac{1}{\beta} \log \frac{1+\beta}{1-\beta} \right] \log \frac{\Delta E}{m}, \quad (4.40)$$

where $\beta = v_i/c$. The photon-energy interval has to be chosen in the range $\Gamma \ll \Delta E \ll k_f^0, m$, where Γ characterizes the energy spread of the incident electrons.

In the REC experiments which are performed at GSI, the effective electron-energy spread is determined by the momentum distribution of the quasi-free target electrons. The width of this spread in the projectile system increases with increasing impact energy. In the case of a N_2 gas target, which is presently being employed in the experiments,

the effective energy spread in the projectile (heavy ion) frame amounts to about 10-40 keV for the impact energy in the range 100-1000 MeV/u. This value will be considerably reduced in the experiments with a H₂ gas target which are under preparation. In the case of a H₂ gas target, to satisfy the conditions on ΔE given above we can choose ΔE to be 50 keV in the projectile frame for the impact energy 1 GeV/u. The corresponding photon-energy interval in the laboratory (gas-target) frame is determined according to the Lorentz transformation,

$$\Delta E_{\text{proj}} = \Delta E_{\text{lab}}(1 - \beta \cos \theta_{\text{lab}})/\sqrt{1 - \beta^2}. \quad (4.41)$$

At a fixed ΔE_{proj} , ΔE_{lab} as a function of the polar angle can be found from this equation. For the photon-energy interval chosen above, $\delta(\Delta E)$ amounts to -0.59% for an impact energy of 1 GeV/u. Adding the Uehling correction and the part of the SE correction presented in Table 6 to the correction $\delta(\Delta E)$, we find the QED correction to the total cross section amounts to -0.92% for an impact energy of 1 GeV/u. For a more accurate evaluation of this effect, complete calculations of all the SE corrections are required.

The results of the numerical evaluation of the differential cross section can be found in Ref. [45]. Here, we note only that the differential cross section at the backward direction vanishes at an impact energy close to 130 MeV/u. In particular, it results in a relatively large contribution of the QED correction to the backward cross section at an energy of 130 MeV/u. At this energy, the QED_{en+bw+ ΔE} correction is about 0.022 mbarn/sr while the zeroth-order cross section amounts only to 0.009 mbarn/sr.

2. Interelectronic-interaction effect on the radiative recombination of an electron with a heavy heliumlike ion

We consider the non-resonant radiative recombination of an electron with momentum $\mathbf{p}_i$ and polarization μ_i with a heavy heliumlike ion in the ground state which is placed at the origin of the coordinate frame. The final state of the system is a lithiumlike ion in the state $(1s)^2v$, where v denotes the valence state. This picture corresponds to the projectile system if we study the radiative recombination of a free target electron with a heavy heliumlike projectile.

To zeroth order, the amplitude of the process is given by

$$\tau^{(0)} = \langle v | e\boldsymbol{\alpha} \cdot \mathbf{A}_f^* | p \rangle, \quad (4.42)$$

where $|v\rangle$ denotes the wave function of the valence electron. The interelectronic-interaction correction of first order in $1/Z$ can easily be derived using the TTGF method with the closed $(1s)^2$ shell regarded as belonging to the vacuum. Such a derivation yields [46]

$$\tau_{\text{int}}^{(1)} = \sum_{l=1}^4 \tau_l^{\text{int}}, \quad (4.43)$$

where

$$\begin{aligned} \tau_1^{\text{int}} = & \sum_{\mu_c} \sum_P (-1)^P \sum_n^{\varepsilon_n \neq \varepsilon_v} \frac{\langle PvPc | I(\Delta_{Pc}) | nc \rangle \langle n | e\boldsymbol{\alpha} \cdot \mathbf{A}_f^* | p \rangle}{\varepsilon_v - \varepsilon_n} \\ & + \sum_{\mu_c} (-\frac{1}{2}) \langle cv | I'(\Delta_{vc}) | vc \rangle \langle v | e\boldsymbol{\alpha} \cdot \mathbf{A}_f^* | p \rangle, \end{aligned} \quad (4.44)$$

$$\tau_2^{\text{int}} = \sum_{\mu_c} \sum_P (-1)^P \sum_n \frac{\langle v | e\boldsymbol{\alpha} \cdot \mathbf{A}_f^* | n \rangle \langle nc | I(\Delta_{Pc}) | PpPc \rangle}{p_i^0 - \varepsilon_n(1 - i0)}, \quad (4.45)$$

$$\tau_3^{\text{int}} = \sum_{\mu_c} \sum_P (-1)^P \sum_n \frac{\langle PvPc | I(\Delta_{pPv}) | pn \rangle \langle n | e\boldsymbol{\alpha} \cdot \mathbf{A}_f^* | c \rangle}{\varepsilon_c - k_f^0 - \varepsilon_n(1 - i0)}, \quad (4.46)$$

$$\tau_4^{\text{int}} = \sum_{\mu_c} \sum_P (-1)^P \sum_n \frac{\langle c | e\boldsymbol{\alpha} \cdot \mathbf{A}_f^* | n \rangle \langle vn | I(\Delta_{pPv}) | PpPc \rangle}{\varepsilon_c + k_f^0 - \varepsilon_n(1 - i0)}. \quad (4.47)$$

Here $|v\rangle$ and $|c\rangle$ are the valence and core states, respectively, $p_i^0 = \sqrt{\mathbf{p}_i^2 + m^2}$ is the energy of the incident electron, $k_f^0 = p_i^0 - \varepsilon_v$ is the energy of the emitted photon, and μ_c indicates the angular momentum projection of the core electron. The expressions (4.44) – (4.47) represent the interelectronic-interaction corrections to the amplitude of the process. The corresponding corrections to the differential cross section are

$$\frac{d\sigma_l^{\text{int}}}{d\Omega_f} = \frac{(2\pi)^4}{v_i} \mathbf{k}_f^2 2\text{Re} \left[\tau^{(0)*} \tau_l^{\text{int}} \right] . \quad (4.48)$$

In addition to this, a contribution originating from a modification of the energy of the emitted photon in the zeroth-order cross section due to the interelectronic interaction should be taken into account which is given by

$$\frac{d\sigma_{\text{en}}^{\text{int}}}{d\Omega_f} = \frac{d\sigma^{(0)}}{d\Omega_f} \Big|_{k_f^0=p_i^0-E_v} - \frac{d\sigma^{(0)}}{d\Omega_f} \Big|_{k_f^0=p_i^0-\varepsilon_v} . \quad (4.49)$$

Here $E_v = \varepsilon_v + \Delta E_{\text{int}}^{(1)}$ is the energy of the valence electron including the first-order interelectronic-interaction correction,

$$\Delta E_{\text{int}}^{(1)} = \sum_{\mu_c} \sum_P (-1)^P \langle P v P c | I(\Delta_{P v v}) | v c \rangle . \quad (4.50)$$

The total interelectronic-interaction correction to the cross section in first order in $1/Z$ is

$$\frac{d\sigma_{\text{int}}^{(1)}}{d\Omega_f} = \frac{d\sigma_{\text{en}}^{\text{int}}}{d\Omega_f} + \sum_{l=1}^4 \frac{d\sigma_l^{\text{int}}}{d\Omega_f} . \quad (4.51)$$

The direct part of the corrections τ_1^{int} and τ_2^{int} can be accounted for by a modification of the incoming and outgoing electron wave functions by the screening potential

$$V_C(x) \rightarrow V_C(x) + V_{\text{scr}}(x) , \quad (4.52)$$

where

$$V_{\text{scr}}(x) = 2\alpha \left\{ \frac{1}{x} \int_0^x dy y^2 (g_{1s}^2(y) + f_{1s}^2(y)) + \int_x^\infty dy y (g_{1s}^2(y) + f_{1s}^2(y)) \right\} . \quad (4.53)$$

Here g_{1s} and f_{1s} are the upper and the lower components of the radial wave function of the ground state, respectively. The screening-potential approximation allows one to account for the dominant part of the interelectronic-interaction effect and is widely used in practical calculations.

Numerical results for the interelectronic-interaction correction to the total cross section of radiative recombination of an electron with heliumlike uranium are presented in Table 7 (for more extensive data we refer to [46]). The calculations are carried out for a capture into the $2s$, $2p_{1/2}$, and $2p_{3/2}$ states of lithiumlike uranium and for various projectile energies. The results of the rigorous relativistic treatment are compared with the calculations based on the screening-potential approximation. A deviation of the complete relativistic results from the screening-potential approximation is mainly determined by the term σ_4^{int} which strongly increases when p_i^0 comes close to the resonance condition ($p_i^0 - (\varepsilon_v - \varepsilon_c) \approx \varepsilon_n$). Numerical results for the differential cross section are given in Ref. [46].

3. Resonance recombination of an electron with a heavy hydrogenlike ion

We consider the process of recombination of an electron with a very heavy ($Z \sim 70 - 110$) hydrogenlike ion in its ground state for the case of resonance with doubly excited $(2s\ 2s)_0$, $(2p_{1/2}\ 2p_{1/2})_0$, $(2s\ 2p_{1/2})_{0,1}$ states of the corresponding heliumlike ion. We assume that the experimentally measured quantity is a part of the total cross section which corresponds to the emission of photons with an energy $\omega \approx E_d - E_r$, where E_d ($d = 1, 2, 3, 4$) are the energies of the doubly excited $(2s\ 2s)_0$, $(2p_{1/2}\ 2p_{1/2})_0$, $(2s\ 2p_{1/2})_{0,1}$ states, respectively, and E_r ($r = 1, 2, 3, 4$) are the energies of singly excited $(1s\ 2p_{1/2})_{0,1}$, $(1s\ 2s)_{0,1}$ states. The amplitude of this process is given by the sum of the amplitudes of the dielectronic recombination (DR) and radiative recombination (RR) processes

$$e^-(p_i^0) + X^{(Z-1)+}(1s) \rightarrow \begin{cases} X^{(Z-2)+}(d)^{**} \rightarrow X^{(Z-2)+}(r)^* + \gamma(\omega) \rightarrow \dots \\ X^{(Z-2)+}(r)^* + \gamma(\omega) \rightarrow \dots \end{cases}$$

Here p_i^0 is the energy of the incident electron. Among the doubly excited states $\{d\}$, there are states with identical quantum numbers ($(2s\ 2s)_0$, $(2p_{1/2}\ 2p_{1/2})_0$), while all the singly excited states $\{r\}$ have different quantum numbers. Main channels of the decay of $(1s\ 2s)_1$, $(1s\ 2p_{1/2})_1$ and $(1s\ 2s)_0$, $(1s\ 2p_{1/2})_0$ states are one- and two-photon transitions

to the ground state, respectively. Therefore, two- and three-photon processes give the dominant contribution to the cross section. The cross section of the process can be derived using the method described in section III(F). In the resonance approximation, such a derivation yields (see Ref. [37] for details)

$$\begin{aligned} \sigma(p_i^0) = & \frac{2\pi^2}{2(j'+1)} \frac{1}{|\mathbf{p}_i|^2} \sum_{j'lJM} \left\{ \sum_d \frac{W_{dd} |\langle d_L | \hat{I} | i_J \rangle|^2}{(E_{1s} + p_i^0 - E_d)^2 + \Gamma_d^2/4} \right. \\ & + 2 \operatorname{Re} \sum_{d < d'} \frac{W_{dd'} \langle d_L | \hat{I} | i_J \rangle \langle d'_L | \hat{I} | i_J \rangle^*}{(E_{1s} + p_i^0 - E_d + i\Gamma_d/2)(E_{1s} + p_i^0 - E_{d'} - i\Gamma_{d'}/2)} \\ & \left. + W_{i_J i_J} + 2 \operatorname{Re} \sum_d \frac{W_{di_J} \langle d_L | \hat{I} | i_J \rangle}{E_{1s} + p_i^0 - E_d + i\Gamma_d/2} \right\}, \end{aligned} \quad (4.54)$$

where $|i_J\rangle \equiv |JMn j' l' p_i^0 j l\rangle$, $(n j' l') = (1 \ 1/2 \ 0)$ are the quantum numbers of the $1s$ state of the hydrogenlike ion, $\hat{I} \equiv I(\varepsilon_{2s} - \varepsilon_{1s})$,

$$W_{dd'} = \sum_r W_{dd'}^{(r)}, \quad W_{di_J} = \sum_r W_{di_J}^{(r)}, \quad W_{i_J i_J} = \sum_r W_{i_J i_J}^{(r)}, \quad (4.55)$$

$$W_{dd'}^{(r)} = 2\pi\omega^2 \sum_{\epsilon} \int d\Omega_f \langle r | \hat{R}_\gamma | d_R \rangle \langle r | \hat{R}_\gamma | d'_R \rangle^*, \quad (4.56)$$

$$W_{di_J}^{(r)} = 2\pi\omega^2 \sum_{\epsilon} \int d\Omega_f \langle r | \hat{R}_\gamma | d_R \rangle \langle r | \hat{R}_\gamma | i_J \rangle^*, \quad (4.57)$$

$$W_{i_J i_J}^{(r)} = 2\pi\omega^2 \sum_{\epsilon} \int d\Omega_f |\langle r | \hat{R}_\gamma | i_J \rangle|^2, \quad (4.58)$$

$$\hat{R}_\gamma = - \sum_{n=1}^2 e\boldsymbol{\alpha} \cdot \mathbf{A}_f^*(\mathbf{x}_n). \quad (4.59)$$

The cross section (4.54) consists of four terms, $\sigma = \sum_{l=1}^4 \sigma_l$. The first two terms (σ_1, σ_2) correspond to the DR process, the third term (σ_3) corresponds to the RR process, and σ_4 describes the interference between the DR and RR processes. The term σ_2 is caused by the interference of the DR amplitudes on the levels with identical quantum numbers ($d, d' = 1, 2$). The magnitude of this term is determined by the overlap of the levels d, d' and can be characterized by the nonorthogonality integral $\langle d'_R | d_R \rangle$ which is connected with $W_{dd'}$ by the identity

$$W_{dd'} = i(\mathcal{E}_d - \mathcal{E}_{d'}^*) \langle d'_R | d_R \rangle. \quad (4.60)$$

Formula (4.54) was used in [34] for the numerical calculation of the cross section of the resonance recombination of an electron with hydrogenlike uranium. Later, a more accurate calculation was performed by Yerokhin [200]. According to these calculations, the ratio σ_2/σ which characterizes the overlap effect amounts up to 30% in the region between the maxima of the curve $\sigma(p_i^0)$. For the parameters of the overlapping levels it was found [34]

$$|\langle d_R | d'_R \rangle| = 0.180, \quad |W_{dd'}|/|E_d - E_{d'}| = 0.183. \quad (4.61)$$

Using (4.60) and the identity

$$W_{di} = i\langle i | (\hat{I} - \hat{I}^\dagger) | d_R \rangle, \quad (4.62)$$

the expression (4.54) can be transformed to the following (see Ref. [37] for details)

$$\begin{aligned} \sigma(p_i^0) = & \frac{2\pi^2}{2(2j'+1)} \frac{1}{|\mathbf{p}_i|^2} \sum_{j'lJM} \left\{ \sum_d \frac{\Gamma_d \operatorname{Re}(\langle i_J | \hat{I} | d_R \rangle \langle d_L | \hat{I} | i_J \rangle)}{(E_{1s} + p_i^0 - E_d)^2 + \frac{\Gamma_d^2}{4}} \right. \\ & - 2 \sum_d \frac{\operatorname{Im}(\langle i_J | \hat{I}^\dagger | d_R \rangle \langle d_L | \hat{I} | i_J \rangle) (E_{1s} + p_i^0 - E_d)}{(E_{1s} + p_i^0 - E_d)^2 + \frac{\Gamma_d^2}{4}} + W_{i_J i_J} \\ & \left. - 2 \sum_d \frac{\operatorname{Im}(\langle i_J | (\hat{I} - \hat{I}^\dagger) | d_R \rangle \langle d_L | \hat{I} | i_J \rangle) (E_{1s} + p_i^0 - E_d)}{(E_{1s} + p_i^0 - E_d)^2 + \frac{\Gamma_d^2}{4}} \right\}. \end{aligned} \quad (4.63)$$

This expression also consists of four terms $\sigma = \sum_{l=1}^4 \bar{\sigma}_l$. But, in contrast to (4.54), here the Breit-Wigner part of the cross section is completely contained in the first term ($\bar{\sigma}_1$) which is a sum of Lorentz-type terms. The terms $\bar{\sigma}_2, \bar{\sigma}_4$ again correspond to the interference of the DR amplitudes on the levels $d, d' = 1, 2$ and the interference of the DR and RR processes, respectively. But, unlike σ_2, σ_4 , the terms $\bar{\sigma}_2, \bar{\sigma}_4$ do not contain any admixture of Lorentz-type terms. They are given by sums of terms which are odd functions of $(E_{1s} + p_i^0 - E_d)$. (We should note that the admixture of Lorentz-type terms in σ_2, σ_4 is very small and hardly affects the values σ_2/σ and σ_4/σ .) The terms σ_2, σ_4 ($\bar{\sigma}_2, \bar{\sigma}_4$) lead to a deviation of the shape of the individual resonances from the Lorentz shape. This deviation can be characterized by the Low parameter [83]

$$\delta = \frac{\sigma(E_d - E_{1s} + \frac{\Gamma_d}{2}) - \sigma(E_d - E_{1s} - \frac{\Gamma_d}{2})}{\sigma(E_d - E_{1s})}. \quad (4.64)$$

Calculation of this parameter for $d = 2$ in the case of recombination with U^{91+} using the results of Refs. [34,200] yields $\delta \approx -0.17$. The contributions to δ from the terms $\bar{\sigma}_2$ and $\bar{\sigma}_4$ are equal -0.24 and 0.07 , respectively.

Summing the interference terms in (4.63), we obtain a compact formula for σ ,

$$\begin{aligned} \sigma(p_i^0) = & \frac{2\pi^2}{2(2j'+1)|\mathbf{p}_i|^2} \sum_{jLM} \left\{ \sum_d \frac{\Gamma_d \text{Re}(\langle i_J | \hat{I} | d_R \rangle \langle d_L | \hat{I} | i_J \rangle)}{(E_{1s} + p_i^0 - E_d)^2 + \frac{\Gamma_d^2}{4}} \right. \\ & \left. - 2 \sum_d \frac{\text{Im}(\langle i_J | \hat{I} | d_R \rangle \langle d_L | \hat{I} | i_J \rangle) (E_{1s} + p_i^0 - E_d)}{(E_{1s} + p_i^0 - E_d)^2 + \frac{\Gamma_d^2}{4}} + W_{i_J i_J} \right\}. \end{aligned} \quad (4.65)$$

The formulas (4.63) and (4.65) are more convenient for the numerical calculations than formula (4.54), since they do not require any calculation of the radiative amplitudes. These formulas were employed in Ref. [86] to calculate the resonance recombination of an electron with hydrogenlike lead.

V. CONCLUSION

In the present paper we have considered in detail the two-time Green function method for high-Z few-electron systems. This method allows one to formulate a perturbation theory for calculations of various physical quantities in a rigorous and systematic way in the framework of quantum electrodynamics. To demonstrate the efficiency of the method, we have derived formulas for QED and interelectronic-interaction corrections to the energy levels, transition and scattering amplitudes in one-, two-, and three-electron atoms.

For the last decade, the TTGF method was intensively employed in calculations of QED effects in heavy ions. An overview of these calculations was also given in the present paper. Details of the calculations and other applications of the method can be found in Refs. [35–46,158]. In particular, in Refs. [42,43] the vacuum-polarization and self-energy screening corrections to the energies of lithiumlike ions were calculated. The two-photon exchange diagrams for lithiumlike ions were evaluated in Ref. [158]. The second-order two-electron diagrams for quasidegenerate states of heliumlike ions are studied in [44,75]. In Ref. [35], the TTGF method was employed to construct an effective-energy operator for a high-Z few-electron atom. In Ref. [45], this method was used to evaluate the QED corrections to the radiative recombination of an electron with a bare nucleus. The interelectronic-interaction corrections to the radiative electron capture for a heliumlike ion were considered in Ref. [46]. In Ref. [80], the interelectronic-interaction corrections to the transition probabilities in heliumlike ions are derived.

Concluding, the two-time Green function method provides a uniform and very efficient approach for deriving QED and interelectronic-interaction corrections to energy levels, transition probabilities, and cross sections of scattering processes in high-Z few-electron atoms. Using an effective potential instead of the Coulomb potential of the nucleus allows one to extend this approach to many-electron atoms.

ACKNOWLEDGEMENTS

Many practical calculations by the TTGF method were carried out in collaboration with Anton Artemyev and Vladimir Yerokhin. Stimulating discussions with T. Beier, E.-O. Le Bigot, J. Eichler, P. Indelicato, U. Jentschura, S. Karshenboim, I. Lindgren, P. Mohr, G. Plunien, J. Sapirstein, G. Soff, and S. Zschocke are gratefully acknowledged. Valuable conversations with O. Andreev, D. Arbatsky, I. Bednyakov, M. Sysak, and O. Zhrebtssov are also acknowledged. This work was supported in part by RFBR (Grant No. 98-02-18350, Grant No. 98-02-0411, and Grant No.

01-02-17248), by the program "Russian Universities - Basic Research" (project No. 3930), by DFG (Grant No. 436 RUS 113/616/0-1), and by GSI.

APPENDIX A: QED IN THE HEISENBERG REPRESENTATION

In the Heisenberg representation, the basic equations of quantum electrodynamics in the presence of a classical time-independent field $A_{\text{cl}}^\nu(\mathbf{x})$ are

$$\begin{aligned} (i \not{\partial} - m - e \mathbf{A}_{\text{cl}}(\mathbf{x}))\psi(x) &= e \mathbf{A}(x)\psi(x) - \delta m \psi(x), \\ \square A_\nu(x) &= j_\nu(x), \end{aligned} \quad (\text{A1})$$

where $j_\nu(x) = (e/2)[\bar{\psi}(x)\gamma_\nu\psi(x)]$ is the electron-positron current operator. The state vectors in the Heisenberg representation are time-independent

$$\partial_t |\Phi\rangle = 0. \quad (\text{A2})$$

The physical state vectors have to obey a subsidiary condition

$$(\partial_\nu A^\nu(x))^{(+)}|\Phi\rangle = 0, \quad (\text{A3})$$

where $(\partial_\nu A^\nu(x))^{(+)}$ is the positive-frequency part of $\partial_\nu A^\nu(x)$. The Heisenberg operators $\psi(x)$, $\bar{\psi}(x)$, and $A^\nu(x)$ obey the same equal-time permutation relations as the corresponding free-field operators. However, in contrast to the free fields, the permutation relations for arbitrary times remain unknown. Due to the time-translation invariance, Heisenberg operators obey the following transformation equation

$$\exp(iHt)F(0, \mathbf{x})\exp(-iHt) = F(t, \mathbf{x}), \quad (\text{A4})$$

where H is the Hamiltonian of the system in the Heisenberg representation. For more details, see [21,56,81].

APPENDIX B: SINGULARITIES OF THE TWO-TIME GREEN FUNCTION IN A FINITE ORDER OF PERTURBATION THEORY

Let us investigate the singularities of the Green function $\mathcal{G}(E)$ in a finite order of perturbation theory. To m -th order in e , which corresponds to order $m/2$ in α , it is given by

$$\begin{aligned} &\mathcal{G}^{(m/2)}(E; \mathbf{x}'_1, \dots, \mathbf{x}'_N; \mathbf{x}_1, \dots, \mathbf{x}_N) \delta(E - E') \\ &= \frac{1}{2\pi i} \frac{1}{N!} \int_{-\infty}^{\infty} dt dt' \exp(iE't' - iEt) \frac{(-i)^m}{m!} e^m \int d^4 y_1 \cdots d^4 y_m \\ &\quad \times \langle 0 | T \psi_{\text{in}}(t', \mathbf{x}'_1) \cdots \psi_{\text{in}}(t', \mathbf{x}'_N) \bar{\psi}_{\text{in}}(t, \mathbf{x}_N) \cdots \bar{\psi}_{\text{in}}(t, \mathbf{x}_1) \\ &\quad \times \bar{\psi}_{\text{in}}(y_1) \gamma_\rho \psi_{\text{in}}(y_1) A_{\text{in}}^\rho(y_1) \cdots \bar{\psi}_{\text{in}}(y_m) \gamma_\sigma \psi_{\text{in}}(y_m) A_{\text{in}}^\sigma(y_m) | 0 \rangle_{\text{con}}, \end{aligned} \quad (\text{B1})$$

where the label "con" means that disconnected vacuum-vacuum subdiagrams must be omitted. For simplicity, we omit here the mass renormalization counterterm. The presence of this term would not change the consideration given below.

Let us consider the contribution of a diagram of m -th order in e . This diagram is defined by a certain order of contractions in equation (B1). The contractions between the electron-positron fields and between the photon fields give the propagators (2.10) and (2.11), respectively. In this Appendix we will use the following representation for these propagators

$$\begin{aligned} \langle 0 | T \psi_{\text{in}}(x) \bar{\psi}_{\text{in}}(y) | 0 \rangle &= \theta(x^0 - y^0) \sum_{\varepsilon_n > 0} \psi_n(\mathbf{x}) \bar{\psi}_n(\mathbf{y}) \exp[-i\varepsilon_n(x^0 - y^0)] \\ &\quad - \theta(y^0 - x^0) \sum_{\varepsilon_n < 0} \psi_n(\mathbf{x}) \bar{\psi}_n(\mathbf{y}) \exp[-i\varepsilon_n(x^0 - y^0)], \end{aligned} \quad (\text{B2})$$

$$\langle 0 | T A_{\text{in}}^\rho(x) A_{\text{in}}^\sigma(y) | 0 \rangle = -g^{\rho\sigma} \int \frac{d\mathbf{k}}{(2\pi)^3} \frac{\exp[-i\sqrt{\mathbf{k}^2 + \mu^2}|x^0 - y^0|] \exp[i\mathbf{k} \cdot (\mathbf{x} - \mathbf{y})]}{2\sqrt{\mathbf{k}^2 + \mu^2}}. \quad (\text{B3})$$

Here, by definition, $\theta(t) = (t + |t|)/(2t)$ for $t \neq 0$ and $\theta(0) = 1/2$, and a non-zero photon mass is introduced. Following [13,201], we will use the formalism of time-ordered diagrams [202,203] to investigate the singularities of $\mathcal{G}^{(m/2)}(E)$. Let us consider a certain order of the time variables,

$$y_{i_m}^0 > y_{i_{m-1}}^0 > \cdots y_{i_l}^0 > t' > y_{i_{l-1}}^0 > \cdots y_{i_s}^0 > t > y_{i_{s-1}}^0 > \cdots y_{i_1}^0$$

which defines a time-ordered version of the Feynman diagram. The contribution of the Feynman diagram is the sum of all time-ordered versions. Each time-ordered version is conveniently represented by a diagram in which the vertices are ordered upwards according to increasing time (see, for example, Fig. 39). According to equations (B2) and (B3), each electron propagator contains a sum over the whole electron-energy spectrum and each photon propagator contains an integral over the photon momentum. Let us place these sums and integrals in front of the expression for the time-ordered version of the Feynman diagram under consideration. Then, an electron line is characterized by an electron energy ε_n and, according to (B2), gives a factor (here we are only interested in time-dependent terms)

$$\exp[-i\varepsilon_n(y_i^0 - y_k^0)] = \exp[-i|\varepsilon_n|(y_i^0 - y_k^0)] \quad \text{for } \varepsilon_n > 0$$

and

$$-\exp[i\varepsilon_n(y_i^0 - y_k^0)] = -\exp[-i|\varepsilon_n|(y_i^0 - y_k^0)] \quad \text{for } \varepsilon_n < 0,$$

where in both cases we consider $y_i^0 > y_k^0$. A photon line gives a factor

$$\exp[-i\sqrt{\mathbf{k}^2 + \mu^2}(y_i^0 - y_k^0)],$$

where we consider again $y_i^0 > y_k^0$. Each time y_i^0 in the diagram is marked by a horizontal dashed line. These lines may intersect other electron and photon lines (see Fig. 39). For the point of intersection with an electron line, we introduce a factor $\exp(i|\varepsilon_n|y_i^0) \exp(-i|\varepsilon_n|y_i^0) = 1$, where ε_n is the energy of the intersected electron. For the point of intersection with a photon line we introduce a factor $\exp(i\sqrt{\mathbf{k}^2 + \mu^2}y_i^0) \exp(-i\sqrt{\mathbf{k}^2 + \mu^2}y_i^0) = 1$, where $\mathbf{k}$ is the momentum of the intersected photon. In addition, we represent the factor $\exp(iE't') \exp(-iEt)$ as

$$\begin{aligned} \exp(iE't') \exp(-iEt) &= \exp[i(E' - E)y_{i_m}^0] \exp[-i(E' - E)y_{i_m}^0] \cdots \\ &\quad \times \exp[i(E' - E)y_{i_l}^0] \exp[-i(E' - E)y_{i_l}^0] \\ &\quad \times \exp(iE't') \exp(-iEt') \exp(iEt') \exp(-iEy_{i_{l-1}}^0) \\ &\quad \times \exp(iEy_{i_{l-1}}^0) \exp(-iEy_{i_{l-2}}^0) \cdots \exp(iEy_{i_s}^0) \exp(-iEt). \end{aligned} \quad (\text{B4})$$

As a result of all these representations, the integral over times at fixed intermediate electron states (n) and photon momenta ($\mathbf{k}$) is

$$\begin{aligned} I_m &\equiv \int_{-\infty}^{\infty} dy_{i_m}^0 \int_{-\infty}^{y_{i_m}^0} dy_{i_{m-1}}^0 \cdots \int_{-\infty}^{y_{i_l}^0} dt' \int_{-\infty}^{t'} dy_{i_{l-1}}^0 \cdots \int_{-\infty}^{y_{i_s}^0} dt \int_{-\infty}^t dy_{i_{s-1}}^0 \cdots \int_{-\infty}^{y_{i_2}^0} dy_{i_1}^0 \\ &\quad \times \exp[i(E' - E)y_{i_m}^0] \exp[i(E - E' - \sum_{(m)} |\varepsilon_n| - \sum_{(m)} \sqrt{\mathbf{k}^2 + \mu^2})(y_{i_m}^0 - y_{i_{m-1}}^0)] \cdots \\ &\quad \times \exp[i(E - E' - \sum_{(l)} |\varepsilon_n| - \sum_{(l)} \sqrt{\mathbf{k}^2 + \mu^2})(y_{i_l}^0 - t')] \\ &\quad \times \exp[i(E - \sum_{(t')} |\varepsilon_n| - \sum_{(t')} \sqrt{\mathbf{k}^2 + \mu^2})(t' - y_{i_{l-1}}^0)] \cdots \\ &\quad \times \exp[i(E - \sum_{(s)} |\varepsilon_n| - \sum_{(s)} \sqrt{\mathbf{k}^2 + \mu^2})(y_{i_s}^0 - t)] \\ &\quad \times \exp[i(-\sum_{(t)} |\varepsilon_n| - \sum_{(t)} \sqrt{\mathbf{k}^2 + \mu^2})(t - y_{i_{s-1}}^0)] \cdots \\ &\quad \times \exp[i(-\sum_{(2)} |\varepsilon_n| - \sum_{(2)} \sqrt{\mathbf{k}^2 + \mu^2})(y_{i_2}^0 - y_{i_1}^0)]. \end{aligned} \quad (\text{B5})$$

Here $\sum_{(m)} |\varepsilon_n|$ denotes the sum of the electron energies from the electron lines which are sandwiched between the horizontal lines corresponding to the times $y_{i_m}^0$ and $y_{i_{m-1}}^0$. $\sum_{(m)} \sqrt{\mathbf{k}^2 + \mu^2}$ denotes the sum of the photon energies from the photon lines which are sandwiched between the horizontal lines corresponding to the times $y_{i_m}^0$ and $y_{i_{m-1}}^0$. Using the identity

$$\int_{-\infty}^0 dx \exp(-i\alpha x) = \frac{i}{\alpha + i0}, \quad (\text{B6})$$

we easily find

$$\begin{aligned}
I_m = 2\pi\delta(E - E') & \frac{i}{-\sum_{(m)} |\varepsilon_n| - \sum_{(m)} \sqrt{\mathbf{k}^2 + \mu^2}} \cdots \frac{i}{-\sum_{(l)} |\varepsilon_n| - \sum_{(l)} \sqrt{\mathbf{k}^2 + \mu^2}} \\
& \times \frac{i}{E - \sum_{(t')} |\varepsilon_n| - \sum_{(t')} \sqrt{\mathbf{k}^2 + \mu^2} + i0} \cdots \frac{i}{E - \sum_{(s)} |\varepsilon_n| - \sum_{(s)} \sqrt{\mathbf{k}^2 + \mu^2} + i0} \\
& \times \frac{i}{-\sum_{(t)} |\varepsilon_n| - \sum_{(t)} \sqrt{\mathbf{k}^2 + \mu^2}} \cdots \frac{i}{-\sum_{(2)} |\varepsilon_n| - \sum_{(2)} \sqrt{\mathbf{k}^2 + \mu^2}}. \tag{B7}
\end{aligned}$$

A similar calculation for $t' < t$ yields an expression which is obtained from (B7) by a replacement $E \rightarrow -E$ in all the denominators.

Because each photon line contracts two vertices, at least $m/2$ denominators in (B7) have to contain the photon-energy terms and therefore do not contribute to the singularities under consideration. It follows that $\mathcal{G}^{(m/2)}(E)$ has isolated poles of all orders till $m/2 + 1$ at the unperturbed positions of the bound state energies. The separation of these poles from the related cuts is provided by keeping a non-zero photon mass μ . As to the cuts starting from the lower energy levels, they can be turned down.

APPENDIX C: TWO-TIME GREEN FUNCTION IN TERMS OF THE FOURIER TRANSFORM OF THE $2N$ -TIME GREEN FUNCTION

To prove the equation (2.56) we have to show that

$$\begin{aligned}
\mathcal{G}(E)\delta(E - E') &= \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \\
&\quad \times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\
&\quad \times G(p_1^0, \dots, p_N^0; p_1'^0, \dots, p_N'^0), \tag{C1}
\end{aligned}$$

where the coordinate variables are omitted, for brevity. According to the definition of G (see equation (2.12)), equation (C1) is equivalent to

$$\begin{aligned}
\mathcal{G}(E)\delta(E - E') &= \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \\
&\quad \times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\
&\quad \times (2\pi)^{-2N} \int_{-\infty}^{\infty} dx_1^0 \cdots dx_N^0 dx_1'^0 \cdots dx_N'^0 \\
&\quad \times \exp(ip_1'^0 x_1'^0 + \cdots + ip_N'^0 x_N'^0 - ip_1^0 x_1^0 - \cdots - ip_N^0 x_N^0) \\
&\quad \times \langle 0 | T \psi(x_1') \cdots \psi(x_N') \bar{\psi}(x_N) \cdots \bar{\psi}(x_1) | 0 \rangle \\
&= (2\pi)^{-2N} \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dp_2^0 \cdots dp_N^0 dp_2'^0 \cdots dp_N'^0 \\
&\quad \times \int_{-\infty}^{\infty} dx_1^0 \cdots dx_N^0 dx_1'^0 \cdots dx_N'^0 \\
&\quad \times \exp[i(E' - p_2'^0 - \cdots - p_N'^0)x_1'^0 + ip_2'^0 x_2'^0 \cdots + ip_N'^0 x_N'^0] \\
&\quad \times \exp[-i(E - p_2^0 - \cdots - p_N^0)x_1^0 - ip_2^0 x_2^0 - \cdots - ip_N^0 x_N^0] \\
&\quad \times \langle 0 | T \psi(x_1') \cdots \psi(x_N') \bar{\psi}(x_N) \cdots \bar{\psi}(x_1) | 0 \rangle. \tag{C2}
\end{aligned}$$

Using the identity

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega \exp(i\omega x) = \delta(x), \tag{C3}$$

we obtain

$$\begin{aligned}
\mathcal{G}(E)\delta(E-E') &= (2\pi)^{-2} \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dx_1^0 \cdots dx_N^0 dx_1'^0 \cdots dx_N'^0 \delta(x_1^0 - x_2'^0) \cdots \delta(x_1'^0 - x_N^0) \\
&\quad \times \delta(x_1^0 - x_2^0) \cdots \delta(x_1^0 - x_N^0) \exp(iE'x_1'^0 - iEx_1^0) \\
&\quad \times \langle 0 | T \psi(x_1') \cdots \psi(x_N') \bar{\psi}(x_N) \cdots \bar{\psi}(x_1) | 0 \rangle \\
&= \frac{1}{2\pi i} \frac{1}{N!} \int_{-\infty}^{\infty} dx^0 dx'^0 \exp(iE'x'^0 - iEx^0) \\
&\quad \times \langle 0 | T \psi(x'^0, \mathbf{x}_1') \cdots \psi(x'^0, \mathbf{x}_N') \bar{\psi}(x^0, \mathbf{x}_N) \cdots \bar{\psi}(x^0, \mathbf{x}_1) | 0 \rangle.
\end{aligned} \tag{C4}$$

The last equation exactly coincides with the definition of $\mathcal{G}(E)$ given by (2.18).

APPENDIX D: MATRIX ELEMENTS OF THE TWO-TIME GREEN FUNCTION BETWEEN ONE-DETERMINANT WAVE FUNCTIONS

To derive the equation (2.61) we use the following two identities. First, if A is a symmetric operator in the coordinates of all electrons, we obtain (see, e.g., [204])

$$\begin{aligned}
A_{ik} \equiv \langle u_i | A | u_k \rangle &= \sum_P (-1)^P \psi_{P i_1}^*(\xi'_1) \cdots \psi_{P i_N}^*(\xi'_N) A(\xi'_1, \dots, \xi'_N; \xi_1, \dots, \xi_N) \\
&\quad \times \psi_{k_1}(\xi_1) \cdots \psi_{k_N}(\xi_N),
\end{aligned} \tag{D1}$$

where repeated variables $\{\xi\}$ imply integration (the integration over $\mathbf{x}$ and the summation over α) and $A(\xi'_1, \dots, \xi'_N; \xi_1, \dots, \xi_N)$ is the kernel of the operator A . Second, if the kernel of the operator A is represented in the form

$$A(\xi'_1, \dots, \xi'_N; \xi_1, \dots, \xi_N) = \sum_Q (-1)^Q a(\xi'_{Q_1}, \dots, \xi'_{Q_N}; \xi_1, \dots, \xi_N), \tag{D2}$$

we can find

$$\begin{aligned}
A_{ik} &= N! \sum_P (-1)^P \psi_{P i_1}^*(\xi'_1) \cdots \psi_{P i_N}^*(\xi'_N) a(\xi'_1, \dots, \xi'_N; \xi_1, \dots, \xi_N) \\
&\quad \times \psi_{k_1}(\xi_1) \cdots \psi_{k_N}(\xi_N).
\end{aligned} \tag{D3}$$

According to (2.60), we have

$$\begin{aligned}
\mathcal{G}(E) \gamma_1^0 \cdots \gamma_N^0 \delta(E-E') &= \frac{2\pi}{i} \frac{1}{N!} \int_{-\infty}^{\infty} dp_1^0 \cdots dp_N^0 dp_1'^0 \cdots dp_N'^0 \\
&\quad \times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\
&\quad \times \sum_P (-1)^P \hat{G}((p_{P1}^0, \xi'_{P1}), \dots, (p_{PN}^0, \xi'_{PN}); (p_1^0, \xi_1), \dots, (p_N^0, \xi_N)) \\
&= \frac{2\pi}{i} \frac{1}{N!} \sum_P (-1)^P \int_{-\infty}^{\infty} dp_1'^0 \cdots dp_N'^0 dp_1^0 \cdots dp_N^0 \\
&\quad \times \delta(E - p_1^0 - \cdots - p_N^0) \delta(E' - p_1'^0 - \cdots - p_N'^0) \\
&\quad \times \hat{G}((p_1^0, \xi'_{P1}), \dots, (p_N^0, \xi'_{PN}); (p_1^0, \xi_1), \dots, (p_N^0, \xi_N)) \\
&\equiv \sum_P (-1)^P \tilde{G}(\xi'_{P1}, \dots, \xi'_{PN}; \xi_1, \dots, \xi_N).
\end{aligned} \tag{D4}$$

Using (D1)-(D4) we easily obtain (2.61).

APPENDIX E: DOUBLE SPECTRAL REPRESENTATION FOR THE TWO-TIME GREEN FUNCTION DESCRIBING A TRANSITION PROCESS

Let us consider the function $G(E', E)$ defined as

$$G(E', E) = \int_{-\infty}^{\infty} dt dt' \exp(iE't' - iEt) \langle 0 | T A(t') B(0) C(t) | 0 \rangle. \quad (\text{E1})$$

Using the transformation rules for the Heisenberg operators and integrating over the time variables, we can derive the following double spectral representation for $G(E', E)$

$$\begin{aligned} G(E', E) = & - \int_{-\infty}^{\infty} dW' dW \left[\frac{K(W', W)}{(E' - W')(E - W)} + \frac{L(W', W)}{(E' + W')(E + W)} \right] \\ & + \int_{-\infty}^{\infty} dW' d\omega \left[\frac{M(W', \omega)}{(E' - W')(k^0 + \omega)} + \frac{N(W', \omega)}{(E' + W')(k^0 - \omega)} \right] \\ & - \int_{-\infty}^{\infty} d\omega dW \left[\frac{P(\omega, W)}{(k^0 - \omega)(E - W)} + \frac{Q(\omega, W)}{(k^0 + \omega)(E + W)} \right], \end{aligned} \quad (\text{E2})$$

where $k^0 = E - E'$,

$$K(W', W) = \sum_{n,m} \delta(W' - E_n) \delta(W - E_m) \langle 0 | A(0) | n \rangle \langle n | B(0) | m \rangle \langle m | C(0) | 0 \rangle, \quad (\text{E3})$$

$$L(W', W) = \sum_{n,m} \delta(W' - E_n) \delta(W - E_m) \langle 0 | C(0) | n \rangle \langle n | B(0) | m \rangle \langle m | A(0) | 0 \rangle, \quad (\text{E4})$$

$$M(W', \omega) = \sum_{n,m} \delta(W' - E_n) \delta(\omega - E_m) \langle 0 | A(0) | n \rangle \langle n | C(0) | m \rangle \langle m | B(0) | 0 \rangle, \quad (\text{E5})$$

$$N(W', \omega) = \sum_{n,m} \delta(W' - E_n) \delta(\omega - E_m) \langle 0 | B(0) | n \rangle \langle n | C(0) | m \rangle \langle m | A(0) | 0 \rangle, \quad (\text{E6})$$

$$P(\omega, W) = \sum_{n,m} \delta(\omega - E_n) \delta(W - E_m) \langle 0 | B(0) | n \rangle \langle n | A(0) | m \rangle \langle m | C(0) | 0 \rangle, \quad (\text{E7})$$

$$Q(\omega, W) = \sum_{n,m} \delta(\omega - E_n) \delta(W - E_m) \langle 0 | C(0) | n \rangle \langle n | A(0) | m \rangle \langle m | B(0) | 0 \rangle. \quad (\text{E8})$$

- [1] H.F. Beyer, H.-J. Kluge, V.P. Shevelko, X-ray Radiation of Highly Charged Ions, Springer, Berlin, 1997.
- [2] H.F. Beyer, V.P. Shevelko (eds.), Atomic Physics with Heavy Ions, Springer, Berlin, 1999.
- [3] V.M. Shabaev, A.N. Artemyev, V.A. Yerokhin, Phys. Scr. T 86 (2000) 7.
- [4] P.J. Mohr, G. Plumien, G. Soff, Phys. Rep. 293 (1998) 227.
- [5] J. Sapirstein, in: D.H.E. Dubin, D. Schneider (eds.), Trapped Charged Particles and Fundamental Physics, American Institute of Physics Conference Proceedings 457, 1999, p. 3.
- [6] T. Beier, Phys. Rep., 339 (2000) 79.
- [7] I. Lindgren, Phys. Scr. T 80 (1999) 131.
- [8] M. Gell-Mann, F. Low, 1951 Phys. Rev. 84 (1951) 350.
- [9] J. Sucher, Phys. Rev. 107 (1957) 1448.
- [10] L.N. Labzowsky, Zh. Eksp. Teor. Fiz. 59 (1970) 168 [Sov. Phys. JETP 32 (1970) 94].
- [11] M.A. Braun, A.D. Gurchumelia, Teor. Mat. Fiz. 45 (1980) 199 [Theor. Math. Phys. 45 (1980) N 2].
- [12] Yu.Yu. Dmitriev, G.L. Klimchitskaya, L.N. Labzowsky, Relativistic Effects in Spectra of Atomic Systems, Energoatomizdat, Moscow, 1984.
- [13] M.A. Braun, A.D. Gurchumelia, U.I. Safronova, Relativistic Atom Theory, Nauka, Moscow, 1984.
- [14] P.J. Mohr, in: W.R. Johnson, P.J. Mohr, J. Sucher (eds.), Relativistic, Quantum Electrodynamics and Weak Interaction Effects in Atoms, American Institute of Physics Conference Series N 189, 1989, p. 47.
- [15] I. Lindgren, in: W.R. Johnson, P.J. Mohr, J. Sucher (eds.), Relativistic, Quantum Electrodynamics and Weak Interaction Effects in Atoms, American Institute of Physics Conference Series N 189, 1989, p. 371.
- [16] L. Labzowsky, G. Klimchitskaya, Yu. Dmitriev, Relativistic Effects in Spectra of Atomic Systems, IOP Publishing, Bristol, 1993.

- [17] J. Sapirstein, Rev. Mod. Phys. 70 (1998) 55.
- [18] F.J. Dyson, Phys. Rev. 75 (1949) 486, 1736.
- [19] E.E. Salpeter, H.A. Bethe, Phys. Rev. 84 (1951) 1232.
- [20] J. Hubbard, Proc. Roy. Soc. A 240 (1957) 539.
- [21] S.S. Schweber, An Introduction to Relativistic Quantum Field Theory, Harper & Row, New York, 1961.
- [22] A.N. Vasil'ev, A.L. Kitanin, Teor. Mat. Fiz. 24 (1975) 219 [Theor. Math. Phys. 24 (1975) N 2].
- [23] S.A. Zapryagaev, N.L. Manakov, V.G. Pal'chikov, Theory of One- and Two-Electron Multicharged Ions, Energoatomizdat, Moscow, 1985.
- [24] J. Sapirstein, Phys. Scr. 36 (1987) 801.
- [25] Yu.Yu. Dmitriev, T.A. Fedorova, Phys. Lett. A 225 (1997) 296.
- [26] Yu.Yu. Dmitriev, T.A. Fedorova, Phys. Lett. A 245 (1998) 555.
- [27] I. Lindgren, Molecular Physics 98 (2000) 1159.
- [28] C. Itzykson, J.-B. Zuber, Quantum Field Theory, McGraw-Hill, New York, 1980.
- [29] V.M. Shabaev, in : U.I. Safronova (ed.), Many-particles Effects in Atoms, AN SSSR, Nauchnyi Sovet po Spektroskopii, Moscow, 1988, p. 15.
- [30] V.M. Shabaev, in : U.I. Safronova (ed.), Many-particles Effects in Atoms, AN SSSR, Nauchnyi Sovet po Spektroskopii, Moscow, 1988, p. 24.
- [31] V.M. Shabaev, Izv. Vuz. Fiz. 33 (1990) 43 [Sov. Phys. Journ. 33 (1990) 660].
- [32] V.M. Shabaev, Teor. Mat. Fiz. 82 (1990) 83 [Theor. Math. Phys. 82 (1990) 57].
- [33] V.M. Shabaev, J. Phys. A 24 (1991) 5665.
- [34] V.V. Karasiov, L.N. Labzowsky, A.V. Nefiodov, V.M. Shabaev, Phys. Lett. A 161 (1992) 453.
- [35] V.M. Shabaev, J. Phys. B 26 (1993) 4703.
- [36] V.M. Shabaev, I.G. Fokeeva, Phys. Rev. A 49 (1994) 4489.
- [37] V.M. Shabaev, Phys. Rev. A 50 (1994) 4521.
- [38] M.B. Shabaeva, V.M. Shabaev, Phys. Rev. A 52 (1995) 2811.
- [39] V.A. Yerokhin, A.N. Artemyev, V.M. Shabaev, Phys. Lett. A 234 (1997) 361.
- [40] A.N. Artemyev, V.M. Shabaev, V.A. Yerokhin, Phys. Rev. A 56 (1997) 3529.
- [41] V.M. Shabaev, Phys. Rev. A 57 (1998) 59.
- [42] A.N. Artemyev, T. Beier, G. Plunien, V.M. Shabaev, G. Soff, V.A. Yerokhin, Phys. Rev. A 60 (1999) 45.
- [43] V.A. Yerokhin, A.N. Artemyev, T. Beier, G. Plunien, V.M. Shabaev, G. Soff, Phys. Rev. A 60 (1999) 3522.
- [44] A.N. Artemyev, T. Beier, G. Plunien, V.M. Shabaev, G. Soff, V.A. Yerokhin, Phys. Rev. A 62 (2000) 022116.
- [45] V.M. Shabaev, V.A. Yerokhin, T. Beier, J. Eichler, Phys. Rev. A 61 (2000) 052112.
- [46] V.A. Yerokhin, V.M. Shabaev, T. Beier, J. Eichler, Phys. Rev. A 62 (2000) 042712.
- [47] M.A. Braun, A.S. Shirokov, Izv. Akad. Nauk SSSR: Ser. Fiz. 41 (1977) 2585 [Bull. Acad. Sci. USSR: Phys. Ser. 41 (1977) 109].
- [48] M.A. Braun, Kh. Parera, Izv. Akad. Nauk SSSR: Ser. Fiz. 50 (1986) 1303 [Bull. Acad. Sci. USSR: Phys. Ser. 50 (1986) 59].
- [49] M.A. Braun, Teor. Mat. Fiz. 72 (1987) 394 [Theor. Math. Phys. 72 (1987) 958].
- [50] V.M. Shabaev, Teor. Mat. Fiz. 63 (1985) 394 [Theor. Math. Phys. 63 (1985) 588].
- [51] V.M. Shabaev, Yad. Fiz. 47 (1988) 107 [Sov. J. Nucl. Phys. 47 (1988) 69].
- [52] S.A. Zapryagaev, D.I. Morgulis, Yad. Fiz. 45 (1987) 716 [Sov. J. Nucl. Phys. 45 (1987) N 3].
- [53] G. Feldman, T. Fulton, Ann. Phys. (N.Y.) 179 (1987) 20.
- [54] T. Fulton, in: W.R. Johnson, P.J. Mohr, J. Sucher (eds.), Relativistic, Quantum Electrodynamics and Weak Interaction Effects in Atoms, American Institute of Physics Conference Series N 189, 1989, p. 429.
- [55] D.R. Yennie, in: W.R. Johnson, P.J. Mohr, J. Sucher (eds.), Relativistic, Quantum Electrodynamics and Weak Interaction Effects in Atoms, American Institute of Physics Conference Series N 189, 1989, p. 393.
- [56] J.D. Bjorken, D. Drell, Relativistic Quantum Fields, McGraw-Hill, New York, 1965.
- [57] P.J. Mohr, Phys. Rev. A 32 (1985) 1949.
- [58] G.S. Adkins, Phys. Rev. D 36, 1929 (1987).
- [59] G.S. Adkins, Phys. Rev. D 27, 1814 (1983).
- [60] V.L. Bonch-Bruевич, S.V. Tyablikov, The Green Function Method in Statistical Mechanics, North-Holland Publishing Company, Amsterdam, 1962.
- [61] D.J. Thouless, The Quantum Mechanics of Many-Body Systems, Academic Press, New York, 1961.
- [62] A.B. Migdal, Theory of Finite Fermi Systems and Properties of Atomic Nuclei, Nauka, Moscow, 1983.
- [63] A.A. Logunov, A.N. Tavkhelidze, Nuovo Cim. 29 (1963) 380.
- [64] R.N. Faustov, Teor. Mat. Fiz. 3 (1970) 240 [Theor. Math. Phys. 3 (1970) N 2].
- [65] B. Sz-Nagy, Comm. Math. Helv. 19 (1946/47) 347.
- [66] T. Kato, Progr. Theor. Phys. 4 (1949) 514.
- [67] T. Kato, Perturbation Theory for Linear Operators, Springer, Berlin, 1966.
- [68] A. Messiah, Quantum Mechanics, Vol. 2, Wiley, New York, 1962.

- [69] M. Reed, B. Simon, *Methods of Modern Mathematical Physics, Vol. 4: Analysis of Operators*, Academic Press, New York, 1978.
- [70] G.P. Lepage, *Phys. Rev. A* 16 (1977) 863.
- [71] S.A. Blundell, P.J. Mohr, W.R. Johnson, J. Sapirstein, *Phys. Rev. A* 48 (1993) 2615.
- [72] I.S. Gradshteyn, I.N. Ryzhik, *Tables of Integrals, Sums, Series, and Products*, Nauka, Moscow, 1977.
- [73] I. Lindgren, H. Persson, S. Salomonson, L.N. Labzowsky, *Phys. Rev. A* 51 (1995) 1167.
- [74] M.H. Mittleman, *Phys. Rev. A* 5 (1972) 2395.
- [75] E.-O. Le Bigot, P. Indelicato, V.M. Shabaev, *Phys. Rev. A* 63 (2001) 040501(R).
- [76] A.S. Yelkhovsky, Budker Institute of Nuclear Physics, Report No. BINP 94-27, Novosibirsk, 1994; arXiv: hep-th/9403095.
- [77] K. Pachucki, H. Grotch, *Phys. Rev. A* 51 (1995) 1854.
- [78] N.N. Bogolyubov, D.V. Shirkov, *Introduction to the Theory of Quantized Fields*, Nauka, Moscow, 1984.
- [79] V.B. Berestetsky, E.M. Lifshitz, L.P. Pitaevsky, *Quantum Electrodynamics*, Pergamon Press, Oxford, 1982.
- [80] P. Indelicato, V.M. Shabaev, to be published.
- [81] A.I. Akhiezer, V.B. Berestetsky, *Quantum Electrodynamics*, Nauka, Moscow, 1969.
- [82] J. Eichler, W. Meyerhof, *Relativistic Atomic Collisions*, Academic Press, San Diego, 1995.
- [83] F.E. Low, *Phys. Rev.* 88 (1952) 53.
- [84] M.A. Braun, *Zh. Eksp. Teor. Fiz.* 94 (1988) 145.
- [85] V.G. Gorshkov, L.N. Labzowsky, A.A. Sultanaev, *Zh. Eksp. Teor. Fiz.* 96 (1989) 53 [*Sov. Phys. JETP* 69 (1989) 28].
- [86] A.V. Nefiodov, V.V. Karasiov, V.A. Yerokhin, *Phys. Rev. A* 50 (1994) 4975.
- [87] E.H. Wichmann, N.M. Kroll, *Phys. Rev.* 101 (1956) 843.
- [88] N.L. Manakov, L.P. Rapoport, S.A. Zapryagaev, *Phys. Lett. A* 43 (1973) 139.
- [89] P.J. Mohr, *Ann. Phys.* 88 (1974) 26, 52.
- [90] M. Gyulassy, *Nucl. Phys. A* 244 (1975) 497.
- [91] G. Soff, P. Mohr, *Phys. Rev. A* 38 (1988) 5066.
- [92] G.W.F. Drake, S.P. Goldman, *Phys. Rev. A* 23 (1981) 2093.
- [93] I.P. Grant, *Phys. Rev. A* 25 (1982) 1230.
- [94] W.R. Johnson, S.A. Blundell, J. Sapirstein, *Phys. Rev. A* 37 (1988) 307.
- [95] S. Salomonson, P. Oster, *Phys. Rev. A* 40 (1989) 5548.
- [96] J. Sapirstein, W.R. Johnson, *J. Phys. B* 29 (1996) 5213.
- [97] I.P. Grant, H.M. Quiney, *Phys. Rev. A* 62 (2000) 022508.
- [98] V.M. Shabaev, *J. Phys. B* 24 (1991) 4479.
- [99] M. Baranger, H.A. Bethe, and R.P. Feynman, *Phys. Rev.* 92 (1953) 482.
- [100] N.J. Snyderman, *Ann. Phys.* 211 (1991) 43.
- [101] V.A. Yerokhin, V.M. Shabaev, *Phys. Rev. A* 60 (1999) 800.
- [102] G.A. Rinker and L. Wilets, *Phys. Rev. A* 12 (1975) 748.
- [103] H. Persson, S. Salomonson, P. Sunnergren, and I. Lindgren, *Phys. Rev. Lett.* 76 (1996) 204.
- [104] P. Sunnergren, Ph. D. thesis, Göteborg University and Chalmers University of Technology, Göteborg, 1998.
- [105] H. Persson, S.M. Schneider, G. Soff, W. Greiner, I. Lindgren, *Phys. Rev. Lett.* 76 (1996) 1433.
- [106] S.A. Blundell, K.T. Cheng, J. Sapirstein, *Phys. Rev. A* 55 (1997) 1857.
- [107] V.M. Shabaev, M. Tomaselli, T. Kühl, A.N. Artemyev, V.A. Yerokhin, *Phys. Rev. A* 56 (1997) 252.
- [108] H. Persson, S. Salomonson, P. Sunnergren, I. Lindgren, *Phys. Rev. A* 56 (1997) R2499.
- [109] V.M. Shabaev, M.B. Shabaeva, I.I. Tupitsyn, V.A. Yerokhin, A.N. Artemyev, T. Kühl, M. Tomaselli, O.M. Zharebtsov, *Phys. Rev. A* 57, (1998) 149; 58 (1998) 1610.
- [110] P. Sunnergren, H. Persson, S. Salomonson, S. M. Schneider, I. Lindgren, G. Soff, *Phys. Rev. A* 58 (1998) 1055.
- [111] T. Beier, I. Lindgren, H. Persson, S. Salomonson, P. Sunnergren, H. Häffner, N. Hermanspahn, *Phys. Rev. A* 62 (2000) 032510.
- [112] T. Franosch, G. Soff, *Z. Phys. D* 18 (1991) 219.
- [113] D. Andrae, *Phys. Rep.* 336 (2000) 413.
- [114] V.M. Shabaev, *J. Phys. B* 26 (1993) 1103.
- [115] A.M. Desiderio, W.R. Johnson, *Phys. Rev. A* 3 (1971) 1287.
- [116] G.E. Brown, J.S. Langer, G.W. Schaefer, *Proc. R. Soc. London, Ser. A* 251 (1959) 92.
- [117] S.A. Blundell, N.J. Snyderman, *Phys. Rev. A* 44 (1991) R1427.
- [118] P. Indelicato, P.J. Mohr, *Phys. Rev. A* 46 (1992) 172.
- [119] H. Persson, I. Lindgren, S. Salomonson, *Phys. Scr. T* 46 (1993) 125.
- [120] H.M. Quiney, I.P. Grant, *Phys. Scr. T* 46 (1993) 132; *J. Phys. B* 27 (1994) L299.
- [121] P.J. Mohr, *Phys. Rev. A* 46 (1992) 4421.
- [122] P. Indelicato, P.J. Mohr, *Phys. Rev. A* 58 (1998) 165.
- [123] P.J. Mohr, G. Soff, *Phys. Rev. Lett* 70 (1993) 158.
- [124] T. Beier, P.J. Mohr, H. Persson, G. Soff, *Phys. Rev. A* 58 (1998) 954.
- [125] U.D. Jentschura, P.J. Mohr, G. Soff, *Phys. Rev. Lett.* 82 (1999) 53.

- [126] U.D. Jentschura, P.J. Mohr, G. Soff, Phys. Rev. A 63 (2001) 042512.
- [127] N.L. Manakov, A.A. Nekipelov, A.G. Fainstein, Sov. Phys. JETP 68 (1989) 673 [Zh. Eksp. Teor. Fiz. 95 (1989) 1167].
- [128] T. Beier, G. Plunien, M. Greiner, G. Soff, J. Phys. B 30 (1997) 2761.
- [129] H. Persson, I. Lindgren, S. Salomonson, P. Sunnergren, Phys. Rev. A 48 (1993) 2772.
- [130] T. Beier, P.J. Mohr, H. Persson, G. Plunien, M. Greiner, G. Soff, Phys. Lett. A 236 (1997) 329.
- [131] A. Mitrushenkov, L.N. Labzowsky, I. Lindgren, H. Persson, S. Salomonson, Phys. Lett. A 200 (1995) 51.
- [132] S. Mallampalli, J. Sapirstein, Phys. Rev. A 57 (1998) 1548.
- [133] S. Mallampalli, J. Sapirstein, Phys. Rev. Lett. 80 (1998) 5297.
- [134] V.A. Yerokhin, Phys. Rev. A 62 (2000) 012508.
- [135] V.A. Yerokhin, Phys. Rev. Lett. 86 (2001) 1990.
- [136] I. Goidenko, L. Labzowsky, A. Nefiodov, G. Plunien, G. Soff, S. Zschocke, Hyperfine Interactions 127 (2000) 293.
- [137] K. Pachucki, Hyperfine Interactions 114 (1998) 55.
- [138] K. Pachucki, Phys. Rev. A 63 (2001) 042503.
- [139] V.M. Artemyev, V.M. Shabaev, V.A. Yerokhin, Phys. Rev. A 52 (1995) 1884.
- [140] V.M. Artemyev, V.M. Shabaev, V.A. Yerokhin, J. Phys. B 28 (1995) 5201.
- [141] V.M. Shabaev, A.N. Artemyev, T. Beier, G. Plunien, V.A. Yerokhin, G. Soff, Phys. Rev. A 57 (1998) 4235.
- [142] V.M. Shabaev, A.N. Artemyev, T. Beier, G. Plunien, V.A. Yerokhin, G. Soff, Phys. Scr. T 80 (1999) 493.
- [143] V.M. Shabaev, A.N. Artemyev, T. Beier, G. Soff, J. Phys. B 31 (1998) L337.
- [144] G. Plunien, G. Soff, Phys. Rev. A 51 (1995) 1119; 53 (1996) 4614.
- [145] A.V. Nefiodov, L.N. Labzowsky, G. Plunien, G. Soff, Phys. Lett. A 222 (1996) 227.
- [146] P.J. Mohr, B.N. Taylor, Rev. Mod. Phys. 72 (2000) 351.
- [147] H.F. Beyer, IEEE Trans. Instrum. Meas. 44 (1995) 510.
- [148] H.F. Beyer, G. Menzel, D. Liesen, A. Gallus, F. Bosch, R. Deslattes, P. Indelicato, T. Stöhlker, O. Klepper, R. Moshhammer, F. Nolden, H. Eickhoff, B. Franzke, M. Steck, Z. Phys. D 35 (1995) 169.
- [149] T. Stöhlker, P.H. Mokler, F. Bosch, R.W. Dunford, O. Klepper, C. Kozhuharov, T. Ludziejewski, F. Franzke, F. Nolden, H. Reich, P. Rymuza, Z. Stachura, M. Steck, P. Swiat, A. Warczak, Phys. Rev. Lett. 85 (2000) 3109.
- [150] G.W. Drake, Can. J. Phys. 66 (1988) 586.
- [151] P.J. Mohr, J. Sapirstein, Phys. Rev. A 62 (2000) 052501.
- [152] R.E. Marrs, S.R. Elliot, T. Stöhlker, Phys. Rev. A 52 (1995) 3577.
- [153] F.C. Sanders, C.W. Scherr, Phys. Rev. 181 (1969) 84.
- [154] J. Schweppe, A. Belkacem, L. Blumenfeld, N. Claytor, B. Feinberg, H. Gould, V. E. Kostroun, L. Levy, S. Misawa, J. R. Mowat, M. H. Prior, Phys. Rev. Lett. 66 (1991) 1434 .
- [155] P. Beiersdorfer, A.L. Osterheld, J.H. Scofield, J.R. Crespo-López-Urrutia, K. Widmann, Phys. Rev. Lett. 80 (1998) 3022.
- [156] U. Staude, Ph. Bosselman, R. Büttner, D. Horn, K.-H. Schartner, F. Folkmann, A.E. Livingston, T. Ludziejewski, P.H. Mokler, Phys. Rev. A 58 (1998) 3516.
- [157] Ph. Bosselmann, U. Staude, D. Horn, K.-H. Schartner, F. Folkmann, A. E. Livingston, P.H. Mokler, Phys. Rev. A 59 (1999) 1874.
- [158] V.A. Yerokhin, A.N. Artemyev, V.M. Shabaev, M.M. Sysak, O.M. Zherebtsov, G. Soff, Phys. Rev. Lett. 85 (2000) 4699.
- [159] P. Indelicato, P.J. Mohr, Theor. Chem. Acta 80 (1991) 207.
- [160] K.T. Cheng, W.R. Johnson, J. Sapirstein, Phys. Rev. Lett. 66 (1991) 2960.
- [161] P.J. Mohr, Phys. Scr. T 46 (1993) 44.
- [162] S.A. Blundell, Phys. Rev. A 47 (1993) 1790.
- [163] I. Lindgren, H. Persson, S. Salomonson, A. Ynnerman, Phys. Rev. A 47 (1993) R4555.
- [164] M.H. Chen, K.T. Cheng, W.R. Johnson, J. Sapirstein, Phys. Rev. A 52 (1995) 266.
- [165] V.M. Shabaev, V.A. Yerokhin, O.M. Zherebtsov, A.N. Artemyev, M.M. Sysak, G. Soff, Hyperfine Interactions 132 (2001) 339.
- [166] V.M. Shabaev, J. Phys. B 27 (1994) 5825.
- [167] V.M. Shabaev, In H.F. Beyer, V.P. Shevelko (eds.), Atomic Physics with Heavy Ions, Springer, 1999, p. 139.
- [168] I. Klaft, S. Borneis, T. Engel, B. Fricke, R. Grieser, G. Huber, T. Kühl, D. Marx, R. Neumann, S. Schröder, P. Seelig, L. Völker, Phys. Rev. Lett. 73 (1994) 2425.
- [169] J.R. Crespo Lopez-Urrutia, P. Beiersdorfer, D. Savin, K. Widmann, Phys. Rev. Lett. 77 (1996) 826.
- [170] J.R. Crespo Lopez-Urrutia, P. Beiersdorfer, K. Widmann, B. Birkel, A.-M. Mårtensson-Pendrill, M.G.H. Gustavsson, Phys. Rev. A 57 (1998) 879.
- [171] P. Seelig, S. Borneis, A. Dax, T. Engel, S. Faber, M. Gerlach, C. Holbrow, G. Huber, T. Kühl, D. Marx, K. Meier, P. Merz, W. Quint, F. Schmitt, M. Tomaselli, L. Völker, M. Würtz, K. Beckert, B. Franzke, F. Nolden, H. Reich, M. Steck, T. Winkler, Phys. Rev. Lett. 81 (1998) 4824.
- [172] P. Raghavan, At. Data Nucl. Data Tables 42, 189 (1989).
- [173] M.G.H. Gustavsson, A.-M. Mårtensson-Pendrill, Phys. Rev. A, 58 (1998) 3611.
- [174] O. Lutz, G. Stricker, Phys. Lett. A 35 (1971) 397.
- [175] H.M. Gibbs, C.M. White, Phys. Rev. A 188 (1969) 180.

- [176] O.P. Sushkov, V.V. Flambaum, I.B. Khriplovich, *Opt. Spectr.* 44 (1978) 2.
- [177] V.M. Shabaev, M.B. Shabaeva, I.I. Tupitsyn, V.A. Yerokhin, *Hyperfine Interactions* **114**, 129 (1998).
- [178] M.B. Shabaeva, *Opt. Spectr.* 86 (1999) 368.
- [179] V.M. Shabaev, A.N. Artemyev, O.M. Zherebtsov, V.A. Yerokhin, G. Plunien, G. Soff, *Hyperfine Interactions* 27 (2000) 279.
- [180] S. Boucard, P. Indelicato, *Eur. Phys. J. D* 8 (2000) 59.
- [181] O.M. Zherebtsov, V.M. Shabaev, *Can. J. Phys.* 78 (2000) 701.
- [182] A.N. Artemyev, V.M. Shabaev, G. Plunien, G. Soff, V.A. Yerokhin, *Phys. Rev. A*, in press.
- [183] J. Sapirstein, K.T. Cheng, *Phys. Rev. A* 63 (2001) 032506.
- [184] G. Breit, *Nature (London)* 122 (1928) 649.
- [185] S.G. Karshenboim, *Phys. Lett. A* 266 (2000) 380.
- [186] W. Quint, *Phys. Scr.* 59 (1995) 203.
- [187] N. Hermanspahn, W. Quint, S. Stahl, M. Tönges, G. Bollen, H.-J. Kluge, R. Ley, R. Mann, G. Werth, *Hyperfine Interactions* 99 (1996) 91.
- [188] N. Hermanspahn, H. Häffner, H.-J. Kluge, W. Quint, S. Stahl, J. Verdu, and G. Werth, *Phys. Rev. Lett.* 84 (2000) 427.
- [189] H. Häffner, T. Beier, N. Hermanspahn, H.-J. Kluge, W. Quint, S. Stahl, J. Verdu, G. Werth, *Phys. Rev. Lett.* 85 (2000) 5308.
- [190] S.G. Karshenboim, in: S. Karshenboim et al. (eds.), *The Hydrogen Atom*, Berlin, Springer, 2001, p.651; arXiv: hep-ph/0008227 (2000) (<http://xxx.lanl.gov>).
- [191] V.M. Shabaev, *Can. J. Phys.* 76 (1998) 907.
- [192] H. Winter, S. Borneis, A. Dax, S. Faber, T. Kühn, D. Marx, F. Schmitt, P. Seelig, W. Seelig, V.M. Shabaev, M. Tomaselli, M. Würtz, in *GSI scientific report 1998*, edited by U. Grundinger (GSI, Darmstadt, Germany, 1999), p. 87.
- [193] V.G. Pal'chikov, *Hyperfine Interactions* 127 (2000) 287.
- [194] A. Ichihara, T. Shirai, J. Eichler, *Phys. Rev. A* 49 (1994) 1875.
- [195] J. Eichler, A. Ichihara, T. Shirai, *Phys. Rev. A* 51 (1995) 3027.
- [196] Th. Stöhlker, C. Kozhuharov, P.H. Mokler, A. Warczak, F. Bosch, H. Geissel, R. Moshhammer, C. Scheidenberger, J. Eichler, A. Ichihara, T. Shirai, Z. Stachura, P. Rymuza, *Phys. Rev. A* 51 (1995) 2098.
- [197] Th. Stöhlker, T. Ludziejewski, F. Bosch, R.W. Dunford, C. Kozhuharov, P.H. Mokler, H.F. Beyer, O. Brinzaescu, F. Franzke, J. Eichler, A. Griegal, S. Hagmann, A. Ichihara, A. Krämer, D. Liesen, H. Reich, P. Rymuza, Z. Stachura, M. Steck, P. Swiat, A. Warczak, *Phys. Rev. Lett.* 82 (1999) 3232.
- [198] J.M. Jauch, F. Rohrlich, *The Theory of Photons and Electrons*, Springer-Verlag, Berlin, 1976.
- [199] D.R. Yennie, S.C. Frautschi, H. Suura, *Ann. Phys.* 13 (1961) 379.
- [200] V.A. Yerokhin, Diploma thesis, St. Petersburg State University, St. Petersburg, 1993.
- [201] A.S. Shirokov, Candidate of Science thesis, Leningrad State University, Leningrad, 1981.
- [202] L. Van Hove, *Physica* 21 (1955) 901.
- [203] N.M. Hugenholtz, *Physica* 23 (1957) 481.
- [204] H.A. Bethe, *Intermediate Quantum Mechanics*, Benjamin, New York, 1964.

TABLE I. The ground-state Lamb shift in $^{238}\text{U}^{91+}$, in eV.

Point nucleus binding energy	-132279.92(1)
Finite nuclear size [39,112]	198.81(38)
First-order SE [123]	355.05
First-order VP [119]	-88.60
Second-order QED	± 1.5
Nuclear recoil [141]	0.46
Nuclear polarization [144,145]	-0.20(10)
Lamb shift theory	$465.52(39) \pm 1.5$
Lamb shift experiment [149]	468(13)

TABLE II. The two-electron contribution to the ground-state energy in $^{209}\text{Bi}^{81+}$, in eV.

One-photon exchange contribution	1897.56(1)
Two-photon exchange within the Breit approximation	-10.64
Two-photon exchange beyond the Breit approximation	-0.30(1)
SE screening	-6.73
VP screening	1.55
Three- and more photon contribution	0.06(7)
Total theory [39]	1881.50(7)
Experiment [152]	1876(14)

TABLE III. The $2p_{1/2} - 2s$ transition energy in $^{238}\text{U}^{89+}$, in eV.

One-photon exchange [43]	368.83
One-electron nuclear size [43]	-33.35(6)
First-order QED [119,123]	-42.93
Two-photon exchange within the Breit approximation [158]	-13.54
Two-photon exchange beyond the Breit approximation [158]	0.17
Self energy screening [43]	1.52
Vacuum polarization screening [42]	-0.36
Three- and more photon exchange [165]	0.16(7)
Nuclear recoil [139]	-0.07
Nuclear polarization [144,145]	0.03(1)
One-electron second-order QED	± 0.20
Total theory	$280.46(9) \pm 0.20$
Experiment [154]	280.59(10)

TABLE IV. Theoretical contributions to the ground state hyperfine splitting in hydrogenlike ions (in eV).

Ion	$^{165}\text{Ho}^{66+}$	$^{185}\text{Re}^{74+}$	$^{187}\text{Re}^{74+}$	$^{207}\text{Pb}^{81+}$	$^{209}\text{Bi}^{82+}$
μ/μ_N	4.177(5)	3.1871(3)	3.2197(5)	0.592583(9)	4.1106(2)
Relativistic value for point nucleus	2.326(3)	3.010	3.041	1.425	5.839
Finite nuclear charge	-0.106(1)	-0.213(2)	-0.215(2)	-0.149	-0.649(2)
Bohr-Weisskopf effect	-0.020(6)	-0.034(10)	-0.035(10)	-0.053(5)	-0.061(27)
QED	-0.011	-0.015	-0.015	-0.007	-0.030
Total theory [179]	2.189(7)	2.748(10)	2.776(10)	1.215(5)	5.100(27)
Experiment [168–171]	2.1645(5)	2.719(2)	2.745(2)	1.2159(2)	5.0840(8)

TABLE V. The bound-electron g factor in $^{209}\text{Bi}^{82+}$.

Relativistic value	1.7276
QED	0.0029
Nuclear size correction	0.0005
Total theory [179]	1.7310
Experiment [192]	1.7343(33)

TABLE VI. The relative values of the QED corrections to the total cross section for the radiative recombination into the K-shell of bare uranium [45], expressed in %.

Impact energy [MeV/u]	Correction	Vacuum polarization, in %	Self energy, in %
100	$\sigma_{\text{en}}^{(1)} + \sigma_{\text{bw}}^{(1)}$	0.126	-0.390
	$\sigma_{\text{cw}}^{(1)}$	-0.006	?
	Total	0.120	-0.390
300	$\sigma_{\text{en}}^{(1)} + \sigma_{\text{bw}}^{(1)}$	0.175	-0.513
	$\sigma_{\text{cw}}^{(1)}$	-0.003	?
	Total	0.173	-0.513
1000	$\sigma_{\text{en}}^{(1)} + \sigma_{\text{bw}}^{(1)}$	0.220	-0.591
	$\sigma_{\text{cw}}^{(1)}$	0.043	?
	Total	0.263	-0.591

TABLE VII. The zeroth-order cross section $\sigma^{(0)}$ and the first-order interelectronic-interaction correction calculated in Ref. [46], in barns. $\sigma_{\text{scr}}^{(1)}$ denotes the interelectronic-interaction correction calculated in the screening potential approximation and $\sigma_{\text{int}}^{(1)}$ indicates the results of the rigorous relativistic calculation.

Impact energy [MeV/u]	$\sigma^{(0)}$	$\sigma_{\text{scr}}^{(1)}$	$\sigma_{\text{int}}^{(1)}$
2s-state:			
100	41.203	-1.393	-2.055
300	9.105	-0.3345	-0.3755
700	2.457	-0.0979	-0.1051
2p _{1/2} -state:			
100	33.041	-2.535	-3.088
300	5.042	-0.4538	-0.3864
700	1.065	-0.1022	-0.0861
2p _{3/2} -state:			
100	31.489	-2.275	-2.896
300	3.646	-0.3132	-0.2804
700	0.622	-0.0568	-0.0489




FIG. 1. Singularities of the two-time Green function in the complex E plane.




FIG. 2. Singularities of the two-time Green function in the bound-state region, if the interaction between the electron-positron field and the electromagnetic field is switched off.




FIG. 3. Singularities of the two-time Green function in the bound-state region, disregarding the instability of excited states.




FIG. 4. Singularities of the two-time Green function in the bound-state region if the cuts are turned down, to the second sheet of the Riemann surface. The instability of excited states is disregarded.




FIG. 5. Singularities of the two-time Green function in the bound-state region if the cuts are turned down, to the second sheet of the Riemann surface. The instability of excited states is taken into account.




FIG. 6. Singularities of the two-time Green function in the bound state region for a non-zero photon mass, including one- and two-photon spectra, if the cuts are turned down, to the second sheet of the Riemann surface. The instability of excited states is disregarded.




FIG. 7. The contour Γ surrounds the pole corresponding to the level under consideration and keeps outside all other singularities. For simplicity, only one- and two-photon spectra are displayed.




FIG. 8. Singularities of the photon propagator in the complex ω plane for a non-zero photon mass μ .



FIG. 9. The contour Γ surrounds the poles corresponding to the quasidegenerate levels under consideration and keeps outside all other singularities. For simplicity, only one-photon spectra are displayed.




FIG. 10. A deformation of the contour Γ that allows drawing the cuts to the related poles in the case of quasidegenerate states when $\mu \rightarrow 0$. For simplicity, only one-photon spectra are displayed.




FIG. 11. The interaction with an external potential $\delta V(\mathbf{x})$.




FIG. 12. The first-order self-energy diagram.




FIG. 13. The first-order vacuum-polarization diagram.



FIG. 14. The mass-counterterm diagram.




FIG. 15. δV - self-energy diagrams.




FIG. 16. δV - mass-counterterm diagrams.




FIG. 17. C is the contour of the integration over the electron energy ω in the formalism with the standard vacuum. C' is the integration contour for the vacuum with the $(1s)^2$ shell included. The integral along the contour $C_{\text{int}} = C' - C$ describes the interaction of the valence electron with the $(1s)^2$ -shell electrons.




FIG. 18. The first-order self-energy diagrams for a two-electron atom.




FIG. 19. The first-order vacuum-polarization diagrams for a two-electron atom.




FIG. 20. One-photon exchange diagram.




FIG. 21. Two-photon exchange diagrams.




FIG. 22. The self-energy screening diagrams.




FIG. 23. The vacuum-polarization screening diagrams.




FIG. 24. δV - interelectronic-interaction diagrams.




FIG. 25. Coulomb nuclear recoil diagram.



FIG. 26. One-transverse-photon nuclear recoil diagrams.




FIG. 27. Two-transverse-photon nuclear recoil diagram.




FIG. 28. Two-electron Coulomb nuclear recoil diagram.




FIG. 29. Two-electron one-transverse-photon nuclear recoil diagrams.




FIG. 30. Two-electron two-transverse-photon nuclear recoil diagram.




FIG. 31. The photon emission by a one-electron atom in zeroth-order approximation.




FIG. 32. The first-order QED corrections to the photon emission by a one-electron atom.




FIG. 33. The mass-counterterm corrections to the photon emission by a one-electron atom.




FIG. 34. The radiative recombination of an electron with a hydrogenlike atom in zeroth-order approximation.




FIG. 35. The interelectronic-interaction corrections of first order in $1/Z$ to the radiative recombination of an electron with a hydrogenlike atom.



FIG. 36. The photon scattering on a one-electron atom in zeroth-order approximation.




FIG. 37. A typical diagram describing two-photon exchange between three electrons in a lithiumlike atom.




FIG. 38. The radiative recombination accompanied by emission of a soft photon.




FIG. 39. A time-ordered version of a Feynman diagram.